

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dichloridobis(5,7-dichloroquinolin-8-olato- κ^2N,O)tin(IV)

Yousef Fazaeli,^a Ezzatollah Najafi,^a Mostafa M. Amini^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 5 February 2009; accepted 6 February 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.026; wR factor = 0.079; data-to-parameter ratio = 17.8.

The Sn^{IV} atom in the title compound, [Sn(C₉H₄Cl₂NO)₂Cl₂], is chelated by the substituted quinolin-8-olate anions in a distorted octahedral geometry. The N-donor atoms are in a *cis* alignment as are the Cl atoms; the O atoms are *trans* to each other.

Related literature

For the structure of dichloridobis(quinolin-8-olato)tin(IV), which shows a very similar coordination geometry, see: Archer *et al.* (1987).



Experimental

Crystal data

[Sn(C₉H₄Cl₂NO)₂Cl₂]
 $M_r = 615.65$
 Monoclinic, $P2_1/c$
 $a = 15.2459$ (2) Å
 $b = 8.9262$ (1) Å
 $c = 15.8541$ (2) Å
 $\beta = 110.381$ (1)°

$V = 2022.48$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.08$ mm⁻¹
 $T = 100$ (2) K
 $0.28 \times 0.22 \times 0.18$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.594$, $T_{\max} = 0.706$

18582 measured reflections
 4651 independent reflections
 4218 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.079$
 $S = 1.06$
 4651 reflections

262 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.59$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.25$ e Å⁻³

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

The authors thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2866).

References

- Archer, S. J., Koch, K. R. & Schmidt, S. (1987). *Inorg. Chim. Acta*, **126**, 209–218.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). publCIF. In preparation.

supporting information

Acta Cryst. (2009). E65, m270 [doi:10.1107/S1600536809004371]

Dichloridobis(5,7-dichloroquinolin-8-olato- κ^2N,O)tin(IV)

Yousef Fazaeli, Ezzatollah Najafi, Mostafa M. Amini and Seik Weng Ng

S1. Experimental

5,7-Dichloro-8-hydroxyquinoline (1 mmol, 0.21 g) was added to a solution of stannous chloride (1 mmol, 0.23) in DMSO (20 ml). The clear solution was set aside for several days to yield yellow crystals. These crystals are stable when heated at 573 K.

S2. Refinement

H-atoms were placed in calculated positions (C—H 0.93 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. The final difference Fourier map had a large peak/deep hole at about 1 Å from the Sn atom.




Figure 1

Anisotropic displacement ellipsoid plot (Barbour, 2001) of $\text{SnCl}_2(\text{C}_9\text{H}_4\text{Cl}_2\text{NO})_2$; ellipsoids are drawn at the 70% probability level and H atoms of arbitrary radius.

Dichloridobis(5,7-dichloroquinolin-8-olato- κ^2N,O)tin(IV)

Crystal data

[Sn(C₉H₄Cl₂NO)₂Cl₂]
M_r = 615.65
 Monoclinic, *P*2₁/*c*
 Hall symbol: -P 2ybc
a = 15.2459 (2) Å
b = 8.9262 (1) Å
c = 15.8541 (2) Å
 β = 110.381 (1)°
V = 2022.48 (4) Å³
Z = 4

F(000) = 1192
D_x = 2.022 Mg m⁻³
 Mo *K* α radiation, λ = 0.71073 Å
 Cell parameters from 9940 reflections
 θ = 2.6–28.3°
 μ = 2.08 mm⁻¹
T = 100 K
 Polyhedron, yellow
 0.28 × 0.22 × 0.18 mm

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
T_{min} = 0.594, *T_{max}* = 0.706

18582 measured reflections
 4651 independent reflections
 4218 reflections with *I* > 2 σ (*I*)
R_{int} = 0.021
 θ_{\max} = 27.5°, θ_{\min} = 1.4°
h = -19→19
k = -11→11
l = -20→20

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2 σ (*F*²)] = 0.026
wR(*F*²) = 0.079
S = 1.06
 4651 reflections
 262 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
w = 1/[$\sigma^2(F_o^2) + (0.0394P)^2 + 4.6706P$]
 where *P* = (*F_o*² + 2*F_c*²)/3
 (Δ/σ)_{max} = 0.001
 $\Delta\rho_{\max}$ = 0.59 e Å⁻³
 $\Delta\rho_{\min}$ = -1.25 e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> */ <i>U_{eq}</i>
Sn1	0.778941 (13)	0.92337 (2)	0.344613 (13)	0.01736 (7)
Cl1	0.93987 (6)	0.32043 (9)	0.64058 (5)	0.02934 (17)
Cl2	0.84159 (6)	0.38698 (8)	0.28352 (5)	0.02702 (17)
Cl3	0.34831 (5)	1.07293 (11)	0.39582 (6)	0.0343 (2)
Cl4	0.66222 (6)	1.38854 (9)	0.45378 (7)	0.03409 (19)
Cl5	0.70590 (6)	0.98399 (10)	0.19137 (5)	0.02889 (17)
Cl6	0.92679 (6)	1.01709 (10)	0.36724 (6)	0.03319 (19)
O1	0.80174 (15)	0.7039 (2)	0.31888 (13)	0.0191 (4)
O2	0.73412 (14)	1.1077 (2)	0.39671 (15)	0.0205 (4)
N1	0.83134 (16)	0.8252 (3)	0.48199 (16)	0.0171 (5)
N2	0.63835 (17)	0.8521 (3)	0.34618 (16)	0.0192 (5)
C1	0.83654 (19)	0.6170 (3)	0.39116 (19)	0.0169 (5)
C2	0.8582 (2)	0.4680 (3)	0.38683 (19)	0.0183 (6)

C3	0.8920 (2)	0.3773 (3)	0.4644 (2)	0.0198 (6)
H3	0.9064	0.2750	0.4591	0.024*
C4	0.9038 (2)	0.4366 (3)	0.5468 (2)	0.0198 (6)
C5	0.8855 (2)	0.5884 (3)	0.55734 (19)	0.0180 (6)
C6	0.85181 (19)	0.6765 (3)	0.47856 (19)	0.0160 (5)
C7	0.8970 (2)	0.6608 (4)	0.64017 (19)	0.0208 (6)
H7	0.9187	0.6058	0.6949	0.025*
C8	0.8765 (2)	0.8106 (4)	0.6412 (2)	0.0230 (6)
H8	0.8843	0.8598	0.6964	0.028*
C9	0.8441 (2)	0.8900 (3)	0.5600 (2)	0.0202 (6)
H9	0.8310	0.9939	0.5612	0.024*
C10	0.6471 (2)	1.1032 (3)	0.39562 (19)	0.0183 (6)
C11	0.6027 (2)	1.2221 (3)	0.4198 (2)	0.0217 (6)
C12	0.5108 (2)	1.2119 (4)	0.4191 (2)	0.0248 (6)
H12	0.4824	1.2963	0.4356	0.030*
C13	0.4617 (2)	1.0813 (4)	0.3950 (2)	0.0250 (7)
C14	0.5014 (2)	0.9536 (4)	0.37007 (19)	0.0218 (6)
C15	0.5937 (2)	0.9681 (3)	0.37025 (19)	0.0185 (5)
C16	0.4580 (2)	0.8118 (4)	0.3460 (2)	0.0288 (7)
H16	0.3959	0.7965	0.3449	0.035*
C17	0.5055 (2)	0.6976 (4)	0.3246 (2)	0.0299 (7)
H17	0.4771	0.6018	0.3099	0.036*
C18	0.5961 (2)	0.7209 (4)	0.3242 (2)	0.0251 (6)
H18	0.6280	0.6409	0.3077	0.030*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.01867 (11)	0.01519 (11)	0.01871 (11)	0.00298 (7)	0.00712 (8)	0.00097 (7)
C11	0.0364 (4)	0.0257 (4)	0.0232 (4)	0.0040 (3)	0.0070 (3)	0.0099 (3)
C12	0.0428 (4)	0.0173 (3)	0.0204 (3)	0.0031 (3)	0.0103 (3)	-0.0020 (3)
C13	0.0169 (3)	0.0521 (5)	0.0357 (4)	0.0022 (3)	0.0114 (3)	0.0054 (4)
C14	0.0308 (4)	0.0211 (4)	0.0559 (5)	-0.0008 (3)	0.0220 (4)	-0.0073 (4)
C15	0.0293 (4)	0.0319 (4)	0.0251 (4)	0.0071 (3)	0.0091 (3)	0.0065 (3)
C16	0.0314 (4)	0.0304 (4)	0.0410 (5)	-0.0019 (3)	0.0167 (4)	-0.0024 (4)
O1	0.0251 (10)	0.0166 (10)	0.0150 (9)	0.0032 (8)	0.0063 (8)	0.0006 (8)
O2	0.0167 (10)	0.0175 (10)	0.0293 (11)	-0.0003 (8)	0.0107 (9)	-0.0020 (8)
N1	0.0153 (11)	0.0178 (12)	0.0189 (11)	0.0009 (9)	0.0068 (9)	-0.0013 (9)
N2	0.0204 (12)	0.0201 (12)	0.0162 (11)	-0.0008 (10)	0.0052 (9)	0.0016 (9)
C1	0.0157 (13)	0.0172 (13)	0.0184 (13)	-0.0007 (10)	0.0066 (10)	0.0006 (10)
C2	0.0206 (14)	0.0166 (13)	0.0186 (13)	0.0006 (11)	0.0078 (11)	-0.0016 (11)
C3	0.0189 (14)	0.0174 (13)	0.0232 (14)	0.0001 (11)	0.0074 (11)	0.0014 (11)
C4	0.0168 (13)	0.0226 (15)	0.0186 (13)	-0.0022 (11)	0.0046 (11)	0.0058 (11)
C5	0.0147 (13)	0.0237 (15)	0.0153 (13)	-0.0010 (11)	0.0047 (10)	0.0021 (11)
C6	0.0132 (12)	0.0164 (13)	0.0193 (13)	-0.0006 (10)	0.0069 (10)	-0.0005 (10)
C7	0.0182 (13)	0.0281 (16)	0.0167 (13)	-0.0008 (12)	0.0067 (11)	0.0017 (12)
C8	0.0203 (14)	0.0306 (17)	0.0191 (14)	-0.0005 (12)	0.0083 (11)	-0.0041 (12)
C9	0.0178 (13)	0.0207 (14)	0.0234 (14)	-0.0012 (11)	0.0089 (11)	-0.0046 (11)

C10	0.0181 (13)	0.0205 (14)	0.0173 (13)	0.0009 (11)	0.0074 (11)	0.0025 (11)
C11	0.0217 (14)	0.0201 (14)	0.0245 (14)	0.0027 (12)	0.0094 (12)	0.0016 (11)
C12	0.0228 (15)	0.0299 (17)	0.0233 (15)	0.0079 (13)	0.0099 (12)	0.0042 (12)
C13	0.0146 (13)	0.0381 (18)	0.0224 (15)	0.0029 (12)	0.0067 (11)	0.0061 (13)
C14	0.0189 (14)	0.0290 (16)	0.0159 (13)	-0.0013 (12)	0.0041 (11)	0.0039 (12)
C15	0.0176 (13)	0.0221 (14)	0.0157 (12)	0.0004 (11)	0.0056 (10)	0.0034 (11)
C16	0.0214 (15)	0.041 (2)	0.0229 (15)	-0.0091 (14)	0.0058 (12)	0.0010 (14)
C17	0.0296 (17)	0.0285 (17)	0.0276 (16)	-0.0131 (14)	0.0052 (13)	-0.0049 (13)
C18	0.0288 (16)	0.0243 (16)	0.0208 (14)	-0.0022 (13)	0.0070 (12)	-0.0010 (12)

Geometric parameters (Å, °)

Sn1—O1	2.055 (2)	C4—C5	1.405 (4)
Sn1—O2	2.061 (2)	C5—C6	1.413 (4)
Sn1—N1	2.222 (2)	C5—C7	1.419 (4)
Sn1—N2	2.244 (2)	C7—C8	1.374 (5)
Sn1—Cl6	2.3122 (9)	C7—H7	0.9500
Sn1—Cl5	2.3561 (8)	C8—C9	1.400 (4)
Cl1—C4	1.737 (3)	C8—H8	0.9500
Cl2—C2	1.726 (3)	C9—H9	0.9500
Cl3—C13	1.735 (3)	C10—C11	1.383 (4)
Cl4—C11	1.726 (3)	C10—C15	1.432 (4)
O1—C1	1.332 (3)	C11—C12	1.401 (4)
O2—C10	1.321 (4)	C12—C13	1.366 (5)
N1—C9	1.316 (4)	C12—H12	0.9500
N1—C6	1.369 (4)	C13—C14	1.411 (5)
N2—C18	1.323 (4)	C14—C15	1.412 (4)
N2—C15	1.365 (4)	C14—C16	1.418 (5)
C1—C2	1.378 (4)	C16—C17	1.361 (5)
C1—C6	1.425 (4)	C16—H16	0.9500
C2—C3	1.411 (4)	C17—C18	1.400 (5)
C3—C4	1.362 (4)	C17—H17	0.9500
C3—H3	0.9500	C18—H18	0.9500
O1—Sn1—O2	160.55 (8)	N1—C6—C1	116.0 (3)
O1—Sn1—N1	77.91 (8)	C5—C6—C1	122.5 (3)
O2—Sn1—N1	88.76 (9)	C8—C7—C5	119.9 (3)
O1—Sn1—N2	87.73 (9)	C8—C7—H7	120.0
O2—Sn1—N2	76.73 (9)	C5—C7—H7	120.0
N1—Sn1—N2	84.03 (9)	C7—C8—C9	119.4 (3)
O1—Sn1—Cl6	98.77 (6)	C7—C8—H8	120.3
O2—Sn1—Cl6	95.19 (6)	C9—C8—H8	120.3
N1—Sn1—Cl6	89.56 (7)	N1—C9—C8	122.0 (3)
N2—Sn1—Cl6	169.75 (7)	N1—C9—H9	119.0
O1—Sn1—Cl5	93.79 (6)	C8—C9—H9	119.0
O2—Sn1—Cl5	97.26 (6)	O2—C10—C11	124.0 (3)
N1—Sn1—Cl5	168.73 (7)	O2—C10—C15	120.0 (3)
N2—Sn1—Cl5	88.07 (6)	C11—C10—C15	116.0 (3)

Cl6—Sn1—C15	99.32 (3)	C10—C11—C12	122.2 (3)
C1—O1—Sn1	115.39 (17)	C10—C11—Cl4	119.5 (2)
C10—O2—Sn1	116.24 (18)	C12—C11—Cl4	118.3 (2)
C9—N1—C6	120.2 (3)	C13—C12—C11	120.4 (3)
C9—N1—Sn1	129.2 (2)	C13—C12—H12	119.8
C6—N1—Sn1	110.64 (18)	C11—C12—H12	119.8
C18—N2—C15	120.1 (3)	C12—C13—C14	121.5 (3)
C18—N2—Sn1	128.8 (2)	C12—C13—Cl3	119.0 (3)
C15—N2—Sn1	111.08 (19)	C14—C13—Cl3	119.5 (3)
O1—C1—C2	123.4 (3)	C15—C14—C13	116.7 (3)
O1—C1—C6	120.0 (3)	C15—C14—C16	117.0 (3)
C2—C1—C6	116.6 (3)	C13—C14—C16	126.3 (3)
C1—C2—C3	122.1 (3)	N2—C15—C14	121.6 (3)
C1—C2—Cl2	119.5 (2)	N2—C15—C10	115.3 (3)
C3—C2—Cl2	118.4 (2)	C14—C15—C10	123.2 (3)
C4—C3—C2	120.0 (3)	C17—C16—C14	119.8 (3)
C4—C3—H3	120.0	C17—C16—H16	120.1
C2—C3—H3	120.0	C14—C16—H16	120.1
C3—C4—C5	121.5 (3)	C16—C17—C18	120.1 (3)
C3—C4—Cl1	119.1 (2)	C16—C17—H17	119.9
C5—C4—Cl1	119.4 (2)	C18—C17—H17	119.9
C4—C5—C6	117.2 (3)	N2—C18—C17	121.4 (3)
C4—C5—C7	125.7 (3)	N2—C18—H18	119.3
C6—C5—C7	117.0 (3)	C17—C18—H18	119.3
N1—C6—C5	121.5 (3)		
O2—Sn1—O1—C1	-49.4 (3)	Sn1—N1—C6—C1	0.3 (3)
N1—Sn1—O1—C1	-1.69 (19)	C4—C5—C6—N1	179.5 (3)
N2—Sn1—O1—C1	-86.08 (19)	C7—C5—C6—N1	0.1 (4)
Cl6—Sn1—O1—C1	85.96 (19)	C4—C5—C6—C1	-0.1 (4)
Cl5—Sn1—O1—C1	-173.99 (18)	C7—C5—C6—C1	-179.5 (3)
O1—Sn1—O2—C10	-45.4 (4)	O1—C1—C6—N1	-1.9 (4)
N1—Sn1—O2—C10	-91.7 (2)	C2—C1—C6—N1	179.0 (2)
N2—Sn1—O2—C10	-7.57 (19)	O1—C1—C6—C5	177.8 (3)
Cl6—Sn1—O2—C10	178.82 (19)	C2—C1—C6—C5	-1.3 (4)
Cl5—Sn1—O2—C10	78.7 (2)	C4—C5—C7—C8	179.9 (3)
O1—Sn1—N1—C9	-179.7 (3)	C6—C5—C7—C8	-0.7 (4)
O2—Sn1—N1—C9	-13.9 (2)	C5—C7—C8—C9	0.2 (4)
N2—Sn1—N1—C9	-90.7 (3)	C6—N1—C9—C8	-1.5 (4)
Cl6—Sn1—N1—C9	81.3 (2)	Sn1—N1—C9—C8	178.9 (2)
Cl5—Sn1—N1—C9	-136.4 (3)	C7—C8—C9—N1	0.9 (5)
O1—Sn1—N1—C6	0.71 (18)	Sn1—O2—C10—C11	-173.6 (2)
O2—Sn1—N1—C6	166.44 (19)	Sn1—O2—C10—C15	7.9 (3)
N2—Sn1—N1—C6	89.65 (19)	O2—C10—C11—C12	-178.9 (3)
Cl6—Sn1—N1—C6	-98.36 (18)	C15—C10—C11—C12	-0.3 (4)
Cl5—Sn1—N1—C6	43.9 (4)	O2—C10—C11—Cl4	0.0 (4)
O1—Sn1—N2—C18	-8.2 (3)	C15—C10—C11—Cl4	178.6 (2)
O2—Sn1—N2—C18	-176.5 (3)	C10—C11—C12—C13	0.6 (5)

N1—Sn1—N2—C18	-86.3 (3)	C14—C11—C12—C13	-178.3 (2)
Cl6—Sn1—N2—C18	-137.9 (3)	C11—C12—C13—C14	-0.1 (5)
Cl5—Sn1—N2—C18	85.6 (3)	C11—C12—C13—Cl3	179.6 (2)
O1—Sn1—N2—C15	174.68 (19)	C12—C13—C14—C15	-0.7 (4)
O2—Sn1—N2—C15	6.47 (18)	Cl3—C13—C14—C15	179.6 (2)
N1—Sn1—N2—C15	96.60 (19)	C12—C13—C14—C16	178.4 (3)
Cl6—Sn1—N2—C15	45.0 (5)	Cl3—C13—C14—C16	-1.3 (4)
Cl5—Sn1—N2—C15	-91.45 (18)	C18—N2—C15—C14	-1.8 (4)
Sn1—O1—C1—C2	-178.4 (2)	Sn1—N2—C15—C14	175.6 (2)
Sn1—O1—C1—C6	2.5 (3)	C18—N2—C15—C10	178.0 (3)
O1—C1—C2—C3	-177.7 (3)	Sn1—N2—C15—C10	-4.6 (3)
C6—C1—C2—C3	1.3 (4)	C13—C14—C15—N2	-179.2 (3)
O1—C1—C2—Cl2	0.8 (4)	C16—C14—C15—N2	1.6 (4)
C6—C1—C2—Cl2	179.9 (2)	C13—C14—C15—C10	1.1 (4)
C1—C2—C3—C4	0.1 (4)	C16—C14—C15—C10	-178.1 (3)
Cl2—C2—C3—C4	-178.4 (2)	O2—C10—C15—N2	-1.7 (4)
C2—C3—C4—C5	-1.7 (4)	C11—C10—C15—N2	179.7 (3)
C2—C3—C4—Cl1	176.8 (2)	O2—C10—C15—C14	178.1 (3)
C3—C4—C5—C6	1.6 (4)	C11—C10—C15—C14	-0.6 (4)
Cl1—C4—C5—C6	-176.9 (2)	C15—C14—C16—C17	0.1 (4)
C3—C4—C5—C7	-179.0 (3)	C13—C14—C16—C17	-179.1 (3)
Cl1—C4—C5—C7	2.5 (4)	C14—C16—C17—C18	-1.5 (5)
C9—N1—C6—C5	1.0 (4)	C15—N2—C18—C17	0.2 (4)
Sn1—N1—C6—C5	-179.3 (2)	Sn1—N2—C18—C17	-176.6 (2)
C9—N1—C6—C1	-179.4 (3)	C16—C17—C18—N2	1.5 (5)
