

9-Oxo-4,5-diazafluoren-4-i um tetra-chloridoaurate(III)-4,5-diazafluoren-9-one (1/1)

Nasser Safari,^a Vahid Amani,^a Behrouz Notash^a and Seik Weng Ng^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 19 February 2009; accepted 24 February 2009

Key indicators: single-crystal X-ray study; $T = 118\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.025; wR factor = 0.067; data-to-parameter ratio = 16.7.

The Au^{III} atom in the title compound, $(\text{C}_{11}\text{H}_7\text{N}_2\text{O})[\text{AuCl}_4]\cdot\text{C}_{11}\text{H}_6\text{N}_2\text{O}$, is in a nearly square-planar environment defined by four Cl atoms. The protonated 9-oxo-4,5-diazafluoren-4-i um cation forms an $\text{N}-\text{H}\cdots\text{N}$ hydrogen bond with the neutral 4,5-diazafluoren-9-one molecule.

Related literature

For other 9-oxo-4,5-diazafluoren-4-i um tetrachloridometalates, see: Kulkarni *et al.* (2003); Menon *et al.* (1994); Ravikumar *et al.* (1995); Ravikumar & Lakshmi (1994); Zhang *et al.* (2003). For the synthesis of 4,5-diazafluoren-9-one, see: Henderson *et al.* (1984).



Experimental

Crystal data

$(\text{C}_{11}\text{H}_7\text{N}_2\text{O})[\text{AuCl}_4]\cdot\text{C}_{11}\text{H}_6\text{N}_2\text{O}$
 $M_r = 704.13$

Triclinic, $P\bar{1}$
 $a = 7.1035 (1)\text{ \AA}$
 $b = 12.6513 (2)\text{ \AA}$
 $c = 13.2366 (2)\text{ \AA}$
 $\alpha = 73.285 (1)^\circ$
 $\beta = 78.410 (1)^\circ$

$\gamma = 88.375 (1)^\circ$
 $V = 1115.51 (3)\text{ \AA}^3$
 $Z = 2$
 $\text{Mo } K\alpha$ radiation
 $\mu = 7.10\text{ mm}^{-1}$
 $T = 118\text{ K}$
 $0.20 \times 0.10 \times 0.10\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.331$, $T_{\max} = 0.537$
(expected range = 0.303–0.491)

9343 measured reflections
5044 independent reflections
4829 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.067$
 $S = 1.04$
5044 reflections
302 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 3.05\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.27\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (Å).

Au1—Cl1	2.2720 (8)	Au1—Cl3	2.2864 (8)
Au1—Cl2	2.2882 (8)	Au1—Cl4	2.2872 (8)

Table 2
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots N3	0.89 (1)	1.88 (1)	2.762 (4)	168 (4)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2184).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Henderson, L. J., Fronczeck, F. R. & Cherry, W. R. (1984). *J. Am. Chem. Soc.* **106**, 5876–5879.
- Kulkarni, P., Padhye, S. & Sinn, E. (2003). *Inorg. Chem. Commun.* **6**, 1129–1132.
- Menon, S., Balagopalakrishna, C., Rajasekharan, M. V. & Ramakrishna, B. L. (1994). *Inorg. Chem.* **33**, 950–954.
- Ravikumar, K. & Lakshmi, N. V. (1994). *Z. Kristallogr.* **209**, 56–57.
- Ravikumar, K., Venkata Lakshmi, N., Swamy, G. Y. S. K. & Chandra Mohan, K. (1995). *Acta Cryst. C* **51**, 1556–1558.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.
- Zhang, R.-L., Liu, H.-M., Zhao, J.-S., Gu, A.-P., He, S.-Y., Liu, J.-N. & Shi, Q.-Z. (2003). *Chin. J. Inorg. Chem.* **19**, 722–726.

supporting information

Acta Cryst. (2009). E65, m344 [doi:10.1107/S1600536809006734]

9-Oxo-4,5-diazafluoren-4-i um tetrachloridoaurate(III)-4,5-diazafluoren-9-one (1/1)

Nasser Safari, Vahid Amani, Behrouz Notash and Seik Weng Ng

S1. Experimental

Chloroauric acid trihydrate (0.25 g, 0.73 mmol) dissolved in acetonitrile (5 ml) was mixed with 4,5-diazafluoren-9-one (Henderson *et al.*, 1984) (0.13 g, 0.73 mmol) dissolved in methanol (5 ml). The yellow solution was set aside for the growth of crystals, which appeared after two weeks (yield 70%; m.p. 483 K).

S2. Refinement

C-bound H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.95 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The N-bound H atom was located in a difference Fourier map and refined isotropically, with a distance restraint of N—H = 0.88 (1) Å. The crystal diffracted strongly owing to the extremely heavy metal atom; however, its presence introduced severe absorption problems that could not be corrected analytically as the crystal did not have regular faces. The final difference Fourier map had a highest peak at 1.00 Å and a deepest hole at 1.00 Å from Au1 atom.




Figure 1

Molecular structure of the title compound. Displacement ellipsoids are drawn at the 70% probability level. The dashed line denotes hydrogen bond.

9-Oxo-4,5-diazafluoren-4-i um tetrachloridoaurate(III)-4,5-diazafluoren-9-one (1/1)*Crystal data*
 $M_r = 704.13$
Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 7.1035 (1) \text{ \AA}$
 $b = 12.6513 (2) \text{ \AA}$
 $c = 13.2366 (2) \text{ \AA}$
 $\alpha = 73.285 (1)^\circ$
 $\beta = 78.410 (1)^\circ$
 $\gamma = 88.375 (1)^\circ$
 $V = 1115.51 (3) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 672$
 $D_x = 2.096 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7741 reflections

 $\theta = 2.7\text{--}28.4^\circ$
 $\mu = 7.10 \text{ mm}^{-1}$
 $T = 118 \text{ K}$

Block, yellow

 $0.20 \times 0.10 \times 0.10 \text{ mm}$
*Data collection*Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.331, T_{\max} = 0.537$

9343 measured reflections

5044 independent reflections

4829 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$
 $\theta_{\max} = 27.5^\circ, \theta_{\min} = 1.6^\circ$
 $h = -9 \rightarrow 9$
 $k = -16 \rightarrow 16$
 $l = -17 \rightarrow 17$
*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.067$
 $S = 1.04$

5044 reflections

302 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0509P)^2 + 0.1124P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 3.05 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.27 \text{ e \AA}^{-3}$
Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Au1	0.298203 (14)	0.304054 (8)	0.394619 (8)	0.01445 (6)
Cl1	0.28189 (13)	0.32712 (7)	0.56013 (7)	0.02448 (17)
Cl2	0.31382 (13)	0.49176 (6)	0.32181 (7)	0.02314 (17)
Cl3	0.31133 (12)	0.28082 (6)	0.22834 (6)	0.01975 (16)
Cl4	0.28622 (13)	0.11653 (6)	0.46788 (7)	0.02373 (17)
O1	0.1850 (4)	0.3950 (2)	0.7782 (2)	0.0258 (5)
O2	0.3590 (4)	1.22613 (19)	0.9995 (2)	0.0239 (5)
N1	0.2353 (4)	0.7025 (2)	0.9069 (2)	0.0163 (5)
H1	0.247 (6)	0.7760 (9)	0.885 (3)	0.029 (11)*
N2	0.2032 (4)	0.7897 (2)	0.6684 (2)	0.0187 (5)
N3	0.3136 (4)	0.9265 (2)	0.8548 (2)	0.0156 (5)
N4	0.1257 (4)	0.8534 (2)	1.0949 (2)	0.0168 (5)

C1	0.2546 (5)	0.6495 (3)	1.0088 (3)	0.0186 (6)
H1A	0.2662	0.6912	1.0567	0.022*
C2	0.2576 (5)	0.5354 (3)	1.0441 (3)	0.0203 (6)
H2	0.2704	0.4994	1.1159	0.024*
C3	0.2420 (5)	0.4737 (3)	0.9747 (3)	0.0205 (6)
H3	0.2447	0.3954	0.9977	0.025*
C4	0.2222 (4)	0.5296 (3)	0.8712 (3)	0.0164 (6)
C5	0.2212 (4)	0.6449 (2)	0.8388 (2)	0.0153 (6)
C6	0.2012 (4)	0.4900 (3)	0.7772 (3)	0.0181 (6)
C7	0.1980 (5)	0.5922 (3)	0.6874 (3)	0.0182 (6)
C8	0.1879 (5)	0.6097 (3)	0.5801 (3)	0.0212 (7)
H8	0.1825	0.5501	0.5504	0.025*
C9	0.1863 (5)	0.7187 (3)	0.5186 (3)	0.0212 (7)
H9	0.1809	0.7349	0.4445	0.025*
C10	0.1924 (5)	0.8047 (3)	0.5642 (3)	0.0223 (7)
H10	0.1888	0.8781	0.5195	0.027*
C11	0.2057 (4)	0.6848 (3)	0.7249 (3)	0.0155 (6)
C12	0.3999 (4)	0.9788 (3)	0.7520 (3)	0.0181 (6)
H12	0.4178	0.9380	0.7009	0.022*
C13	0.4636 (4)	1.0882 (3)	0.7171 (3)	0.0180 (6)
H13	0.5212	1.1206	0.6436	0.022*
C14	0.4435 (4)	1.1511 (2)	0.7894 (3)	0.0176 (6)
H14	0.4858	1.2264	0.7675	0.021*
C15	0.3585 (4)	1.0977 (2)	0.8947 (3)	0.0154 (6)
C16	0.2951 (4)	0.9874 (2)	0.9225 (2)	0.0139 (6)
C17	0.3142 (4)	1.1372 (2)	0.9930 (3)	0.0165 (6)
C18	0.2124 (4)	1.0415 (2)	1.0813 (3)	0.0162 (6)
C19	0.1345 (4)	1.0290 (3)	1.1887 (3)	0.0181 (6)
H19	0.1383	1.0876	1.2200	0.022*
C20	0.0503 (4)	0.9262 (3)	1.2485 (3)	0.0199 (6)
H20	-0.0062	0.9133	1.3226	0.024*
C21	0.0490 (4)	0.8422 (3)	1.1996 (3)	0.0195 (6)
H21	-0.0094	0.7729	1.2426	0.023*
C22	0.2039 (4)	0.9522 (2)	1.0394 (2)	0.0143 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Au1	0.01648 (8)	0.01391 (8)	0.01188 (8)	0.00013 (5)	-0.00036 (5)	-0.00370 (5)
Cl1	0.0377 (5)	0.0222 (4)	0.0142 (4)	0.0028 (3)	-0.0047 (3)	-0.0066 (3)
Cl2	0.0343 (4)	0.0151 (4)	0.0193 (4)	-0.0012 (3)	-0.0052 (3)	-0.0038 (3)
Cl3	0.0262 (4)	0.0187 (4)	0.0144 (4)	-0.0011 (3)	-0.0020 (3)	-0.0061 (3)
Cl4	0.0333 (4)	0.0151 (4)	0.0194 (4)	0.0019 (3)	-0.0005 (3)	-0.0030 (3)
O1	0.0391 (14)	0.0147 (11)	0.0241 (13)	-0.0037 (10)	-0.0044 (11)	-0.0072 (10)
O2	0.0348 (13)	0.0135 (11)	0.0251 (13)	0.0004 (9)	-0.0077 (11)	-0.0074 (10)
N1	0.0197 (12)	0.0112 (12)	0.0171 (13)	0.0000 (9)	-0.0020 (10)	-0.0038 (10)
N2	0.0205 (13)	0.0146 (12)	0.0195 (14)	-0.0029 (10)	-0.0032 (11)	-0.0030 (10)
N3	0.0171 (12)	0.0149 (12)	0.0136 (12)	-0.0002 (9)	-0.0016 (10)	-0.0033 (10)

N4	0.0180 (12)	0.0155 (12)	0.0162 (13)	-0.0018 (9)	-0.0016 (10)	-0.0048 (10)
C1	0.0229 (15)	0.0171 (14)	0.0161 (15)	-0.0012 (11)	-0.0014 (12)	-0.0068 (12)
C2	0.0255 (16)	0.0161 (15)	0.0177 (16)	0.0011 (12)	-0.0038 (13)	-0.0031 (12)
C3	0.0228 (16)	0.0151 (14)	0.0211 (16)	0.0004 (12)	-0.0026 (13)	-0.0024 (12)
C4	0.0178 (14)	0.0136 (14)	0.0189 (15)	-0.0004 (11)	-0.0024 (12)	-0.0071 (12)
C5	0.0146 (13)	0.0143 (14)	0.0164 (15)	-0.0007 (10)	0.0001 (11)	-0.0054 (12)
C6	0.0189 (14)	0.0171 (15)	0.0191 (15)	0.0002 (11)	-0.0012 (12)	-0.0081 (12)
C7	0.0190 (14)	0.0161 (14)	0.0199 (16)	0.0005 (11)	-0.0021 (12)	-0.0071 (12)
C8	0.0212 (15)	0.0241 (16)	0.0198 (16)	-0.0026 (12)	-0.0037 (13)	-0.0089 (13)
C9	0.0194 (15)	0.0265 (17)	0.0169 (16)	-0.0016 (12)	-0.0025 (12)	-0.0054 (13)
C10	0.0224 (16)	0.0218 (16)	0.0197 (17)	-0.0023 (12)	-0.0048 (13)	-0.0007 (13)
C11	0.0137 (13)	0.0137 (14)	0.0181 (15)	-0.0001 (10)	-0.0023 (11)	-0.0035 (12)
C12	0.0190 (14)	0.0188 (15)	0.0162 (15)	0.0008 (11)	-0.0023 (12)	-0.0056 (12)
C13	0.0184 (14)	0.0176 (14)	0.0141 (14)	-0.0006 (11)	-0.0007 (12)	-0.0002 (12)
C14	0.0169 (14)	0.0130 (13)	0.0214 (16)	-0.0009 (11)	-0.0040 (12)	-0.0024 (12)
C15	0.0156 (13)	0.0128 (13)	0.0177 (15)	0.0032 (10)	-0.0046 (12)	-0.0035 (12)
C16	0.0122 (13)	0.0133 (13)	0.0155 (14)	0.0014 (10)	-0.0027 (11)	-0.0032 (11)
C17	0.0178 (14)	0.0131 (13)	0.0190 (15)	0.0030 (11)	-0.0057 (12)	-0.0040 (12)
C18	0.0149 (13)	0.0157 (14)	0.0183 (15)	0.0011 (11)	-0.0043 (11)	-0.0049 (12)
C19	0.0193 (14)	0.0184 (14)	0.0193 (15)	0.0028 (11)	-0.0060 (12)	-0.0084 (12)
C20	0.0170 (14)	0.0270 (16)	0.0148 (15)	0.0024 (12)	-0.0009 (12)	-0.0064 (13)
C21	0.0181 (14)	0.0172 (14)	0.0206 (16)	-0.0011 (11)	-0.0021 (12)	-0.0025 (12)
C22	0.0120 (12)	0.0149 (13)	0.0165 (14)	0.0020 (10)	-0.0036 (11)	-0.0047 (11)

Geometric parameters (\AA , $^\circ$)

Au1—Cl1	2.2720 (8)	C7—C8	1.389 (5)
Au1—Cl2	2.2882 (8)	C7—C11	1.405 (4)
Au1—Cl3	2.2864 (8)	C8—C9	1.386 (5)
Au1—Cl4	2.2872 (8)	C8—H8	0.9500
O1—C6	1.208 (4)	C9—C10	1.393 (5)
O2—C17	1.209 (4)	C9—H9	0.9500
N1—C5	1.331 (4)	C10—H10	0.9500
N1—C1	1.355 (4)	C12—C13	1.385 (4)
N1—H1	0.89 (1)	C12—H12	0.9500
N2—C11	1.325 (4)	C13—C14	1.396 (4)
N2—C10	1.355 (4)	C13—H13	0.9500
N3—C16	1.326 (4)	C14—C15	1.381 (4)
N3—C12	1.354 (4)	C14—H14	0.9500
N4—C22	1.326 (4)	C15—C16	1.400 (4)
N4—C21	1.352 (4)	C15—C17	1.498 (4)
C1—C2	1.384 (4)	C16—C22	1.493 (4)
C1—H1A	0.9500	C17—C18	1.500 (4)
C2—C3	1.387 (5)	C18—C19	1.382 (4)
C2—H2	0.9500	C18—C22	1.402 (4)
C3—C4	1.383 (5)	C19—C20	1.392 (4)
C3—H3	0.9500	C19—H19	0.9500
C4—C5	1.397 (4)	C20—C21	1.396 (4)

C4—C6	1.503 (4)	C20—H20	0.9500
C5—C11	1.471 (4)	C21—H21	0.9500
C6—C7	1.486 (5)		
Cl1—Au1—Cl3	179.43 (3)	N2—C10—C9	123.9 (3)
Cl1—Au1—Cl4	90.27 (3)	N2—C10—H10	118.1
Cl3—Au1—Cl4	89.73 (3)	C9—C10—H10	118.1
Cl1—Au1—Cl2	89.47 (3)	N2—C11—C7	126.7 (3)
Cl3—Au1—Cl2	90.55 (3)	N2—C11—C5	125.6 (3)
Cl4—Au1—Cl2	179.30 (3)	C7—C11—C5	107.7 (3)
C5—N1—C1	120.1 (3)	N3—C12—C13	123.8 (3)
C5—N1—H1	122 (3)	N3—C12—H12	118.1
C1—N1—H1	118 (3)	C13—C12—H12	118.1
C11—N2—C10	114.0 (3)	C12—C13—C14	120.3 (3)
C16—N3—C12	115.2 (3)	C12—C13—H13	119.8
C22—N4—C21	115.0 (3)	C14—C13—H13	119.8
N1—C1—C2	120.9 (3)	C15—C14—C13	116.1 (3)
N1—C1—H1A	119.6	C15—C14—H14	121.9
C2—C1—H1A	119.6	C13—C14—H14	121.9
C1—C2—C3	120.1 (3)	C14—C15—C16	119.8 (3)
C1—C2—H2	120.0	C14—C15—C17	131.3 (3)
C3—C2—H2	120.0	C16—C15—C17	108.9 (3)
C4—C3—C2	118.0 (3)	N3—C16—C15	124.8 (3)
C4—C3—H3	121.0	N3—C16—C22	126.9 (3)
C2—C3—H3	121.0	C15—C16—C22	108.3 (3)
C3—C4—C5	120.0 (3)	O2—C17—C15	126.5 (3)
C3—C4—C6	132.0 (3)	O2—C17—C18	128.0 (3)
C5—C4—C6	107.9 (3)	C15—C17—C18	105.4 (2)
N1—C5—C4	121.0 (3)	C19—C18—C22	119.4 (3)
N1—C5—C11	129.2 (3)	C19—C18—C17	132.1 (3)
C4—C5—C11	109.8 (3)	C22—C18—C17	108.5 (3)
O1—C6—C7	128.9 (3)	C18—C19—C20	116.5 (3)
O1—C6—C4	126.0 (3)	C18—C19—H19	121.8
C7—C6—C4	105.1 (3)	C20—C19—H19	121.8
C8—C7—C11	118.2 (3)	C19—C20—C21	120.0 (3)
C8—C7—C6	132.5 (3)	C19—C20—H20	120.0
C11—C7—C6	109.4 (3)	C21—C20—H20	120.0
C9—C8—C7	116.6 (3)	N4—C21—C20	124.0 (3)
C9—C8—H8	121.7	N4—C21—H21	118.0
C7—C8—H8	121.7	C20—C21—H21	118.0
C8—C9—C10	120.7 (3)	N4—C22—C18	125.2 (3)
C8—C9—H9	119.7	N4—C22—C16	126.0 (3)
C10—C9—H9	119.7	C18—C22—C16	108.8 (3)
C5—N1—C1—C2	-0.8 (5)	C16—N3—C12—C13	-0.8 (5)
N1—C1—C2—C3	0.4 (5)	N3—C12—C13—C14	1.0 (5)
C1—C2—C3—C4	-0.4 (5)	C12—C13—C14—C15	0.1 (5)
C2—C3—C4—C5	0.9 (5)	C13—C14—C15—C16	-1.3 (4)

C2—C3—C4—C6	−179.7 (3)	C13—C14—C15—C17	179.7 (3)
C1—N1—C5—C4	1.2 (5)	C12—N3—C16—C15	−0.5 (4)
C1—N1—C5—C11	−177.8 (3)	C12—N3—C16—C22	179.1 (3)
C3—C4—C5—N1	−1.3 (5)	C14—C15—C16—N3	1.5 (5)
C6—C4—C5—N1	179.2 (3)	C17—C15—C16—N3	−179.3 (3)
C3—C4—C5—C11	177.9 (3)	C14—C15—C16—C22	−178.0 (3)
C6—C4—C5—C11	−1.6 (3)	C17—C15—C16—C22	1.2 (3)
C3—C4—C6—O1	5.2 (6)	C14—C15—C17—O2	−5.5 (6)
C5—C4—C6—O1	−175.3 (3)	C16—C15—C17—O2	175.4 (3)
C3—C4—C6—C7	−176.8 (3)	C14—C15—C17—C18	177.2 (3)
C5—C4—C6—C7	2.8 (3)	C16—C15—C17—C18	−1.9 (3)
O1—C6—C7—C8	−4.5 (6)	O2—C17—C18—C19	3.9 (6)
C4—C6—C7—C8	177.6 (4)	C15—C17—C18—C19	−178.8 (3)
O1—C6—C7—C11	175.0 (3)	O2—C17—C18—C22	−175.3 (3)
C4—C6—C7—C11	−2.9 (3)	C15—C17—C18—C22	2.0 (3)
C11—C7—C8—C9	0.0 (5)	C22—C18—C19—C20	−0.3 (4)
C6—C7—C8—C9	179.4 (3)	C17—C18—C19—C20	−179.5 (3)
C7—C8—C9—C10	−0.7 (5)	C18—C19—C20—C21	0.4 (5)
C11—N2—C10—C9	−0.6 (5)	C22—N4—C21—C20	−0.4 (5)
C8—C9—C10—N2	1.0 (5)	C19—C20—C21—N4	−0.1 (5)
C10—N2—C11—C7	−0.2 (5)	C21—N4—C22—C18	0.5 (4)
C10—N2—C11—C5	178.5 (3)	C21—N4—C22—C16	−178.9 (3)
C8—C7—C11—N2	0.5 (5)	C19—C18—C22—N4	−0.1 (5)
C6—C7—C11—N2	−179.1 (3)	C17—C18—C22—N4	179.2 (3)
C8—C7—C11—C5	−178.4 (3)	C19—C18—C22—C16	179.3 (3)
C6—C7—C11—C5	2.0 (4)	C17—C18—C22—C16	−1.3 (3)
N1—C5—C11—N2	0.0 (5)	N3—C16—C22—N4	0.0 (5)
C4—C5—C11—N2	−179.1 (3)	C15—C16—C22—N4	179.5 (3)
N1—C5—C11—C7	178.9 (3)	N3—C16—C22—C18	−179.5 (3)
C4—C5—C11—C7	−0.2 (4)	C15—C16—C22—C18	0.1 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···N3	0.89 (1)	1.88 (1)	2.762 (4)	168 (4)