

## *N,N'-Bis(2-hydroxy-3-ethoxybenzylidene)butane-1,4-diamine*

Hoong-Kun Fun,<sup>a\*</sup> Reza Kia,<sup>a‡</sup> Hadi Kargar<sup>b</sup> and Arezoo Jamshidvand<sup>b</sup>

<sup>a</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>b</sup>Department of Chemistry, School of Science, Payame Noor University (PNU), Ardakan, Yazd, Iran  
Correspondence e-mail: hkfun@usm.my


Received 27 February 2009; accepted 2 March 2009

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.049;  $wR$  factor = 0.131; data-to-parameter ratio = 22.9.

The title Schiff base compound,  $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4$ , lies across a crystallographic inversion centre and adopts an *E* configuration with respect to the  $\text{C}=\text{N}$  bond. Pairs of weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  interactions link neighbouring molecules into dimers with an  $R_2^2(28)$  ring motif. The crystal structure is stabilized by intermolecular  $\text{C}-\text{H}\cdots\pi$  interactions. An intramolecular  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bond occurs.

### Related literature

For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For information on Schiff base ligands and complexes and their applications, see, for example: Calligaris & Randaccio (1987); Casellato & Vigato (1977); Fun & Kia (2008a,b). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



### Experimental

#### Crystal data

$\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4$

$M_r = 384.46$

‡ Additional correspondence author: e-mail: zsrkk@yahoo.com.

Triclinic, $P\bar{1}$	$V = 493.23 (2)\text{ \AA}^3$
$a = 6.8647 (2)\text{ \AA}$	$Z = 1$
$b = 6.9052 (2)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 10.8083 (3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$\alpha = 92.779 (2)^\circ$	$T = 100\text{ K}$
$\beta = 99.908 (2)^\circ$	$0.45 \times 0.19 \times 0.07\text{ mm}$
$\gamma = 101.239 (2)^\circ$	

#### Data collection

Bruker SMART APEXII CCD area-detector diffractometer	8962 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2005)	2954 independent reflections
$T_{\min} = 0.961$ , $T_{\max} = 0.994$	2157 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.033$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$	129 parameters
$wR(F^2) = 0.131$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.43\text{ e \AA}^{-3}$
2954 reflections	$\Delta\rho_{\min} = -0.27\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots\text{A}$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
O1—H1…N1	0.84	1.82	2.5638 (14)	147
C5—H5A…O1 <sup>i</sup>	0.95	2.59	3.2268 (14)	125
C11—H11B…Cg1 <sup>ii</sup>	0.98	2.96	3.5403 (15)	145

Symmetry codes: (i)  $x, y - 1, z$ ; (ii)  $-x + 2, -y + 1, -z + 1$ . Cg1 is the centroid of the C1–C6 benzene ring.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. RK thanks Universiti Sains Malaysia for the award of a post-doctoral research fellowship. HK and AJ thank PNU for financial support. HKF also thanks Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2734).

### References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Calligaris, M. & Randaccio, L. (1987). *Comprehensive Coordination Chemistry*, Vol. 2, edited by G. Wilkinson, pp. 715–738. London: Pergamon.
- Casellato, U. & Vigato, P. A. (1977). *Coord. Chem. Rev.* **23**, 31–50.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Fun, H.-K. & Kia, R. (2008a). *Acta Cryst. E64*, m1081–m1082.
- Fun, H.-K. & Kia, R. (2008b). *Acta Cryst. E64*, m1116–m1117.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

# supporting information

*Acta Cryst.* (2009). E65, o706 [doi:10.1107/S1600536809007545]

## **N,N'-Bis(2-hydroxy-3-ethoxybenzylidene)butane-1,4-diamine**

**Hoong-Kun Fun, Reza Kia, Hadi Kargar and Arezoo Jamshidvand**

### **S1. Comment**

The condensation of primary amines with carbonyl compounds yields Schiff base (Casellato & Vigato, 1977) that are still one of the most prevalent mixed-donor ligands in coordination chemistry. In the past two decades, the synthesis, structure and properties of Schiff base complexes have stimulated much interest for their noteworthy contributions in single molecule-based magnetism, materials science, catalysis of many reactions like carbonylation, hydroformylation, reduction, oxidation, epoxidation and hydrolysis (Casellato & Vigato 1977). In comparison to the Schiff base metal complexes, only a relatively small number of free Schiff base ligands have been characterized (Calligaris & Randaccio, 1987). As an extension of our work (Fun & Kia 2008*a,b*) on the structural characterization of Schiff base ligands, the title compound is reported here.


The molecule of the title compound, Fig 1, lies across a crystallographic inversion centre and adopts an *E* configuration with respect to the azomethine ( $\text{C}=\text{N}$ ) bond. The asymmetric unit of the compound is composed of one-half of the molecule. The imino group is coplanar with the benzene ring. Pairs of intermolecular  $\text{C}-\text{H}\cdots\text{O}$  interactions link neighbouring molecules into dimers with a  $\text{R}_2^2(28)$  ring motif (Bernstein *et al.*, 1995). The crystal structure is stabilized by intermolecular  $\text{C}-\text{H}\cdots\pi$  interactions [ $Cg1$  is the centroid of the C1–C6 benzene ring] (Table 1).

### **S2. Experimental**


The synthetic method has been described earlier (Fun, Kia & Kargar *et al.*, 2008*b*), except that 2-hydroxy-3-ethoxy-salicylaldehyde was used. Single crystals suitable for *X*-ray diffraction were obtained by evaporation of an ethanol solution at room temperature.

### **S3. Refinement**

H atom of the hydroxy group was positioned by a freely rotating O—H bond and constrained with a fixed distance of 0.84 Å. The rest of the hydrogen atoms were positioned geometrically with a riding model approximation with  $\text{C}-\text{H} = 0.95\text{--}0.99$  Å and  $U_{\text{iso}}(\text{H}) = 1.2$  or 1.5 (C & O). A rotating group model was used for methyl group.

**Figure 1**

The molecular structure of the title compound with atom labels and 50% probability ellipsoids for non-H atoms. The suffix A corresponds to symmetry code  $(-x + 2, -y + 1, -z + 2)$ .

**Figure 2**

The crystal packing of the title compound, viewed down the *c* axis showing dimer formation by R<sup>2</sup>(28) ring motif.

### *N,N'*-Bis(2-hydroxy-3-ethoxybenzylidene)butane-1,4-diamine

#### Crystal data

C<sub>22</sub>H<sub>28</sub>N<sub>2</sub>O<sub>4</sub>

*M*<sub>r</sub> = 384.46

Triclinic, P $\bar{1}$

Hall symbol: -P 1

*a* = 6.8647 (2) Å

*b* = 6.9052 (2) Å

*c* = 10.8083 (3) Å

$\alpha$  = 92.779 (2) $^\circ$

$\beta$  = 99.908 (2) $^\circ$

$\gamma$  = 101.239 (2) $^\circ$

*V* = 493.23 (2) Å<sup>3</sup>

*Z* = 1

*F*(000) = 206

*D*<sub>x</sub> = 1.294 Mg m<sup>-3</sup>

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 2475 reflections

$\theta$  = 2.5–29.4 $^\circ$

$\mu$  = 0.09 mm<sup>-1</sup>

$T = 100$  K

Plate, yellow

 $0.45 \times 0.19 \times 0.07$  mm*Data collection*Bruker SMART APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\varphi$  and  $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Bruker, 2005) $T_{\min} = 0.961$ ,  $T_{\max} = 0.994$ 

8962 measured reflections

2954 independent reflections

2157 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.033$  $\theta_{\max} = 30.4^\circ$ ,  $\theta_{\min} = 1.9^\circ$  $h = -9 \rightarrow 9$  $k = -9 \rightarrow 9$  $l = -14 \rightarrow 15$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$  $wR(F^2) = 0.131$  $S = 1.04$ 

2954 reflections

129 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0681P)^2 + 0.047P]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.43$  e  $\text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.27$  e  $\text{\AA}^{-3}$ *Special details*

**Experimental.** The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1)K.

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.69496 (13)	0.51130 (12)	0.23213 (9)	0.0191 (2)
H1	0.5792	0.4902	0.1870	0.029*
O2	1.05734 (12)	0.52787 (12)	0.36073 (8)	0.0178 (2)
N1	0.35928 (15)	0.30726 (15)	0.10848 (10)	0.0181 (2)
C1	0.75200 (17)	0.33702 (16)	0.25124 (11)	0.0149 (2)
C2	0.94642 (17)	0.34181 (16)	0.32117 (11)	0.0157 (2)
C3	1.00920 (18)	0.16571 (17)	0.34438 (12)	0.0177 (2)
H3A	1.1398	0.1690	0.3921	0.021*
C4	0.88179 (18)	-0.01729 (17)	0.29812 (12)	0.0188 (3)
H4A	0.9259	-0.1372	0.3146	0.023*
C5	0.69162 (18)	-0.02237 (17)	0.22845 (12)	0.0178 (2)
H5A	0.6058	-0.1462	0.1964	0.021*

C6	0.62455 (17)	0.15406 (16)	0.20476 (11)	0.0157 (2)
C7	0.42169 (18)	0.14831 (17)	0.13323 (11)	0.0169 (2)
H7A	0.3346	0.0238	0.1046	0.020*
C8	0.15522 (17)	0.29668 (17)	0.03663 (12)	0.0185 (3)
H8A	0.0561	0.2041	0.0735	0.022*
H8B	0.1478	0.2459	-0.0517	0.022*
C9	0.10389 (17)	0.50109 (17)	0.03934 (11)	0.0170 (2)
H9A	0.2076	0.5946	0.0065	0.020*
H9B	0.1064	0.5488	0.1276	0.020*
C10	1.26442 (17)	0.54481 (17)	0.42136 (12)	0.0182 (3)
H10A	1.3370	0.4755	0.3676	0.022*
H10B	1.2713	0.4857	0.5032	0.022*
C11	1.35712 (19)	0.76329 (19)	0.44083 (13)	0.0227 (3)
H11A	1.5003	0.7823	0.4795	0.034*
H11B	1.2866	0.8291	0.4964	0.034*
H11C	1.3448	0.8205	0.3593	0.034*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0164 (4)	0.0136 (4)	0.0265 (5)	0.0048 (3)	-0.0003 (3)	0.0029 (3)
O2	0.0134 (4)	0.0146 (4)	0.0234 (5)	0.0009 (3)	0.0006 (3)	0.0005 (3)
N1	0.0145 (5)	0.0188 (5)	0.0215 (5)	0.0055 (4)	0.0021 (4)	0.0024 (4)
C1	0.0163 (6)	0.0129 (5)	0.0168 (6)	0.0040 (4)	0.0054 (4)	0.0031 (4)
C2	0.0154 (6)	0.0149 (5)	0.0172 (6)	0.0021 (4)	0.0051 (4)	0.0015 (4)
C3	0.0143 (6)	0.0190 (6)	0.0203 (6)	0.0050 (4)	0.0023 (5)	0.0029 (4)
C4	0.0194 (6)	0.0149 (5)	0.0237 (6)	0.0069 (4)	0.0044 (5)	0.0039 (4)
C5	0.0175 (6)	0.0133 (5)	0.0223 (6)	0.0030 (4)	0.0029 (5)	0.0015 (4)
C6	0.0142 (6)	0.0157 (5)	0.0176 (6)	0.0038 (4)	0.0033 (4)	0.0021 (4)
C7	0.0154 (6)	0.0154 (5)	0.0192 (6)	0.0019 (4)	0.0027 (5)	0.0011 (4)
C8	0.0137 (6)	0.0187 (6)	0.0221 (6)	0.0043 (4)	0.0001 (5)	0.0009 (5)
C9	0.0157 (6)	0.0176 (5)	0.0186 (6)	0.0055 (4)	0.0032 (5)	0.0025 (4)
C10	0.0128 (6)	0.0193 (6)	0.0221 (6)	0.0029 (4)	0.0030 (5)	0.0009 (4)
C11	0.0150 (6)	0.0219 (6)	0.0291 (7)	0.0006 (4)	0.0021 (5)	0.0027 (5)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C1	1.3507 (12)	C6—C7	1.4641 (15)
O1—H1	0.8400	C7—H7A	0.9500
O2—C2	1.3662 (13)	C8—C9	1.5201 (15)
O2—C10	1.4391 (13)	C8—H8A	0.9900
N1—C7	1.2776 (14)	C8—H8B	0.9900
N1—C8	1.4666 (14)	C9—C9 <sup>i</sup>	1.527 (2)
C1—C6	1.4037 (16)	C9—H9A	0.9900
C1—C2	1.4096 (16)	C9—H9B	0.9900
C2—C3	1.3871 (15)	C10—C11	1.5081 (17)
C3—C4	1.4033 (17)	C10—H10A	0.9900
C3—H3A	0.9500	C10—H10B	0.9900

C4—C5	1.3833 (16)	C11—H11A	0.9800
C4—H4A	0.9500	C11—H11B	0.9800
C5—C6	1.4036 (15)	C11—H11C	0.9800
C5—H5A	0.9500		
C1—O1—H1	109.5	N1—C8—C9	109.97 (10)
C2—O2—C10	117.43 (9)	N1—C8—H8A	109.7
C7—N1—C8	120.11 (11)	C9—C8—H8A	109.7
O1—C1—C6	122.32 (10)	N1—C8—H8B	109.7
O1—C1—C2	118.03 (10)	C9—C8—H8B	109.7
C6—C1—C2	119.65 (10)	H8A—C8—H8B	108.2
O2—C2—C3	125.83 (11)	C8—C9—C9 <sup>i</sup>	111.80 (13)
O2—C2—C1	114.48 (9)	C8—C9—H9A	109.3
C3—C2—C1	119.70 (11)	C9 <sup>i</sup> —C9—H9A	109.3
C2—C3—C4	120.70 (11)	C8—C9—H9B	109.3
C2—C3—H3A	119.7	C9 <sup>i</sup> —C9—H9B	109.3
C4—C3—H3A	119.7	H9A—C9—H9B	107.9
C5—C4—C3	119.72 (11)	O2—C10—C11	106.44 (9)
C5—C4—H4A	120.1	O2—C10—H10A	110.4
C3—C4—H4A	120.1	C11—C10—H10A	110.4
C4—C5—C6	120.48 (11)	O2—C10—H10B	110.4
C4—C5—H5A	119.8	C11—C10—H10B	110.4
C6—C5—H5A	119.8	H10A—C10—H10B	108.6
C5—C6—C1	119.75 (10)	C10—C11—H11A	109.5
C5—C6—C7	120.42 (11)	C10—C11—H11B	109.5
C1—C6—C7	119.83 (10)	H11A—C11—H11B	109.5
N1—C7—C6	121.38 (11)	C10—C11—H11C	109.5
N1—C7—H7A	119.3	H11A—C11—H11C	109.5
C6—C7—H7A	119.3	H11B—C11—H11C	109.5
C10—O2—C2—C3	6.39 (17)	C4—C5—C6—C7	-178.73 (11)
C10—O2—C2—C1	-173.76 (10)	O1—C1—C6—C5	-179.51 (11)
O1—C1—C2—O2	-0.84 (16)	C2—C1—C6—C5	0.09 (17)
C6—C1—C2—O2	179.55 (10)	O1—C1—C6—C7	-0.25 (17)
O1—C1—C2—C3	179.02 (10)	C2—C1—C6—C7	179.35 (10)
C6—C1—C2—C3	-0.60 (17)	C8—N1—C7—C6	-179.99 (10)
O2—C2—C3—C4	-179.65 (11)	C5—C6—C7—N1	-178.57 (11)
C1—C2—C3—C4	0.51 (18)	C1—C6—C7—N1	2.18 (18)
C2—C3—C4—C5	0.10 (18)	C7—N1—C8—C9	170.13 (11)
C3—C4—C5—C6	-0.61 (18)	N1—C8—C9—C9 <sup>i</sup>	177.44 (12)
C4—C5—C6—C1	0.51 (18)	C2—O2—C10—C11	173.37 (10)

Symmetry code: (i)  $-x, -y+1, -z$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^{\circ}$ )

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1—H1 $\cdots$ N1	0.84	1.82	2.5638 (14)	147

---

C5—H5A···O1 <sup>ii</sup>	0.95	2.59	3.2268 (14)	125
C11—H11B···Cg1 <sup>iii</sup>	0.98	2.96	3.5403 (15)	145

---

Symmetry codes: (ii)  $x, y-1, z$ ; (iii)  $-x+2, -y+1, -z+1$ .