

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (3-Anilino-1-phenyliminothioureato)-chloridodimethyltin(IV)

Jorge A. Delgado D.,<sup>a</sup> Coco K. Y. A. Okio<sup>a\*</sup> and Richard Welter<sup>b</sup><sup>a</sup>Departamento de Química, Universidad Nacional de Colombia, Sede Bogotá, Bogotá, Colombia, and <sup>b</sup>Laboratoire DECOMET, UMR-CNRS 7177, Université Louis Pasteur, 4 rue Blaise Pascal, 67000 Strasbourg, France

Correspondence e-mail: kaokio@unal.edu.co


Received 10 March 2009; accepted 17 March 2009

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.137; data-to-parameter ratio = 20.8.

In the title compound,  $[\text{Sn}(\text{CH}_3)_2(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})\text{Cl}]$ , the Sn atom is five-coordinated in a distorted trigonal-bipyramidal geometry, with two methyl groups and one S atom in the equatorial plane, and one N atom and the Cl atom occupying the apical positions.

## Related literature

For related structures, see: Labib *et al.* (1996). For the biological and pharmaceutical applications of organotin derivatives, see: Davies & Smith (1982); Diop *et al.* (2003); Okio *et al.* (2003).



## Experimental

## Crystal data

$[\text{Sn}(\text{CH}_3)_2(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})\text{Cl}]$   
 $M_r = 439.53$   
 Orthorhombic, *Pbna*  
 $a = 11.6850$  (2) Å  
 $b = 14.8920$  (5) Å  
 $c = 20.6790$  (7) Å

$V = 3598.42$  (18) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.69$  mm<sup>-1</sup>  
 $T = 173$  K  
 $0.10 \times 0.10 \times 0.10$  mm

## Data collection

Nonius KappaCCD diffractometer  
 Absorption correction: none  
 25644 measured reflections

4130 independent reflections  
 2731 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.115$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.137$   
 $S = 1.09$   
 4130 reflections

199 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 1.13$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -1.84$  e Å<sup>-3</sup>

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2197).

## References

- Davies, A. G. & Smith, P. G. (1982). *Comprehensive Organometallic Chemistry*, edited by G. Wilkinson, F. Gordon, A. Stone & E. W. Abel, pp. 519–616. New York: Pergamon Press.
- Diop, L., Mahieu, B., Mahon, M. F., Molloy, K. C. & Okio, K. Y. A. (2003). *Appl. Organomet. Chem.* **17**, 881–882.
- Labib, L., Khalil, T. E., Iskander, M. F. & Refaar, L. S. (1996). *Polyhedron*, **15**, 349–357.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Okio, K. Y. A., Diop, L., Barry, A. H., Jouini, T. & Ortega, S. H. (2003). *Appl. Organomet. Chem.* **17**, 317–318.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2009). E65, m426 [doi:10.1107/S160053680900988X]

**(3-Anilino-1-phenyliminothioureato)chloridodimethyltin(IV)**

**Jorge A. Delgado D., Coco K. Y. A. Okio and Richard Welter**

**S1. Comment**


The interest in the synthesis of new organotin derivatives is related to the diversity of structures that such compounds can form, and on the other side, to their biological and pharmaceutical applications (Davies & Smith, 1982, Okio *et al.*, 2003; Diop *et al.*, 2003). We report here the crystal structure of the title compound (I). The structure consists of discrete molecules (Fig. 1). The Sn atom is five-coordinate in a distorted trigonal-bipyramidal geometry. The distorted trigonal-bipyramidal coordination polyhedron has two methyl groups and one S atom in the equatorial plane, the N1 and Cl atom occupying the apical positions. The Sn—Cl and Sn—S bond distances are shorter than the values of 2.672 (1) Å and 2.478 (2) Å found in dimethylmonochloro[(*N*-(2-pyridinylmethylene) hydrazinecarbothioamidato)NS(− 1)]tin(IV) hemihydrate ( Labib *et al.*, 1996), while the Sn—N and Sn—C bonds are longer than the corresponding ones reported for the same previous compound, (2.359 (4) Å, 2.108 (7) Å and 2.105 (7) Å), as representative example.

**S2. Experimental**

Compound (I) was obtained by reacting dimethyltin (IV) dichloride (220 mg, 1 mmol) with 1,5 diphenylthiocarbazono (128 mg, 0.5 mmol) in dichloromethane under reflux for 3 h. Dark red crystals suitable for X-ray analysis were grown by slow solvent evaporation.

**S3. Refinement**

H atoms were positioned geometrically, with C—H distances in the range 0.95 - 0.98 Å, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2-1.5U_{\text{eq}}(\text{C})$ . H atom bonded to N4 was found in difference maps and positioned geometrically (N4-H = 0.88 Å) and was refined with a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms have been omitted.

### (3-Anilino-1-phenyliminothioureato)chloridodimethyltin(IV)

#### Crystal data

[Sn(CH<sub>3</sub>)<sub>2</sub>(C<sub>13</sub>H<sub>11</sub>N<sub>4</sub>S)Cl]

$M_r = 439.53$

Orthorhombic, *Pbna*

Hall symbol: -P 2ac 2b

$a = 11.6850$  (2) Å

$b = 14.8920$  (5) Å

$c = 20.6790$  (7) Å

$V = 3598.42$  (18) Å<sup>3</sup>

$Z = 8$

$F(000) = 1744$

$D_x = 1.623$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 14356 reflections

$\theta = 1.0$ – $27.5^\circ$

$\mu = 1.69$  mm<sup>-1</sup>

$T = 173$  K

Prism, dark red

$0.10 \times 0.10 \times 0.10$  mm

#### Data collection

Nonius KappaCCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\pi$  scans

25644 measured reflections

4130 independent reflections

2731 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.115$

$\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 2.0^\circ$

$h = -15 \rightarrow 15$

$k = -18 \rightarrow 19$

$l = -26 \rightarrow 22$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.137$   
 $S = 1.09$   
 4130 reflections  
 199 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.07P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 1.13 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.84 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn	0.18590 (3)	0.56103 (2)	0.509606 (17)	0.03083 (14)
Cl	0.03058 (11)	0.57242 (10)	0.59081 (6)	0.0489 (4)
S	0.30578 (10)	0.64147 (10)	0.58678 (6)	0.0440 (4)
N1	0.3690 (3)	0.5795 (2)	0.45454 (18)	0.0308 (9)
N2	0.4565 (3)	0.6110 (3)	0.48494 (17)	0.0325 (9)
N3	0.5342 (3)	0.6667 (2)	0.57432 (18)	0.0343 (9)
N4	0.5270 (3)	0.6917 (2)	0.63558 (19)	0.0364 (9)
H4N	0.4606	0.6889	0.6556	0.044*
C1	0.3037 (4)	0.5110 (3)	0.3553 (2)	0.0368 (11)
H1	0.2316	0.5011	0.3754	0.044*
C2	0.3220 (4)	0.4841 (3)	0.2918 (3)	0.0444 (13)
H2	0.2620	0.4559	0.2683	0.053*
C3	0.4263 (5)	0.4979 (3)	0.2626 (2)	0.0516 (14)
H3	0.4380	0.4797	0.2191	0.062*
C4	0.5137 (5)	0.5383 (4)	0.2966 (3)	0.0484 (14)
H4	0.5856	0.5474	0.2761	0.058*
C5	0.4985 (4)	0.5659 (3)	0.3598 (2)	0.0364 (11)
H5	0.5595	0.5934	0.3829	0.044*
C6	0.3923 (4)	0.5527 (3)	0.3896 (2)	0.0316 (10)
C7	0.6144 (4)	0.7333 (3)	0.7357 (2)	0.0423 (13)
H7	0.5451	0.7188	0.7573	0.051*
C8	0.7073 (5)	0.7652 (3)	0.7698 (3)	0.0474 (14)
H8	0.7011	0.7735	0.8152	0.057*
C9	0.8086 (4)	0.7852 (4)	0.7392 (3)	0.0503 (14)
H9	0.8718	0.8075	0.7633	0.060*

C10	0.8178 (4)	0.7729 (4)	0.6738 (3)	0.0518 (15)
H10	0.8884	0.7856	0.6529	0.062*
C11	0.7263 (5)	0.7424 (3)	0.6376 (3)	0.0425 (13)
H11	0.7330	0.7351	0.5921	0.051*
C12	0.6243 (4)	0.7226 (3)	0.6692 (2)	0.0342 (11)
C13	0.4384 (4)	0.6391 (3)	0.5470 (2)	0.0329 (10)
C14	0.2130 (5)	0.4204 (4)	0.5157 (3)	0.0438 (14)
H14A	0.1517	0.3890	0.4926	0.066*
H14B	0.2129	0.4019	0.5612	0.066*
H14C	0.2869	0.4053	0.4961	0.066*
C15	0.1051 (3)	0.6462 (3)	0.4410 (2)	0.0374 (11)
H15A	0.0860	0.6115	0.4022	0.056*
H15B	0.1572	0.6952	0.4293	0.056*
H15C	0.0350	0.6712	0.4598	0.056*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sn	0.0292 (2)	0.0351 (2)	0.0282 (2)	0.00016 (13)	-0.00196 (12)	0.00342 (13)
Cl	0.0390 (7)	0.0750 (10)	0.0326 (7)	0.0030 (6)	0.0051 (5)	0.0065 (6)
S	0.0344 (6)	0.0634 (9)	0.0342 (8)	0.0036 (6)	-0.0021 (5)	-0.0164 (6)
N1	0.033 (2)	0.029 (2)	0.031 (2)	-0.0025 (16)	-0.0031 (17)	0.0010 (16)
N2	0.035 (2)	0.030 (2)	0.032 (2)	-0.0017 (18)	-0.0044 (17)	-0.0016 (17)
N3	0.037 (2)	0.033 (2)	0.033 (2)	-0.0034 (16)	-0.0054 (17)	-0.0021 (18)
N4	0.039 (2)	0.037 (2)	0.034 (2)	0.0020 (18)	-0.0062 (17)	-0.0077 (18)
C1	0.044 (3)	0.035 (3)	0.032 (3)	0.002 (2)	-0.007 (2)	-0.001 (2)
C2	0.069 (4)	0.031 (3)	0.033 (3)	0.008 (3)	-0.017 (3)	-0.006 (2)
C3	0.082 (4)	0.042 (3)	0.031 (3)	0.013 (3)	0.000 (3)	-0.001 (2)
C4	0.066 (3)	0.043 (3)	0.036 (3)	0.011 (3)	0.018 (3)	0.006 (2)
C5	0.043 (3)	0.033 (3)	0.033 (3)	0.002 (2)	0.005 (2)	0.002 (2)
C6	0.040 (2)	0.027 (3)	0.028 (3)	0.0068 (19)	-0.002 (2)	0.0025 (19)
C7	0.045 (3)	0.046 (3)	0.036 (3)	0.012 (2)	-0.007 (2)	-0.009 (2)
C8	0.061 (3)	0.041 (3)	0.040 (3)	0.019 (2)	-0.021 (3)	-0.012 (2)
C9	0.050 (3)	0.041 (3)	0.060 (4)	-0.001 (3)	-0.024 (3)	-0.007 (3)
C10	0.048 (3)	0.048 (3)	0.060 (4)	-0.018 (2)	-0.013 (3)	0.006 (3)
C11	0.049 (3)	0.038 (3)	0.040 (3)	-0.007 (2)	-0.012 (2)	0.005 (2)
C12	0.037 (3)	0.025 (2)	0.040 (3)	0.004 (2)	-0.010 (2)	-0.004 (2)
C13	0.035 (2)	0.032 (3)	0.032 (3)	0.000 (2)	-0.004 (2)	-0.004 (2)
C14	0.042 (3)	0.037 (3)	0.052 (4)	0.003 (2)	-0.002 (2)	0.010 (2)
C15	0.036 (2)	0.038 (3)	0.038 (3)	0.000 (2)	-0.003 (2)	0.006 (2)

*Geometric parameters (Å, °)*

Sn—C14	2.122 (5)	C4—H4	0.9500
Sn—C15	2.125 (5)	C5—C6	1.399 (6)
Sn—S	2.4380 (13)	C5—H5	0.9500
Sn—N1	2.439 (4)	C7—C8	1.379 (7)
Sn—Cl	2.4784 (13)	C7—C12	1.388 (7)

S—C13	1.755 (4)	C7—H7	0.9500
N1—N2	1.289 (5)	C8—C9	1.375 (8)
N1—C6	1.428 (6)	C8—H8	0.9500
N2—C13	1.366 (6)	C9—C10	1.370 (8)
N3—C13	1.320 (5)	C9—H9	0.9500
N3—N4	1.323 (5)	C10—C11	1.382 (7)
N4—C12	1.410 (6)	C10—H10	0.9500
N4—H4N	0.8800	C11—C12	1.392 (7)
C1—C2	1.390 (7)	C11—H11	0.9500
C1—C6	1.400 (6)	C14—H14A	0.9800
C1—H1	0.9500	C14—H14B	0.9800
C2—C3	1.375 (7)	C14—H14C	0.9800
C2—H2	0.9500	C15—H15A	0.9800
C3—C4	1.377 (7)	C15—H15B	0.9800
C3—H3	0.9500	C15—H15C	0.9800
C4—C5	1.383 (7)		
C14—Sn—C15	134.0 (2)	C5—C6—N1	122.9 (4)
C14—Sn—S	111.14 (15)	C1—C6—N1	117.3 (4)
C15—Sn—S	113.52 (14)	C8—C7—C12	118.8 (5)
C14—Sn—N1	90.45 (17)	C8—C7—H7	120.6
C15—Sn—N1	90.60 (14)	C12—C7—H7	120.6
S—Sn—N1	75.30 (9)	C9—C8—C7	121.1 (5)
C14—Sn—Cl	97.86 (15)	C9—C8—H8	119.4
C15—Sn—Cl	94.95 (13)	C7—C8—H8	119.4
S—Sn—Cl	86.77 (4)	C10—C9—C8	119.5 (5)
N1—Sn—Cl	161.98 (9)	C10—C9—H9	120.2
C13—S—Sn	100.99 (16)	C8—C9—H9	120.2
N2—N1—C6	114.2 (4)	C9—C10—C11	121.3 (5)
N2—N1—Sn	120.6 (3)	C9—C10—H10	119.4
C6—N1—Sn	125.1 (3)	C11—C10—H10	119.4
N1—N2—C13	116.6 (4)	C10—C11—C12	118.5 (5)
C13—N3—N4	116.3 (4)	C10—C11—H11	120.7
N3—N4—C12	120.8 (4)	C12—C11—H11	120.7
N3—N4—H4N	119.6	C7—C12—C11	120.8 (4)
C12—N4—H4N	119.6	C7—C12—N4	117.3 (4)
C2—C1—C6	119.5 (4)	C11—C12—N4	121.9 (4)
C2—C1—H1	120.3	N3—C13—N2	111.5 (4)
C6—C1—H1	120.3	N3—C13—S	122.8 (4)
C3—C2—C1	120.5 (5)	N2—C13—S	125.7 (3)
C3—C2—H2	119.7	Sn—C14—H14A	109.5
C1—C2—H2	119.7	Sn—C14—H14B	109.5
C2—C3—C4	120.0 (5)	H14A—C14—H14B	109.5
C2—C3—H3	120.0	Sn—C14—H14C	109.5
C4—C3—H3	120.0	H14A—C14—H14C	109.5
C3—C4—C5	121.1 (5)	H14B—C14—H14C	109.5
C3—C4—H4	119.4	Sn—C15—H15A	109.5
C5—C4—H4	119.4	Sn—C15—H15B	109.5

C4—C5—C6	119.2 (5)	H15A—C15—H15B	109.5
C4—C5—H5	120.4	Sn—C15—H15C	109.5
C6—C5—H5	120.4	H15A—C15—H15C	109.5
C5—C6—C1	119.8 (4)	H15B—C15—H15C	109.5
C14—Sn—S—C13	77.8 (2)	C2—C1—C6—N1	179.6 (4)
C15—Sn—S—C13	-91.0 (2)	N2—N1—C6—C5	-4.7 (6)
N1—Sn—S—C13	-6.86 (18)	Sn—N1—C6—C5	179.1 (3)
Cl—Sn—S—C13	175.02 (17)	N2—N1—C6—C1	174.9 (4)
C14—Sn—N1—N2	-104.0 (3)	Sn—N1—C6—C1	-1.3 (5)
C15—Sn—N1—N2	121.9 (3)	C12—C7—C8—C9	-0.8 (7)
S—Sn—N1—N2	7.7 (3)	C7—C8—C9—C10	-0.3 (8)
Cl—Sn—N1—N2	13.8 (5)	C8—C9—C10—C11	1.3 (9)
C14—Sn—N1—C6	71.9 (3)	C9—C10—C11—C12	-1.2 (8)
C15—Sn—N1—C6	-62.1 (3)	C8—C7—C12—C11	1.0 (7)
S—Sn—N1—C6	-176.3 (3)	C8—C7—C12—N4	-178.6 (4)
Cl—Sn—N1—C6	-170.2 (2)	C10—C11—C12—C7	0.0 (7)
C6—N1—N2—C13	179.0 (4)	C10—C11—C12—N4	179.6 (5)
Sn—N1—N2—C13	-4.6 (5)	N3—N4—C12—C7	-169.6 (4)
C13—N3—N4—C12	-179.8 (4)	N3—N4—C12—C11	10.9 (7)
C6—C1—C2—C3	0.2 (7)	N4—N3—C13—N2	-177.1 (4)
C1—C2—C3—C4	0.3 (8)	N4—N3—C13—S	4.1 (6)
C2—C3—C4—C5	-0.3 (8)	N1—N2—C13—N3	177.5 (4)
C3—C4—C5—C6	-0.3 (7)	N1—N2—C13—S	-3.8 (6)
C4—C5—C6—C1	0.8 (7)	Sn—S—C13—N3	-172.3 (4)
C4—C5—C6—N1	-179.6 (4)	Sn—S—C13—N2	9.1 (4)
C2—C1—C6—C5	-0.8 (7)		