

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Isopropyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate–ethyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate (0.6/0.4)

Anas Lataifeh, Heinz-Bernhard Kraatz and Michael C. Jennings*

Department of Chemistry, The University of Western Ontario, London, Ontario, Canada N6A 5B7


Correspondence e-mail: mjenning@uwo.ca

Received 13 February 2009; accepted 5 March 2009

 Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.040; wR factor = 0.100; data-to-parameter ratio = 13.1.

Herein we report the crystal structure and synthesis of two cocrystallized complexes, $[\text{Fe}(\text{C}_7\text{H}_7\text{O}_2)(\text{C}_9\text{H}_{12}\text{NO}_2)]_{0.6}$ – $[\text{Fe}(\text{C}_7\text{H}_7\text{O}_2)(\text{C}_8\text{H}_{10}\text{NO}_2)]_{0.4}$. The molecules crystallize as layers in the bc plane with van der Waals interactions allowing the alkyl chains to interact and the ferrocene units to form a herringbone pattern up the c axis. Every second layer is linked via $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonding. The two complexes were modelled as disordered in a ratio of 0.60:0.40.

Related literature

 For background information, see: Barišić *et al.* (2002, 2006); Pavlović *et al.* (2003).


Experimental

Crystal data

$[\text{Fe}(\text{C}_7\text{H}_7\text{O}_2)(\text{C}_9\text{H}_{12}\text{NO}_2)]_{0.6}$
 $[\text{Fe}(\text{C}_7\text{H}_7\text{O}_2)(\text{C}_8\text{H}_{10}\text{NO}_2)]_{0.4}$
 $M_r = 339.56$
 Monoclinic, $P2_1/c$
 $a = 9.7494$ (19) Å
 $b = 15.624$ (3) Å
 $c = 9.860$ (2) Å
 $\beta = 100.82$ (3)°
 $V = 1475.2$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.04$ mm⁻¹
 $T = 150$ K
 $0.40 \times 0.22 \times 0.10$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (SORTAV; Blessing, 1995)
 $T_{\min} = 0.740$, $T_{\max} = 0.906$
 14609 measured reflections
 3377 independent reflections
 2817 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.063$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.100$
 $S = 1.05$
 3377 reflections
 257 parameters
 391 restraints
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.44$ e Å⁻³
 $\Delta\rho_{\min} = -0.46$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N15A}-\text{H15A}\cdots\text{O7}^i$	0.88	1.93	2.793 (14)	168
$\text{N15B}-\text{H15B}\cdots\text{O7}^i$	0.88	2.21	2.962 (19)	143

 Symmetry code: (i) $-x + 2, -y + 2, -z + 1$.

Data collection: COLLECT (Nonius, 2001); cell refinement: DENZO-SMN (Otwinowski & Minor, 1997); data reduction: DENZO-SMN; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL/PC (Sheldrick, 2008); software used to prepare material for publication: SHELXTL/PC.

This work was supported by the NSERC.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KJ2117).

References

- Barišić, L., Rapić, V. & Kovač, V. (2002). *Croat. Chem. Acta*, **75**, 199–210.
 Barišić, L., Rapić, V. & Metzler-Nolte, N. (2006). *Eur. J. Inorg. Chem.* **20**, 4019–4021.
 Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Nonius (2001). COLLECT. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326, New York: Academic Press.
 Pavlović, G., Barišić, L., Rapić, V. & Kovač, V. (2003). *Acta Cryst.* **C59**, m55–m57.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, m401 [doi:10.1107/S1600536809008204]

Isopropyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate–ethyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate (0.6/0.4)

Anas Lataifeh, Heinz-Bernhard Kraatz and Michael C. Jennings

S1. Comment

Methoxy-2'-(tertbutyloxycarbonylamino)ferrocene-1-carboxylate [$\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpN(H)C(O)OCMe}_3\}$] (where Cp = $\eta^5\text{-C}_5\text{H}_4$) has been used as a synthon in the preparation of ferrocene amino acid (Fca) peptide conjugates. This unnatural amino acid provides a convenient route to C- or N-terminal labelling of α -amino acids and peptides in both solution and solid phase (Barišić *et al.* 2006). The C-terminal conjugation of the natural amino acid and peptides to Fca requires the removal of the *tert*-butyloxy (t-Boc) group under acidic conditions. The resultant product $\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpNH}_2\}$ is highly unstable, and usually coupled *in situ* to the active ester derivative of amino acids. In our attempts to trap and crystallize this species, the title compounds (1) and (2) were produced, which might give an insight into the decomposition pathways of the t-Boc group in acidic medium. It is believed that crystallization occurred during partial decomposition and two intermediates in the stepwise decomposition were isolated crystallographically.


There were two molecules co-crystallized in the asymmetric unit. The two molecules were crystallographically identical except for the side chain; one was -N(H)C(O)OCHMe_2 (60%) and the other was $\text{-N(H)C(O)OCH}_2\text{Me}$ (40%). The molecules revealed close contacts between the hydrogen atom attached to the nitrogen and the carbonyl atoms of the adjacent molecule. Essentially, dimers are formed *via* hydrogen bonding (Table 1).

S2. Experimental

The ferrocene compound $\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpN(H)C(O)OCMe}_3\}$ was prepared by standard procedures reported by Rapić and coworkers (Barišić *et al.*, 2002; Pavlović *et al.*, 2003). The synthesis of the ferrocene derivative $\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpNH}_2\}$ requires the removal of the t-Boc group using trifluoroacetic acid (TFA) under argon. After several minutes, the reaction progress was quenched with a base (triethyl amine). Orange plates of the mixed crystal of the title compounds [$\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpN(H)C(O)OCHMe}_2\}$] and [$\{\text{MeOC(O)Cp}\}\text{Fe}\{\text{CpN(H)C(O)OCH}_2\text{Me}\}$] were grown from a concentrated methylene chloride solution by slow diffusion of hexane.

S3. Refinement

All H atoms were positioned geometrically and constrained as riding atoms with C—H = 1.00 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for methyne H atoms and C—H = 0.99 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for methylene H atoms and C—H = 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and N—H = 0.88 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for amine H atoms. Soft proximity (SIMU) and rigid-bond restraints (DELU) were applied to the anisotropic displacement parameters.

**Figure 1**

The molecular structure of the title compound with 30% probability displacement ellipsoids and the atom labelling scheme.

Isopropyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate–ethyl *N*-[1'-(methoxycarbonyl)ferrocenyl]carbamate (0.6/0.4)

Crystal data

[Fe(C₇H₇O₂)(C₉H₁₂NO₂)]_{0.6}[Fe(C₇H₇O₂)(C₈H₁₀NO₂)]_{0.4}

M_r = 339.56

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 9.7494 (19) Å

b = 15.624 (3) Å

c = 9.860 (2) Å

β = 100.82 (3)°

V = 1475.2 (5) Å³

Z = 4

F(000) = 707

D_x = 1.529 Mg m⁻³

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 13419 reflections

θ = 2.0–27.5°

μ = 1.04 mm⁻¹

T = 150 K

Plate, orange

0.40 × 0.22 × 0.10 mm

Data collection

Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ scans, and ω scans with κ offsets

Absorption correction: multi-scan

from symmetry-related measurements

(*SORTAV*; Blessing, 1995)

T_{min} = 0.740, *T_{max}* = 0.906

14609 measured reflections

3377 independent reflections

2817 reflections with *I* > 2 σ (*I*)

R_{int} = 0.063

θ_{\max} = 27.5°, θ_{\min} = 2.6°

h = -12→12

k = -20→20

l = -12→12

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.100$
 $S = 1.05$
 3377 reflections
 257 parameters
 391 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0444P)^2 + 1.0113P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Fe	0.78558 (3)	1.126971 (18)	0.20906 (3)	0.02654 (11)	
C1	0.6804 (2)	1.19504 (13)	0.3330 (2)	0.0321 (4)	
H1A	0.5807	1.1867	0.3418	0.039*	
C2	0.7297 (3)	1.25123 (14)	0.2403 (2)	0.0367 (5)	
H2A	0.6705	1.2887	0.1706	0.044*	
C3	0.8778 (3)	1.24367 (14)	0.2608 (2)	0.0372 (5)	
H3A	0.9403	1.2750	0.2080	0.045*	
C4	0.9215 (2)	1.18269 (14)	0.3677 (2)	0.0323 (4)	
H4A	1.0197	1.1642	0.4040	0.039*	
C5	0.7997 (2)	1.15270 (13)	0.4133 (2)	0.0281 (4)	
C6	0.7987 (2)	1.08701 (13)	0.5188 (2)	0.0279 (4)	
O7	0.90170 (15)	1.05702 (10)	0.59097 (17)	0.0360 (4)	
O8	0.66914 (15)	1.06384 (10)	0.53090 (16)	0.0329 (3)	
C9	0.6609 (3)	1.00237 (16)	0.6387 (3)	0.0401 (5)	
H9A	0.7121	0.9504	0.6229	0.060*	
H9B	0.5629	0.9880	0.6377	0.060*	
H9C	0.7022	1.0270	0.7285	0.060*	
C10	0.8967 (2)	1.03436 (15)	0.1286 (2)	0.0364 (5)	
H10A	0.9987	1.0223	0.1575	0.044*	
C11	0.8348 (3)	1.09292 (16)	0.0240 (2)	0.0390 (5)	
H11A	0.8860	1.1292	-0.0336	0.047*	
C12	0.6877 (2)	1.09098 (15)	0.0158 (2)	0.0366 (5)	
H12A	0.6178	1.1258	-0.0484	0.044*	
C13	0.6566 (2)	1.03129 (14)	0.1156 (2)	0.0326 (5)	
H13A	0.5618	1.0167	0.1336	0.039*	

C14	0.7866 (2)	0.99586 (13)	0.1840 (2)	0.0308 (4)	
N15A	0.8170 (15)	0.9348 (8)	0.2832 (15)	0.0298 (18)	0.60
H15A	0.9036	0.9299	0.3281	0.036*	0.60
C16A	0.7063 (14)	0.8747 (8)	0.3191 (11)	0.034 (2)	0.60
O17A	0.5884 (19)	0.8780 (12)	0.2670 (11)	0.039 (2)	0.60
O18A	0.7847 (3)	0.80918 (19)	0.3703 (3)	0.0288 (6)	0.60
C19A	0.6961 (4)	0.7395 (3)	0.4008 (5)	0.0297 (8)	0.60
H19A	0.6203	0.7287	0.3191	0.036*	0.60
C20A	0.6332 (6)	0.7639 (4)	0.5229 (6)	0.0466 (13)	0.60
H20A	0.7078	0.7762	0.6019	0.070*	0.60
H20B	0.5759	0.7166	0.5462	0.070*	0.60
H20C	0.5747	0.8149	0.5005	0.070*	0.60
C21A	0.7913 (5)	0.6619 (3)	0.4267 (7)	0.0431 (12)	0.60
H21A	0.8368	0.6530	0.3473	0.065*	0.60
H21B	0.7360	0.6113	0.4398	0.065*	0.60
H21C	0.8624	0.6715	0.5098	0.065*	0.60
N15B	0.797 (2)	0.9282 (13)	0.289 (2)	0.032 (3)	0.40
H15B	0.8683	0.9303	0.3582	0.038*	0.40
C16B	0.7224 (19)	0.8744 (13)	0.2867 (15)	0.024 (2)	0.40
O17B	0.595 (3)	0.8684 (18)	0.2333 (18)	0.039 (3)	0.40
O18B	0.7611 (6)	0.8251 (3)	0.4349 (6)	0.0426 (12)	0.40
C19B	0.6752 (9)	0.7564 (6)	0.4736 (12)	0.052 (2)	0.40
H19B	0.5754	0.7688	0.4374	0.062*	0.40
H19C	0.6887	0.7526	0.5755	0.062*	0.40
C20B	0.7132 (13)	0.6753 (6)	0.4178 (12)	0.078 (3)	0.40
H20D	0.6956	0.6784	0.3167	0.117*	0.40
H20E	0.6570	0.6292	0.4469	0.117*	0.40
H20F	0.8124	0.6638	0.4522	0.117*	0.40

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe	0.02407 (16)	0.03112 (17)	0.02215 (17)	0.00247 (11)	-0.00153 (11)	-0.00056 (11)
C1	0.0310 (10)	0.0354 (11)	0.0288 (11)	0.0068 (8)	0.0026 (9)	-0.0032 (8)
C2	0.0463 (13)	0.0294 (10)	0.0315 (12)	0.0075 (9)	-0.0005 (10)	-0.0002 (9)
C3	0.0430 (12)	0.0326 (11)	0.0338 (12)	-0.0065 (9)	0.0012 (10)	0.0016 (9)
C4	0.0280 (10)	0.0373 (11)	0.0286 (11)	-0.0048 (8)	-0.0024 (8)	-0.0038 (9)
C5	0.0295 (10)	0.0305 (9)	0.0222 (10)	0.0007 (8)	-0.0008 (8)	-0.0044 (8)
C6	0.0258 (10)	0.0331 (10)	0.0229 (10)	0.0007 (8)	-0.0004 (8)	-0.0056 (8)
O7	0.0279 (8)	0.0424 (9)	0.0335 (9)	0.0016 (6)	-0.0052 (6)	0.0066 (7)
O8	0.0255 (7)	0.0429 (8)	0.0287 (8)	-0.0008 (6)	0.0008 (6)	0.0046 (6)
C9	0.0379 (12)	0.0491 (13)	0.0325 (12)	-0.0051 (10)	0.0045 (10)	0.0053 (10)
C10	0.0332 (11)	0.0426 (12)	0.0324 (12)	0.0082 (9)	0.0042 (9)	-0.0071 (9)
C11	0.0418 (13)	0.0499 (13)	0.0255 (11)	0.0050 (10)	0.0066 (9)	-0.0041 (9)
C12	0.0390 (12)	0.0428 (12)	0.0228 (10)	0.0048 (9)	-0.0074 (9)	-0.0022 (9)
C13	0.0283 (10)	0.0349 (11)	0.0298 (11)	0.0012 (8)	-0.0072 (8)	-0.0048 (9)
C14	0.0284 (10)	0.0302 (10)	0.0291 (11)	0.0031 (8)	-0.0068 (9)	-0.0061 (8)
N15A	0.022 (4)	0.027 (2)	0.036 (3)	-0.010 (2)	-0.005 (3)	-0.003 (2)

C16A	0.033 (4)	0.027 (2)	0.040 (6)	-0.004 (2)	-0.004 (3)	0.003 (3)
O17A	0.027 (2)	0.045 (4)	0.040 (5)	-0.002 (2)	-0.011 (4)	0.000 (4)
O18A	0.0233 (13)	0.0283 (14)	0.0325 (17)	-0.0025 (10)	-0.0009 (13)	0.0076 (13)
C19A	0.032 (2)	0.028 (2)	0.027 (2)	-0.0064 (14)	0.0018 (17)	0.0043 (16)
C20A	0.044 (3)	0.056 (3)	0.044 (3)	-0.010 (2)	0.021 (2)	0.000 (2)
C21A	0.040 (2)	0.031 (2)	0.055 (3)	0.0009 (19)	0.000 (2)	0.0122 (19)
N15B	0.013 (5)	0.030 (5)	0.047 (5)	0.004 (3)	-0.012 (3)	0.008 (4)
C16B	0.017 (4)	0.037 (4)	0.017 (5)	0.002 (3)	0.001 (3)	0.006 (3)
O17B	0.031 (4)	0.048 (6)	0.032 (7)	-0.006 (3)	-0.008 (5)	-0.002 (5)
O18B	0.038 (3)	0.038 (3)	0.045 (3)	-0.007 (2)	-0.009 (2)	0.014 (2)
C19B	0.045 (5)	0.054 (5)	0.057 (7)	-0.010 (4)	0.012 (4)	0.023 (4)
C20B	0.115 (9)	0.047 (5)	0.069 (6)	-0.020 (6)	0.011 (7)	0.004 (4)

Geometric parameters (Å, °)

Fe—C5	2.033 (2)	C12—C13	1.429 (3)
Fe—C1	2.037 (2)	C12—H12A	1.0000
Fe—C11	2.042 (2)	C13—C14	1.431 (3)
Fe—C12	2.043 (2)	C13—H13A	1.0000
Fe—C4	2.045 (2)	C14—N15A	1.358 (13)
Fe—C10	2.054 (2)	C14—N15B	1.467 (18)
Fe—C3	2.055 (2)	N15A—C16A	1.522 (17)
Fe—C2	2.055 (2)	N15A—H15A	0.8800
Fe—C13	2.057 (2)	C16A—O17A	1.17 (2)
Fe—C14	2.064 (2)	C16A—O18A	1.320 (14)
C1—C2	1.415 (3)	O18A—C19A	1.455 (5)
C1—C5	1.439 (3)	C19A—C20A	1.500 (6)
C1—H1A	1.0000	C19A—C21A	1.518 (6)
C2—C3	1.425 (3)	C19A—H19A	1.0000
C2—H2A	1.0000	C20A—H20A	0.9800
C3—C4	1.425 (3)	C20A—H20B	0.9800
C3—H3A	1.0000	C20A—H20C	0.9800
C4—C5	1.426 (3)	C21A—H21A	0.9800
C4—H4A	1.0000	C21A—H21B	0.9800
C5—C6	1.463 (3)	C21A—H21C	0.9800
C6—O7	1.211 (2)	N15B—C16B	1.11 (3)
C6—O8	1.340 (2)	N15B—H15B	0.8800
O8—C9	1.446 (3)	C16B—O17B	1.26 (3)
C9—H9A	0.9800	C16B—O18B	1.632 (16)
C9—H9B	0.9800	O18B—C19B	1.455 (9)
C9—H9C	0.9800	C19B—C20B	1.457 (15)
C10—C11	1.425 (3)	C19B—H19B	0.9900
C10—C14	1.425 (3)	C19B—H19C	0.9900
C10—H10A	1.0000	C20B—H20D	0.9800
C11—C12	1.421 (3)	C20B—H20E	0.9800
C11—H11A	1.0000	C20B—H20F	0.9800
C5—Fe—C1	41.42 (8)	O7—C6—O8	122.3 (2)

C5—Fe—C11	162.38 (9)	O7—C6—C5	125.03 (19)
C1—Fe—C11	154.35 (9)	O8—C6—C5	112.62 (17)
C5—Fe—C12	155.67 (10)	C6—O8—C9	115.33 (17)
C1—Fe—C12	120.25 (9)	O8—C9—H9A	109.5
C11—Fe—C12	40.72 (10)	O8—C9—H9B	109.5
C5—Fe—C4	40.93 (9)	H9A—C9—H9B	109.5
C1—Fe—C4	69.25 (9)	O8—C9—H9C	109.5
C11—Fe—C4	124.55 (10)	H9A—C9—H9C	109.5
C12—Fe—C4	162.28 (10)	H9B—C9—H9C	109.5
C5—Fe—C10	125.42 (9)	C11—C10—C14	107.5 (2)
C1—Fe—C10	163.99 (9)	C11—C10—Fe	69.20 (13)
C11—Fe—C10	40.71 (9)	C14—C10—Fe	70.10 (12)
C12—Fe—C10	68.51 (9)	C11—C10—H10A	126.2
C4—Fe—C10	106.41 (9)	C14—C10—H10A	126.2
C5—Fe—C3	68.50 (9)	Fe—C10—H10A	126.2
C1—Fe—C3	68.52 (10)	C12—C11—C10	108.2 (2)
C11—Fe—C3	106.72 (10)	C12—C11—Fe	69.68 (13)
C12—Fe—C3	125.58 (10)	C10—C11—Fe	70.09 (13)
C4—Fe—C3	40.69 (9)	C12—C11—H11A	125.9
C10—Fe—C3	118.89 (10)	C10—C11—H11A	125.9
C5—Fe—C2	68.59 (9)	Fe—C11—H11A	125.9
C1—Fe—C2	40.46 (9)	C11—C12—C13	108.5 (2)
C11—Fe—C2	119.57 (10)	C11—C12—Fe	69.60 (13)
C12—Fe—C2	108.02 (9)	C13—C12—Fe	70.12 (12)
C4—Fe—C2	68.59 (9)	C11—C12—H12A	125.8
C10—Fe—C2	153.80 (10)	C13—C12—H12A	125.8
C3—Fe—C2	40.57 (9)	Fe—C12—H12A	125.8
C5—Fe—C13	120.63 (9)	C12—C13—C14	107.0 (2)
C1—Fe—C13	108.29 (9)	C12—C13—Fe	69.08 (12)
C11—Fe—C13	68.71 (10)	C14—C13—Fe	69.93 (12)
C12—Fe—C13	40.80 (9)	C12—C13—H13A	126.5
C4—Fe—C13	154.95 (9)	C14—C13—H13A	126.5
C10—Fe—C13	68.74 (9)	Fe—C13—H13A	126.5
C3—Fe—C13	163.45 (9)	N15A—C14—C10	119.6 (6)
C2—Fe—C13	126.64 (9)	N15A—C14—C13	131.7 (6)
C5—Fe—C14	108.30 (9)	C10—C14—C13	108.7 (2)
C1—Fe—C14	127.19 (9)	C10—C14—N15B	128.0 (8)
C11—Fe—C14	68.11 (10)	C13—C14—N15B	123.3 (8)
C12—Fe—C14	68.11 (9)	N15A—C14—Fe	128.0 (6)
C4—Fe—C14	119.81 (9)	C10—C14—Fe	69.39 (12)
C10—Fe—C14	40.51 (9)	C13—C14—Fe	69.43 (12)
C3—Fe—C14	154.02 (9)	N15B—C14—Fe	129.3 (10)
C2—Fe—C14	164.38 (9)	C14—N15A—C16A	122.2 (11)
C13—Fe—C14	40.65 (8)	C14—N15A—H15A	118.9
C2—C1—C5	107.62 (19)	C16A—N15A—H15A	118.9
C2—C1—Fe	70.47 (13)	O17A—C16A—O18A	130.7 (14)
C5—C1—Fe	69.14 (12)	O17A—C16A—N15A	123.6 (14)
C2—C1—H1A	126.2	O18A—C16A—N15A	100.6 (10)

C5—C1—H1A	126.2	C16A—O18A—C19A	109.7 (6)
Fe—C1—H1A	126.2	O18A—C19A—C20A	108.9 (4)
C1—C2—C3	108.39 (19)	O18A—C19A—C21A	105.3 (4)
C1—C2—Fe	69.07 (12)	C20A—C19A—C21A	113.4 (4)
C3—C2—Fe	69.68 (12)	O18A—C19A—H19A	109.7
C1—C2—H2A	125.8	C20A—C19A—H19A	109.7
C3—C2—H2A	125.8	C21A—C19A—H19A	109.7
Fe—C2—H2A	125.8	C16B—N15B—C14	125.1 (19)
C2—C3—C4	108.3 (2)	C16B—N15B—H15B	117.5
C2—C3—Fe	69.74 (12)	C14—N15B—H15B	117.5
C4—C3—Fe	69.29 (12)	N15B—C16B—O17B	131 (2)
C2—C3—H3A	125.8	N15B—C16B—O18B	107.6 (17)
C4—C3—H3A	125.8	O17B—C16B—O18B	112.6 (15)
Fe—C3—H3A	125.8	C19B—O18B—C16B	122.3 (9)
C3—C4—C5	107.58 (19)	O18B—C19B—C20B	110.0 (9)
C3—C4—Fe	70.02 (13)	O18B—C19B—H19B	109.7
C5—C4—Fe	69.08 (12)	C20B—C19B—H19B	109.7
C3—C4—H4A	126.2	O18B—C19B—H19C	109.7
C5—C4—H4A	126.2	C20B—C19B—H19C	109.7
Fe—C4—H4A	126.2	H19B—C19B—H19C	108.2
C4—C5—C1	108.09 (19)	C19B—C20B—H20D	109.5
C4—C5—C6	124.89 (19)	C19B—C20B—H20E	109.5
C1—C5—C6	126.95 (19)	H20D—C20B—H20E	109.5
C4—C5—Fe	69.99 (12)	C19B—C20B—H20F	109.5
C1—C5—Fe	69.44 (12)	H20D—C20B—H20F	109.5
C6—C5—Fe	123.90 (14)	H20E—C20B—H20F	109.5
C5—Fe—C1—C2	118.61 (18)	C4—Fe—C10—C14	116.94 (13)
C11—Fe—C1—C2	-46.8 (3)	C3—Fe—C10—C14	159.18 (12)
C12—Fe—C1—C2	-82.43 (16)	C2—Fe—C10—C14	-168.33 (18)
C4—Fe—C1—C2	80.95 (14)	C13—Fe—C10—C14	-37.06 (13)
C10—Fe—C1—C2	157.9 (3)	C14—C10—C11—C12	0.4 (3)
C3—Fe—C1—C2	37.20 (13)	Fe—C10—C11—C12	-59.44 (16)
C13—Fe—C1—C2	-125.52 (14)	C14—C10—C11—Fe	59.87 (15)
C14—Fe—C1—C2	-166.73 (13)	C5—Fe—C11—C12	163.0 (3)
C11—Fe—C1—C5	-165.4 (2)	C1—Fe—C11—C12	-50.5 (3)
C12—Fe—C1—C5	158.97 (13)	C4—Fe—C11—C12	-166.63 (13)
C4—Fe—C1—C5	-37.65 (12)	C10—Fe—C11—C12	119.3 (2)
C10—Fe—C1—C5	39.3 (4)	C3—Fe—C11—C12	-125.60 (15)
C3—Fe—C1—C5	-81.41 (14)	C2—Fe—C11—C12	-83.45 (16)
C2—Fe—C1—C5	-118.61 (18)	C13—Fe—C11—C12	37.56 (13)
C13—Fe—C1—C5	115.87 (13)	C14—Fe—C11—C12	81.42 (15)
C14—Fe—C1—C5	74.67 (15)	C5—Fe—C11—C10	43.7 (4)
C5—C1—C2—C3	0.7 (2)	C1—Fe—C11—C10	-169.82 (19)
Fe—C1—C2—C3	-58.74 (16)	C12—Fe—C11—C10	-119.3 (2)
C5—C1—C2—Fe	59.40 (14)	C4—Fe—C11—C10	74.08 (17)
C5—Fe—C2—C1	-38.60 (13)	C3—Fe—C11—C10	115.10 (15)
C11—Fe—C2—C1	158.74 (13)	C2—Fe—C11—C10	157.25 (14)

C12—Fe—C2—C1	115.77 (14)	C13—Fe—C11—C10	-81.73 (15)
C4—Fe—C2—C1	-82.72 (14)	C14—Fe—C11—C10	-37.87 (14)
C10—Fe—C2—C1	-166.43 (19)	C10—C11—C12—C13	0.2 (3)
C3—Fe—C2—C1	-120.12 (19)	Fe—C11—C12—C13	-59.54 (16)
C13—Fe—C2—C1	74.38 (16)	C10—C11—C12—Fe	59.71 (16)
C14—Fe—C2—C1	42.8 (4)	C5—Fe—C12—C11	-167.56 (19)
C5—Fe—C2—C3	81.52 (14)	C1—Fe—C12—C11	157.25 (14)
C1—Fe—C2—C3	120.12 (19)	C4—Fe—C12—C11	38.7 (4)
C11—Fe—C2—C3	-81.14 (16)	C10—Fe—C12—C11	-37.68 (14)
C12—Fe—C2—C3	-124.11 (14)	C3—Fe—C12—C11	73.22 (17)
C4—Fe—C2—C3	37.40 (14)	C2—Fe—C12—C11	114.68 (15)
C10—Fe—C2—C3	-46.3 (3)	C13—Fe—C12—C11	-119.62 (19)
C13—Fe—C2—C3	-165.50 (14)	C14—Fe—C12—C11	-81.43 (15)
C14—Fe—C2—C3	162.9 (3)	C5—Fe—C12—C13	-47.9 (3)
C1—C2—C3—C4	-0.4 (3)	C1—Fe—C12—C13	-83.13 (15)
Fe—C2—C3—C4	-58.72 (15)	C11—Fe—C12—C13	119.62 (19)
C1—C2—C3—Fe	58.36 (15)	C4—Fe—C12—C13	158.4 (3)
C5—Fe—C3—C2	-81.77 (15)	C10—Fe—C12—C13	81.94 (14)
C1—Fe—C3—C2	-37.10 (14)	C3—Fe—C12—C13	-167.16 (13)
C11—Fe—C3—C2	116.20 (15)	C2—Fe—C12—C13	-125.70 (14)
C12—Fe—C3—C2	75.49 (17)	C14—Fe—C12—C13	38.19 (13)
C4—Fe—C3—C2	-119.8 (2)	C11—C12—C13—C14	-0.7 (2)
C10—Fe—C3—C2	158.62 (14)	Fe—C12—C13—C14	-59.91 (15)
C13—Fe—C3—C2	44.9 (4)	C11—C12—C13—Fe	59.22 (16)
C14—Fe—C3—C2	-169.58 (19)	C5—Fe—C13—C12	159.17 (13)
C5—Fe—C3—C4	38.06 (13)	C1—Fe—C13—C12	115.40 (14)
C1—Fe—C3—C4	82.74 (14)	C11—Fe—C13—C12	-37.49 (14)
C11—Fe—C3—C4	-123.97 (14)	C4—Fe—C13—C12	-164.6 (2)
C12—Fe—C3—C4	-164.67 (13)	C10—Fe—C13—C12	-81.33 (15)
C10—Fe—C3—C4	-81.55 (16)	C3—Fe—C13—C12	39.4 (4)
C2—Fe—C3—C4	119.8 (2)	C2—Fe—C13—C12	74.24 (17)
C13—Fe—C3—C4	164.7 (3)	C14—Fe—C13—C12	-118.27 (19)
C14—Fe—C3—C4	-49.7 (3)	C5—Fe—C13—C14	-82.56 (15)
C2—C3—C4—C5	-0.1 (2)	C1—Fe—C13—C14	-126.33 (14)
Fe—C3—C4—C5	-59.08 (14)	C11—Fe—C13—C14	80.78 (15)
C2—C3—C4—Fe	59.00 (15)	C12—Fe—C13—C14	118.27 (19)
C5—Fe—C4—C3	-118.89 (19)	C4—Fe—C13—C14	-46.4 (3)
C1—Fe—C4—C3	-80.80 (15)	C10—Fe—C13—C14	36.94 (14)
C11—Fe—C4—C3	74.64 (17)	C3—Fe—C13—C14	157.6 (3)
C12—Fe—C4—C3	44.9 (3)	C2—Fe—C13—C14	-167.49 (13)
C10—Fe—C4—C3	115.47 (15)	C11—C10—C14—N15A	177.8 (7)
C2—Fe—C4—C3	-37.30 (14)	Fe—C10—C14—N15A	-122.9 (7)
C13—Fe—C4—C3	-169.8 (2)	C11—C10—C14—C13	-0.9 (2)
C14—Fe—C4—C3	157.33 (14)	Fe—C10—C14—C13	58.44 (15)
C1—Fe—C4—C5	38.09 (12)	C11—C10—C14—N15B	176.4 (12)
C11—Fe—C4—C5	-166.47 (13)	Fe—C10—C14—N15B	-124.3 (13)
C12—Fe—C4—C5	163.8 (3)	C11—C10—C14—Fe	-59.30 (15)
C10—Fe—C4—C5	-125.64 (13)	C12—C13—C14—N15A	-177.5 (9)

C3—Fe—C4—C5	118.89 (19)	Fe—C13—C14—N15A	123.2 (9)
C2—Fe—C4—C5	81.58 (14)	C12—C13—C14—C10	0.9 (2)
C13—Fe—C4—C5	-50.9 (3)	Fe—C13—C14—C10	-58.42 (15)
C14—Fe—C4—C5	-83.78 (14)	C12—C13—C14—N15B	-176.4 (12)
C3—C4—C5—C1	0.5 (2)	Fe—C13—C14—N15B	124.2 (12)
Fe—C4—C5—C1	-59.19 (14)	C12—C13—C14—Fe	59.37 (15)
C3—C4—C5—C6	177.69 (19)	C5—Fe—C14—N15A	-11.5 (8)
Fe—C4—C5—C6	118.0 (2)	C1—Fe—C14—N15A	-53.7 (8)
C3—C4—C5—Fe	59.68 (15)	C11—Fe—C14—N15A	150.1 (8)
C2—C1—C5—C4	-0.7 (2)	C12—Fe—C14—N15A	-165.8 (8)
Fe—C1—C5—C4	59.53 (15)	C4—Fe—C14—N15A	31.8 (8)
C2—C1—C5—C6	-177.8 (2)	C10—Fe—C14—N15A	112.1 (8)
Fe—C1—C5—C6	-117.6 (2)	C3—Fe—C14—N15A	66.8 (8)
C2—C1—C5—Fe	-60.24 (15)	C2—Fe—C14—N15A	-87.3 (8)
C1—Fe—C5—C4	-119.31 (18)	C13—Fe—C14—N15A	-127.5 (8)
C11—Fe—C5—C4	39.5 (4)	C5—Fe—C14—C10	-123.56 (14)
C12—Fe—C5—C4	-168.1 (2)	C1—Fe—C14—C10	-165.78 (13)
C10—Fe—C5—C4	73.07 (16)	C11—Fe—C14—C10	38.06 (14)
C3—Fe—C5—C4	-37.84 (13)	C12—Fe—C14—C10	82.10 (15)
C2—Fe—C5—C4	-81.59 (14)	C4—Fe—C14—C10	-80.25 (15)
C13—Fe—C5—C4	157.56 (12)	C3—Fe—C14—C10	-45.3 (3)
C14—Fe—C5—C4	114.70 (13)	C2—Fe—C14—C10	160.6 (3)
C11—Fe—C5—C1	158.8 (3)	C13—Fe—C14—C10	120.42 (19)
C12—Fe—C5—C1	-48.8 (3)	C5—Fe—C14—C13	116.01 (14)
C4—Fe—C5—C1	119.31 (18)	C1—Fe—C14—C13	73.79 (16)
C10—Fe—C5—C1	-167.61 (13)	C11—Fe—C14—C13	-82.37 (15)
C3—Fe—C5—C1	81.47 (14)	C12—Fe—C14—C13	-38.33 (14)
C2—Fe—C5—C1	37.73 (13)	C4—Fe—C14—C13	159.33 (13)
C13—Fe—C5—C1	-83.12 (14)	C10—Fe—C14—C13	-120.42 (19)
C14—Fe—C5—C1	-125.99 (13)	C3—Fe—C14—C13	-165.7 (2)
C1—Fe—C5—C6	121.4 (2)	C2—Fe—C14—C13	40.2 (4)
C11—Fe—C5—C6	-79.7 (4)	C5—Fe—C14—N15B	-0.8 (10)
C12—Fe—C5—C6	72.6 (3)	C1—Fe—C14—N15B	-43.0 (10)
C4—Fe—C5—C6	-119.3 (2)	C11—Fe—C14—N15B	160.8 (10)
C10—Fe—C5—C6	-46.2 (2)	C12—Fe—C14—N15B	-155.1 (10)
C3—Fe—C5—C6	-157.1 (2)	C4—Fe—C14—N15B	42.5 (10)
C2—Fe—C5—C6	159.2 (2)	C10—Fe—C14—N15B	122.8 (10)
C13—Fe—C5—C6	38.3 (2)	C3—Fe—C14—N15B	77.5 (10)
C14—Fe—C5—C6	-4.56 (19)	C2—Fe—C14—N15B	-76.6 (11)
C4—C5—C6—O7	8.7 (3)	C13—Fe—C14—N15B	-116.8 (10)
C1—C5—C6—O7	-174.7 (2)	C10—C14—N15A—C16A	-163.1 (9)
Fe—C5—C6—O7	96.7 (2)	C13—C14—N15A—C16A	15.1 (18)
C4—C5—C6—O8	-172.83 (19)	N15B—C14—N15A—C16A	9 (10)
C1—C5—C6—O8	3.8 (3)	Fe—C14—N15A—C16A	110.8 (11)
Fe—C5—C6—O8	-84.8 (2)	C14—N15A—C16A—O17A	-2 (2)
O7—C6—O8—C9	1.7 (3)	C14—N15A—C16A—O18A	155.1 (11)
C5—C6—O8—C9	-176.83 (18)	O17A—C16A—O18A—C19A	-20.4 (14)
C5—Fe—C10—C11	-165.14 (14)	N15A—C16A—O18A—C19A	-174.9 (7)

C1—Fe—C10—C11	163.9 (3)	C16A—O18A—C19A—C20A	-70.8 (7)
C12—Fe—C10—C11	37.69 (15)	C16A—O18A—C19A—C21A	167.3 (6)
C4—Fe—C10—C11	-124.34 (15)	N15A—C14—N15B—C16B	-149 (13)
C3—Fe—C10—C11	-82.10 (16)	C10—C14—N15B—C16B	-140.6 (19)
C2—Fe—C10—C11	-49.6 (3)	C13—C14—N15B—C16B	36 (3)
C13—Fe—C10—C11	81.65 (15)	Fe—C14—N15B—C16B	126 (2)
C14—Fe—C10—C11	118.7 (2)	C14—N15B—C16B—O17B	-27 (4)
C5—Fe—C10—C14	76.14 (15)	C14—N15B—C16B—O18B	-170.7 (18)
C1—Fe—C10—C14	45.2 (4)	N15B—C16B—O18B—C19B	173.6 (16)
C11—Fe—C10—C14	-118.7 (2)	O17B—C16B—O18B—C19B	22 (2)
C12—Fe—C10—C14	-81.02 (14)	C16B—O18B—C19B—C20B	84.4 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N15A—H15A...O7 ⁱ	0.88	1.93	2.793 (14)	168
N15B—H15B...O7 ⁱ	0.88	2.21	2.962 (19)	143

Symmetry code: (i) $-x+2, -y+2, -z+1$.