

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetra(chlorido/iodido)(1,10-phenanthroline)platinum(IV) hemi[di(chlorine/iodine)]

Nam-Ho Kim and Kwang Ha*

 School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
 Correspondence e-mail: hakwang@chonnam.ac.kr


Received 14 February 2009; accepted 3 March 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; disorder in main residue; R factor = 0.031; wR factor = 0.062; data-to-parameter ratio = 16.7.

The asymmetric unit of the title compound, $[\text{PtCl}_{3.66}\text{I}_{0.34}(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5(\text{Cl}_{0.13}\text{I}_{1.87})$, contains a neutral Pt^{IV} complex and one half of a halogen molecule. The Pt^{IV} ion is six-coordinated in a distorted octahedral environment by two N atoms of the 1,10-phenanthroline ligand and Cl or I atoms. The refinement of the structure and the EDX analysis indicate that the compound is a solid solution in which there is some substitution of Cl for I and *vice versa*. The chemical formula of the pure state of the compound would have been $[\text{PtCl}_4(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5\text{I}_2$. In the analysed crystal, two Cl atoms are partially (*ca* 25% and 9%) replaced by I atoms, and the I_2 molecule has a minor component modelled as ICl. As a result of the disorder, the different *trans* effects of the N and Cl/I atoms are not distinct. The complex displays intermolecular π - π interactions between the six-membered rings, with a centroid-centroid distance of 3.771 (4) Å. There are also weak intramolecular $\text{C}-\text{H} \cdots \text{Cl}$ hydrogen bonds.

Related literature

For details of some other Pt-phenanthroline complexes, see: Buse *et al.* (1977); Fanizzi *et al.* (1996); Kim *et al.* (2009*a,b*). For related Pt-bipyridine complexes, see: Hambley (1986); Hojjat Kashani *et al.* (2008). For bond-length data, see: Orpen *et al.* (1989).



Experimental

Crystal data

$[\text{PtCl}_{3.66}\text{I}_{0.34}(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5(\text{Cl}_{0.13}\text{I}_{1.87})$
 $M_r = 669.26$
 Orthorhombic, *Pbca*
 $a = 14.215$ (5) Å
 $b = 12.733$ (5) Å
 $c = 17.575$ (6) Å
 $V = 3180.8$ (19) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 11.92$ mm⁻¹
 $T = 293$ K
 $0.25 \times 0.17 \times 0.15$ mm

Data collection

Bruker SMART 1000 CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.111$, $T_{\max} = 0.168$
 17485 measured reflections
 3246 independent reflections
 2169 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.062$
 $S = 0.86$
 3246 reflections
 194 parameters
 8 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.29$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.50$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{Cl1}-\text{H1} \cdots \text{Cl2}$	0.93	2.73	3.320 (8)	122
$\text{Cl10}-\text{H10} \cdots \text{Cl1}$	0.93	2.66	3.240 (7)	121

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by a Korea Research Foundation grant funded by the Korean Government (MOEHRD) (KRF-2007-412-J02001).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2156).

References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Buse, K. D., Keller, H. J. & Pritzkow, H. (1977). *Inorg. Chem.* **16**, 1072–1076.
 Fanizzi, F. P., Natile, G., Lanfranchi, M., Tiripicchio, A., Laschi, F. & Zanello, P. (1996). *Inorg. Chem.* **35**, 3173–3182.
 Hambley, T. W. (1986). *Acta Cryst.* **C42**, 49–51.
 Hojjat Kashani, L., Amani, V., Yousefi, M. & Khavasi, H. R. (2008). *Acta Cryst.* **E64**, m905–m906.
 Kim, N.-H., Hwang, I.-C. & Ha, K. (2009*a*). *Acta Cryst.* **E65**, m224.
 Kim, N.-H., Hwang, I.-C. & Ha, K. (2009*b*). *Acta Cryst.* **E65**, m230.
 Orpen, G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1989). *J. Chem. Soc. Dalton Trans.* pp. S1–83.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2009). E65, m379 [doi:10.1107/S1600536809007703]

Tetra(chlorido/iodido)(1,10-phenanthroline)platinum(IV) hemi[di(chlorine/iodine)]

Nam-Ho Kim and Kwang Ha

S1. Comment

The asymmetric unit of the title compound contains a neutral Pt^{IV} complex and one half-molecule of iodine which includes some Cl atoms (*ca* 6%). The Pt^{IV} ion is six-coordinated in a distorted octahedral environment by two N atoms of the 1,10-phenanthroline ligand and Cl or I atoms. The chemical formula of the pure state of the title compound would have been [PtCl₄(C₁₂H₈N₂)]·0.5I₂. In the particular crystal of the compound used, two Cl atoms (Cl3 and Cl4) are partially (*ca* 25% and 9%, respectively) displaced by the I atoms (I3 and I4) through the substitution reaction between the Cl⁻ and I⁻ ligand, and the I₂ molecule also appears to have a minor component, that is I—Cl (Fig. 1 and 2). The chemical formula which resulted from the refinement of the structure was [PtCl_{3.66}I_{0.34}(C₁₂H₈N₂)]·0.5(Cl_{0.13}I_{1.87}), and in this case the ratio of the Cl atom to I atom is 2.91:1. An EDX analysis of the compound, however, gave a ratio of Cl:I = 2.47:1. Accordingly, the exact composition may very well be variable, and likely dependent on the exact conditions present during crystal formation. Even though these data are slightly different, they indicate clearly that the crystals are a solid solution in which there was some substitution of Cl for I and *vice versa*.

As a result of the different *trans* effects of the N and Cl atoms, the Pt—Cl bonds *trans* to the N atom are in general slightly shorter than bond lengths to mutually *trans* Cl atoms (Kim *et al.* 2009*a* and 2009*b*). But the *trans* effects of the N and Cl/I atoms in the crystal are not distinct owing to the disordered atoms. The Pt—I distance is restrained to the value given in table 9.6.3.3 of the International Tables Vol. C (Orpen *et al.*, 1989) (2.658 Å). The main contributor to the distortion from a true octahedral structure is the tight N1—Pt1—N2 chelate angle (81.3 (2)°), which result in non-linear *trans* axes (<Cl1—Pt1—N1 = 174.14 (16)° and <Cl2—Pt1—N2 = 175.97 (17)°). The complex displays intermolecular π - π interactions between the six-membered rings, with a shortest centroid-centroid distance of 3.771 (4) Å and with a dihedral angle between the ring planes of 2.1 (3)°. There are also weak intramolecular C—H...Cl hydrogen bonds (Table 1).

The iodine molecule was presumed formed as a consequence of the oxidation of the iodide ion by the Pt⁴⁺ ion, and crystallized with the partially substituted complex. The bond distance between the I atoms is 2.708 (2) Å.

S2. Experimental

To a solution of [PtCl₄(C₁₂H₈N₂)]·H₂O (0.0821 g, 0.153 mmol) in H₂O (20 ml) was added KI (0.1318 g, 0.794 mmol), and stirred for 2 h at room temperature. The precipitate was separated by filtration and washed with water (20 ml) and MeOH (20 ml) and dried under vacuum, to give a dark brown powder (0.0846 g). Black crystals suitable for X-ray analysis were isolated from an acetone solution of the reaction products. EDX analysis (%atom): Cl 62.30%, I 25.18%, Pt 12.52%.

S3. Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [$C-H = 0.93 \text{ \AA}$ and $U_{iso}(H) = 1.2U_{eq}(C)$]. The disordered Cl5 atom was refined isotropically. Eight restraints instructions were used for the refinement using the following *SHELXL97* (Sheldrick, 2008) commands: EADP C13 I3 and C14 I4, SIMU 0.010 I1 Cl5, BIND I1 Cl5a and C15 I1a, FREE C15 Cl5a, *DFIX* 2.658 0.010 Pt1 I3 and Pt1 I4.




Figure 1

The disordered structure of the title compound, with displacement ellipsoids drawn at the 30% probability level for non-H atoms [Symmetry code: (a) $1 - x, 1 - y, -z$]. The minor bonds are drawn with dashed lines.




Figure 2

View of a packing detail of the title compound. For the sake of clarity, only the major disorder component is shown.

Chloridoiodido(1,10-phenanthroline)platinum(IV) hemi[di(chloride/iodide)]

Crystal data

$[\text{PtCl}_{3.66}\text{I}_{0.34}(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5(\text{Cl}_{0.13}\text{I}_{1.87})$

$M_r = 669.26$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 14.215 (5) \text{ \AA}$

$b = 12.733 (5) \text{ \AA}$

$c = 17.575 (6) \text{ \AA}$

$V = 3180.8 (19) \text{ \AA}^3$

$Z = 8$

$F(000) = 2424$

$D_x = 2.795 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 973 reflections

$\theta = 2.4\text{--}24.5^\circ$

$\mu = 11.92 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Plate, black

$0.25 \times 0.17 \times 0.15 \text{ mm}$

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube
Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)

$T_{\min} = 0.111$, $T_{\max} = 0.168$

17485 measured reflections

3246 independent reflections

2169 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.049$
 $\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.3^\circ$
 $h = -15 \rightarrow 17$

$k = -10 \rightarrow 15$
 $l = -17 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.062$
 $S = 0.86$
 3246 reflections
 194 parameters
 8 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0244P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 1.29 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.50 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Pt1	-0.134096 (18)	0.30394 (2)	0.177609 (15)	0.03991 (10)	
Cl1	-0.17047 (13)	0.14177 (15)	0.22685 (10)	0.0521 (5)	
Cl2	-0.28869 (12)	0.36077 (15)	0.19550 (10)	0.0512 (5)	
Cl3	-0.0903 (14)	0.3607 (17)	0.2949 (8)	0.0494 (13)	0.746 (3)
I3	-0.0759 (11)	0.3695 (13)	0.3094 (6)	0.0494 (13)	0.254 (3)
Cl4	-0.1730 (7)	0.2478 (8)	0.0572 (3)	0.0518 (8)	0.913 (3)
I4	-0.177 (2)	0.238 (2)	0.0398 (9)	0.0518 (8)	0.087 (3)
N1	-0.0875 (4)	0.4446 (4)	0.1355 (3)	0.0393 (14)	
N2	0.0029 (4)	0.2642 (4)	0.1585 (3)	0.0388 (13)	
C1	-0.1341 (5)	0.5328 (6)	0.1283 (4)	0.0517 (19)	
H1	-0.1978	0.5341	0.1403	0.062*	
C2	-0.0911 (6)	0.6243 (6)	0.1031 (4)	0.058 (2)	
H2	-0.1256	0.6863	0.0999	0.070*	
C3	0.0015 (6)	0.6235 (6)	0.0831 (4)	0.055 (2)	
H3	0.0305	0.6846	0.0660	0.066*	
C4	0.0535 (5)	0.5285 (6)	0.0887 (4)	0.0438 (18)	
C5	0.1496 (5)	0.5174 (6)	0.0688 (4)	0.0484 (19)	
H5	0.1820	0.5741	0.0481	0.058*	
C6	0.1946 (5)	0.4255 (6)	0.0796 (4)	0.050 (2)	
H6	0.2581	0.4210	0.0672	0.061*	
C7	0.1485 (5)	0.3347 (5)	0.1093 (4)	0.0403 (17)	
C8	0.1899 (5)	0.2382 (6)	0.1246 (4)	0.052 (2)	

H8	0.2534	0.2282	0.1141	0.062*	
C9	0.1393 (5)	0.1590 (6)	0.1546 (4)	0.0494 (19)	
H9	0.1678	0.0947	0.1647	0.059*	
C10	0.0446 (5)	0.1735 (5)	0.1703 (4)	0.0447 (18)	
H10	0.0098	0.1177	0.1897	0.054*	
C11	0.0538 (5)	0.3460 (6)	0.1273 (4)	0.0384 (17)	
C12	0.0055 (5)	0.4415 (5)	0.1167 (3)	0.0368 (16)	
I1	0.56522 (7)	0.45323 (7)	0.04476 (5)	0.0912 (4)	0.936 (3)
Cl5	0.558 (3)	0.496 (4)	0.021 (2)	0.062 (9)*	0.064 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.04138 (16)	0.03167 (16)	0.04668 (17)	-0.00138 (13)	0.00129 (14)	0.00057 (14)
Cl1	0.0561 (11)	0.0392 (11)	0.0608 (12)	-0.0060 (9)	0.0047 (9)	0.0072 (9)
Cl2	0.0490 (11)	0.0431 (12)	0.0616 (12)	0.0050 (9)	0.0083 (9)	0.0040 (9)
Cl3	0.064 (4)	0.045 (3)	0.039 (4)	-0.005 (2)	-0.007 (2)	-0.014 (3)
I3	0.064 (4)	0.045 (3)	0.039 (4)	-0.005 (2)	-0.007 (2)	-0.014 (3)
Cl4	0.0649 (15)	0.052 (2)	0.038 (3)	-0.0073 (14)	-0.006 (3)	-0.004 (2)
I4	0.0649 (15)	0.052 (2)	0.038 (3)	-0.0073 (14)	-0.006 (3)	-0.004 (2)
N1	0.044 (4)	0.029 (4)	0.045 (3)	-0.001 (3)	-0.001 (3)	-0.002 (3)
N2	0.037 (3)	0.032 (3)	0.046 (3)	0.000 (3)	-0.005 (3)	-0.001 (3)
C1	0.053 (5)	0.048 (5)	0.055 (5)	0.001 (4)	-0.002 (4)	-0.007 (4)
C2	0.057 (5)	0.031 (5)	0.086 (6)	0.006 (4)	-0.009 (5)	-0.004 (4)
C3	0.071 (6)	0.034 (5)	0.060 (5)	-0.011 (4)	-0.003 (4)	0.003 (4)
C4	0.052 (5)	0.036 (5)	0.043 (4)	-0.006 (4)	-0.007 (4)	-0.003 (3)
C5	0.050 (5)	0.044 (5)	0.051 (5)	-0.017 (4)	-0.006 (4)	0.007 (4)
C6	0.045 (5)	0.052 (6)	0.054 (5)	-0.012 (4)	0.006 (4)	-0.011 (4)
C7	0.044 (5)	0.036 (4)	0.041 (4)	-0.001 (3)	-0.002 (3)	-0.006 (3)
C8	0.046 (5)	0.053 (5)	0.056 (5)	0.007 (4)	0.006 (4)	-0.004 (4)
C9	0.047 (5)	0.046 (5)	0.055 (5)	0.009 (4)	-0.007 (4)	-0.010 (4)
C10	0.050 (5)	0.032 (5)	0.052 (5)	-0.006 (3)	0.002 (4)	-0.002 (3)
C11	0.040 (4)	0.037 (4)	0.038 (4)	-0.007 (3)	-0.008 (3)	-0.001 (3)
C12	0.042 (4)	0.034 (4)	0.034 (4)	-0.002 (3)	-0.004 (3)	0.002 (3)
I1	0.1295 (7)	0.0635 (6)	0.0805 (6)	-0.0192 (5)	0.0312 (5)	-0.0058 (4)

Geometric parameters (Å, °)

Pt1—N2	2.040 (5)	C4—C12	1.391 (9)
Pt1—N1	2.048 (5)	C4—C5	1.417 (9)
Pt1—Cl3	2.272 (9)	C5—C6	1.346 (9)
Pt1—Cl1	2.2977 (19)	C5—H5	0.9300
Pt1—Cl4	2.301 (4)	C6—C7	1.428 (9)
Pt1—Cl2	2.3347 (19)	C6—H6	0.9300
Pt1—I3	2.598 (7)	C7—C8	1.389 (9)
Pt1—I4	2.635 (9)	C7—C11	1.391 (8)
N1—C1	1.309 (8)	C8—C9	1.346 (10)
N1—C12	1.364 (8)	C8—H8	0.9300

N2—C10	1.316 (8)	C9—C10	1.387 (9)
N2—C11	1.381 (8)	C9—H9	0.9300
C1—C2	1.388 (10)	C10—H10	0.9300
C1—H1	0.9300	C11—C12	1.408 (9)
C2—C3	1.364 (10)	I1—C15 ⁱ	2.19 (4)
C2—H2	0.9300	I1—I1 ⁱ	2.708 (2)
C3—C4	1.421 (10)	C15—I1 ⁱ	2.19 (4)
C3—H3	0.9300		
N2—Pt1—N1	81.3 (2)	N1—C1—H1	119.0
N2—Pt1—C13	88.1 (6)	C2—C1—H1	119.0
N1—Pt1—C13	87.8 (6)	C3—C2—C1	120.0 (7)
N2—Pt1—C11	93.11 (17)	C3—C2—H2	120.0
N1—Pt1—C11	174.14 (16)	C1—C2—H2	120.0
C13—Pt1—C11	90.3 (6)	C2—C3—C4	119.4 (7)
N2—Pt1—C14	90.0 (3)	C2—C3—H3	120.3
N1—Pt1—C14	91.0 (3)	C4—C3—H3	120.3
C13—Pt1—C14	177.9 (6)	C12—C4—C5	118.7 (7)
C11—Pt1—C14	90.8 (3)	C12—C4—C3	116.6 (7)
N2—Pt1—C12	175.97 (17)	C5—C4—C3	124.7 (7)
N1—Pt1—C12	94.69 (16)	C6—C5—C4	120.7 (7)
C13—Pt1—C12	92.1 (5)	C6—C5—H5	119.7
C11—Pt1—C12	90.91 (7)	C4—C5—H5	119.7
C14—Pt1—C12	89.7 (3)	C5—C6—C7	122.5 (7)
N2—Pt1—I3	85.6 (4)	C5—C6—H6	118.8
N1—Pt1—I3	86.5 (4)	C7—C6—H6	118.8
C13—Pt1—I3	2.7 (9)	C8—C7—C11	117.2 (7)
C11—Pt1—I3	91.4 (4)	C8—C7—C6	126.4 (7)
C14—Pt1—I3	175.2 (4)	C11—C7—C6	116.3 (6)
C12—Pt1—I3	94.6 (4)	C9—C8—C7	120.8 (7)
N2—Pt1—I4	89.4 (8)	C9—C8—H8	119.6
N1—Pt1—I4	91.3 (7)	C7—C8—H8	119.6
C13—Pt1—I4	177.5 (9)	C8—C9—C10	119.8 (7)
C11—Pt1—I4	90.4 (7)	C8—C9—H9	120.1
C14—Pt1—I4	0.7 (10)	C10—C9—H9	120.1
C12—Pt1—I4	90.3 (7)	N2—C10—C9	121.5 (7)
I3—Pt1—I4	174.8 (8)	N2—C10—H10	119.2
C1—N1—C12	119.5 (6)	C9—C10—H10	119.2
C1—N1—Pt1	128.4 (5)	N2—C11—C7	121.3 (6)
C12—N1—Pt1	112.1 (4)	N2—C11—C12	116.6 (6)
C10—N2—C11	119.3 (6)	C7—C11—C12	122.0 (6)
C10—N2—Pt1	128.5 (5)	N1—C12—C4	122.5 (6)
C11—N2—Pt1	112.2 (4)	N1—C12—C11	117.7 (6)
N1—C1—C2	121.9 (7)	C4—C12—C11	119.7 (6)
N2—Pt1—N1—C1	-176.8 (6)	C3—C4—C5—C6	176.6 (7)
C13—Pt1—N1—C1	-88.4 (8)	C4—C5—C6—C7	1.6 (11)
C14—Pt1—N1—C1	93.3 (6)	C5—C6—C7—C8	-177.6 (7)

C12—Pt1—N1—C1	3.6 (6)	C5—C6—C7—C11	-0.4 (10)
I3—Pt1—N1—C1	-90.8 (7)	C11—C7—C8—C9	1.1 (10)
I4—Pt1—N1—C1	93.9 (9)	C6—C7—C8—C9	178.3 (7)
N2—Pt1—N1—C12	0.2 (4)	C7—C8—C9—C10	0.1 (11)
Cl3—Pt1—N1—C12	88.7 (7)	C11—N2—C10—C9	2.1 (9)
Cl4—Pt1—N1—C12	-89.7 (5)	Pt1—N2—C10—C9	-179.7 (5)
C12—Pt1—N1—C12	-179.4 (4)	C8—C9—C10—N2	-1.7 (10)
I3—Pt1—N1—C12	86.3 (5)	C10—N2—C11—C7	-0.9 (9)
I4—Pt1—N1—C12	-89.0 (8)	Pt1—N2—C11—C7	-179.4 (5)
N1—Pt1—N2—C10	-179.6 (6)	C10—N2—C11—C12	-179.4 (6)
Cl3—Pt1—N2—C10	92.4 (8)	Pt1—N2—C11—C12	2.1 (7)
Cl1—Pt1—N2—C10	2.2 (5)	C8—C7—C11—N2	-0.7 (9)
Cl4—Pt1—N2—C10	-88.6 (6)	C6—C7—C11—N2	-178.2 (6)
I3—Pt1—N2—C10	93.4 (7)	C8—C7—C11—C12	177.7 (6)
I4—Pt1—N2—C10	-88.2 (9)	C6—C7—C11—C12	0.3 (9)
N1—Pt1—N2—C11	-1.3 (4)	C1—N1—C12—C4	-1.0 (9)
Cl3—Pt1—N2—C11	-89.3 (7)	Pt1—N1—C12—C4	-178.3 (5)
Cl1—Pt1—N2—C11	-179.5 (4)	C1—N1—C12—C11	178.2 (6)
Cl4—Pt1—N2—C11	89.7 (5)	Pt1—N1—C12—C11	0.9 (7)
I3—Pt1—N2—C11	-88.3 (6)	C5—C4—C12—N1	-178.4 (6)
I4—Pt1—N2—C11	90.1 (8)	C3—C4—C12—N1	2.3 (9)
C12—N1—C1—C2	-1.2 (10)	C5—C4—C12—C11	2.4 (9)
Pt1—N1—C1—C2	175.7 (5)	C3—C4—C12—C11	-176.9 (6)
N1—C1—C2—C3	1.8 (11)	N2—C11—C12—N1	-2.0 (9)
C1—C2—C3—C4	-0.4 (11)	C7—C11—C12—N1	179.5 (6)
C2—C3—C4—C12	-1.6 (10)	N2—C11—C12—C4	177.2 (6)
C2—C3—C4—C5	179.2 (7)	C7—C11—C12—C4	-1.3 (10)
C12—C4—C5—C6	-2.6 (10)		

Symmetry code: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 \cdots C12	0.93	2.73	3.320 (8)	122
C10—H10 \cdots C11	0.93	2.66	3.240 (7)	121