

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N'-(4-Bromophenylsulfonyl)isonicotinohydrazide

Islam Ullah Khan,^a Muhammad Ashfaq,^a
 Muhammad Nadeem Arshad,^{a*} Hamad Ahmad^a and
 Ghulam Mustafa^b

^aMaterials Chemistry Laboratory, Department of Chemistry, GC University, Lahore, Pakistan, and ^bDepartment of Chemistry, University of Gujrat, Gujrat, Pakistan
 Correspondence e-mail: mnachemist@hotmail.com


Received 3 July 2009; accepted 18 July 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å;
 R factor = 0.053; wR factor = 0.146; data-to-parameter ratio = 17.7.

The title compound, $\text{C}_{12}\text{H}_{10}\text{BrN}_3\text{O}_3\text{S}$, crystallizes with two crystallographically independent molecules in the asymmetric unit. The dihedral angles between the two six-membered rings in the molecules are 34.1 (3) and 45.1 (2)°. In the crystal structure, molecules are connected *via* N—H···O and N—H···N hydrogen bonding.

Related literature

For general background to isonicotinic acid hydrazides, see: Carlton (1967). For a related structure, see: Wang *et al.* (2008). For the synthesis and biological activity of isoniazid and hydrazide derivatives, see: Lourenco *et al.* (2008); Kucukguzel *et al.* (2003); Carvalho *et al.* (2008). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{10}\text{BrN}_3\text{O}_3\text{S}$
 $M_r = 356.20$
 Monoclinic, $P2_1/c$
 $a = 10.1229$ (6) Å
 $b = 19.0440$ (12) Å
 $c = 15.0640$ (7) Å
 $\beta = 96.862$ (2)°

$V = 2883.2$ (3) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 3.01$ mm⁻¹
 $T = 296$ K
 $0.36 \times 0.30 \times 0.15$ mm

Data collection

Bruker Kappa APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2007)
 $T_{\min} = 0.349$, $T_{\max} = 0.641$

29398 measured reflections
 6601 independent reflections
 3473 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.146$
 $S = 1.01$
 6601 reflections
 373 parameters

H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 1.06$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.88$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2···O4 ⁱ	0.76 (4)	2.15 (4)	2.882 (4)	165.09
N3—H3···N4 ⁱⁱ	0.78 (4)	2.10 (4)	2.868 (5)	168.21
N6—H6···O6 ⁱⁱⁱ	0.83 (4)	2.26 (4)	2.998 (4)	147.35
N5—H5···N1 ^{iv}	0.83 (4)	2.03 (4)	2.847 (4)	168.32
N3—H3···O1	0.78 (4)	2.49 (4)	2.732 (4)	100 (3)

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x + 1, -y, -z + 1$; (iv) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

The authors acknowledge the Higher Education Commission of Pakistan for providing a grant for the project of strengthening the Materials Chemistry Laboratory at GC University, Lahore, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2151).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2007). SADABS, APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Carlton, W. W. (1967). *Avian Dis.* **11**, 241–254.
- Carvalho, S. A., de Silva, E. F., de Souza, M. V. N., Lourenco, M. C. S. & Vicenate, F. R. (2008). *Bio. Med. Chem. Lett.* **18**, 538–541.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Kucukguzel, S. G., Mazi, A., Sahin, F., Ozturk, S. & Stables, J. (2003). *Eur. J. Med. Chem.* **38**, 1005–1013.
- Lourenco, M. C. S., Ferreira, M. L., de Souza, M. V. N., Peralta, M. A., Vasconcelos, T. R. A. & Henriques, M. G. M. O. (2008). *Eur. J. Med. Chem.* **43**, 1344–1347.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Wang, S.-Y., Song, X.-M. & Duan, L.-X. (2008). *Acta Cryst.* **E64**, o1880.

supporting information

Acta Cryst. (2009). E65, o1991 [doi:10.1107/S1600536809028475]

***N'*-(4-Bromophenylsulfonyl)isonicotinohydrazide**

Islam Ullah Khan, Muhammad Ashfaq, Muhammad Nadeem Arshad, Hamad Ahmad and Ghulam Mustafa

S1. Comment

Isonicotinic acid hydrazide (INH) commonly known as isoniazid is a drug being used for the treatment of tuberculosis (TB) for long time (Carlton, 1967). Different approaches have been made for the synthesis of biologically active derivatives of isoniazid (Lourenco *et al.*, 2008), (Kucukguzel *et al.*, 2007), (Carvalho, *et al.*, 2008) and their crystallographic studies (Wang *et al.*, 2008). In this context we report the crystal structure of title compound (*N'*-[(4-bromophenyl)sulfonyl]-pyridine-4-carbohydrazide) a sulfonamide derivative of Isoniazid.

The title compound crystallizes with two crystallographically independent molecules in the asymmetric unit. The dihedral angle in each of these molecules amount to 34.1 (3) ° in molecule A, while it is 45.1 (2) ° (Fig. 1).

In the crystal structure the molecules are connected via intermolecular and intramolecular N–H···O and N–H···N hydrogen bonding (Fig. 2 and Tab. 1). One of the two independent molecules is connected into dimers via N–H···O hydrogen bonding of the sulfonamide group into $R_2^2(8)$ rings (Bernstein *et al.*, 1995). These dimers are further linked by additional N–H···O and N–H···N hydrogen bonding.

S2. Experimental

To the solution of Isoniazid (0.5 g, 3.646 mmol) in distilled water (10 ml), 4-Bromobenzenesulfonyl chloride (0.9316 g, 3.65 mmol) was suspended. The reaction mixture was stirred at room temperature for 4 hrs at constant pH 8–9, which was adjusted by 1M sodium carbonate solution. After completion of the reaction which was observed by the consumption of suspended 4-Bromobenzenesulfonyl chloride, the pH was adjusted at 2–3 using 1 N HCl solution, which results the formation of a light yellow coloured precipitate, which was filtered off and dried. The product was recrystallized from methanol.

S3. Refinement

The C–H bonds were positioned with idealized geometry with C—H = 0.93 Å and were refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$. The N–H atoms were located in difference map and refined isotropic ($U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{N})$) with varying coordinates.




Figure 1

The structure of (I) with labeling and displacement ellipsoids drawn at the 50% probability level.




Figure 2

Unit cell packing for (I) with hydrogen bonding shown as dashed lines.

***N'*-(4-Bromophenylsulfonyl)isonicotinohydrazide**

Crystal data

$C_{12}H_{10}BrN_3O_3S$

$M_r = 356.20$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 10.1229\ (6)\ \text{\AA}$

$b = 19.0440\ (12)\ \text{\AA}$

$c = 15.0640\ (7)\ \text{\AA}$

$\beta = 96.862\ (2)^\circ$

$V = 2883.2\ (3)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1424$

$D_x = 1.641\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5184 reflections

$\theta = 2.1\text{--}22.9^\circ$

$\mu = 3.01\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Needle, white yellow

$0.36 \times 0.30 \times 0.15\ \text{mm}$

Data collection

Bruker Kappa APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2007)
 $T_{\min} = 0.349$, $T_{\max} = 0.641$

29398 measured reflections
6601 independent reflections
3473 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -13 \rightarrow 11$
 $k = -24 \rightarrow 23$
 $l = -17 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.146$
 $S = 1.01$
6601 reflections
373 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0597P)^2 + 2.5287P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.06 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.88 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.01531 (7)	0.40221 (5)	0.59324 (5)	0.1083 (3)
Br2	1.07622 (6)	0.14607 (4)	0.37352 (6)	0.1046 (3)
S1	0.46424 (10)	0.38694 (6)	0.33172 (7)	0.0392 (3)
S2	0.46755 (9)	0.10499 (5)	0.43498 (6)	0.0328 (2)
O1	0.1616 (3)	0.49151 (15)	0.28196 (19)	0.0467 (7)
O2	0.4756 (3)	0.31692 (15)	0.2999 (2)	0.0510 (8)
O3	0.5802 (3)	0.42527 (17)	0.3664 (2)	0.0566 (9)
O4	0.2996 (2)	0.20910 (13)	0.59686 (17)	0.0376 (7)
O5	0.3909 (3)	0.16199 (14)	0.39574 (18)	0.0420 (7)
O6	0.4282 (3)	0.03415 (14)	0.41370 (17)	0.0413 (7)
N1	-0.2166 (3)	0.3447 (2)	0.1433 (3)	0.0533 (10)
N2	0.2699 (3)	0.41007 (18)	0.2107 (2)	0.0340 (8)
H2	0.270 (4)	0.383 (2)	0.174 (3)	0.041*
N3	0.3969 (3)	0.43457 (18)	0.2456 (2)	0.0353 (8)
H3	0.398 (4)	0.474 (2)	0.259 (3)	0.042*

N4	0.5582 (4)	0.42464 (18)	0.6902 (2)	0.0474 (9)
N5	0.5095 (3)	0.17527 (17)	0.5802 (2)	0.0327 (8)
H5	0.590 (4)	0.176 (2)	0.600 (3)	0.039*
N6	0.4645 (3)	0.11035 (17)	0.5443 (2)	0.0336 (8)
H6	0.493 (4)	0.077 (2)	0.577 (3)	0.040*
C1	0.0288 (4)	0.4054 (2)	0.1991 (3)	0.0352 (9)
C2	0.0182 (4)	0.3454 (3)	0.1500 (4)	0.0685 (16)
H2A	0.0942	0.3235	0.1342	0.082*
C3	-0.1055 (5)	0.3172 (3)	0.1236 (4)	0.0732 (17)
H3A	-0.1102	0.2761	0.0900	0.088*
C4	-0.2061 (5)	0.4033 (3)	0.1895 (4)	0.0774 (18)
H4	-0.2837	0.4244	0.2038	0.093*
C5	-0.0851 (4)	0.4357 (3)	0.2184 (4)	0.0668 (15)
H5A	-0.0828	0.4775	0.2505	0.080*
C6	0.1592 (4)	0.4400 (2)	0.2344 (3)	0.0351 (9)
C7	0.3474 (4)	0.3869 (2)	0.4092 (2)	0.0368 (9)
C8	0.2620 (4)	0.3310 (2)	0.4126 (3)	0.0453 (11)
H8	0.2713	0.2909	0.3786	0.054*
C9	0.1620 (5)	0.3355 (3)	0.4677 (3)	0.0574 (13)
H9	0.1031	0.2984	0.4711	0.069*
C10	0.1508 (5)	0.3954 (3)	0.5172 (3)	0.0559 (13)
C11	0.2374 (5)	0.4504 (3)	0.5151 (3)	0.0557 (13)
H11	0.2291	0.4899	0.5504	0.067*
C12	0.3362 (4)	0.4468 (2)	0.4607 (3)	0.0460 (11)
H12	0.3953	0.4840	0.4581	0.055*
C13	0.4719 (4)	0.29175 (19)	0.6327 (2)	0.0301 (8)
C14	0.5961 (4)	0.3167 (2)	0.6178 (3)	0.0482 (11)
H14	0.6526	0.2894	0.5877	0.058*
C15	0.6342 (5)	0.3825 (2)	0.6483 (3)	0.0556 (12)
H15	0.7183	0.3984	0.6390	0.067*
C16	0.4389 (4)	0.4011 (2)	0.7026 (3)	0.0424 (10)
H16	0.3832	0.4305	0.7306	0.051*
C17	0.3926 (4)	0.3353 (2)	0.6764 (2)	0.0363 (9)
H17	0.3089	0.3205	0.6881	0.044*
C18	0.4181 (4)	0.22155 (19)	0.6023 (2)	0.0288 (8)
C19	0.6340 (4)	0.1167 (2)	0.4150 (2)	0.0346 (9)
C20	0.6736 (4)	0.1792 (3)	0.3808 (3)	0.0514 (11)
H20	0.6125	0.2151	0.3669	0.062*
C21	0.8050 (5)	0.1878 (3)	0.3674 (3)	0.0654 (14)
H21	0.8330	0.2298	0.3443	0.078*
C22	0.8945 (5)	0.1344 (3)	0.3881 (3)	0.0606 (14)
C23	0.8545 (5)	0.0717 (3)	0.4216 (3)	0.0581 (13)
H23	0.9158	0.0357	0.4350	0.070*
C24	0.7237 (4)	0.0623 (2)	0.4352 (3)	0.0477 (11)
H24	0.6957	0.0201	0.4576	0.057*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0866 (5)	0.1511 (8)	0.0966 (5)	0.0053 (5)	0.0490 (4)	-0.0107 (5)
Br2	0.0489 (3)	0.1069 (6)	0.1655 (7)	-0.0228 (3)	0.0441 (4)	-0.0377 (5)
S1	0.0289 (5)	0.0374 (6)	0.0504 (6)	0.0021 (4)	0.0004 (4)	-0.0027 (5)
S2	0.0309 (5)	0.0262 (5)	0.0410 (5)	0.0004 (4)	0.0034 (4)	-0.0016 (4)
O1	0.0373 (16)	0.0419 (19)	0.0602 (18)	0.0023 (13)	0.0028 (14)	-0.0199 (15)
O2	0.0537 (18)	0.0331 (17)	0.0663 (19)	0.0106 (14)	0.0074 (15)	-0.0016 (15)
O3	0.0308 (16)	0.064 (2)	0.072 (2)	-0.0041 (15)	-0.0075 (14)	-0.0053 (17)
O4	0.0277 (14)	0.0306 (16)	0.0552 (17)	-0.0032 (12)	0.0079 (12)	0.0012 (13)
O5	0.0397 (16)	0.0371 (17)	0.0483 (16)	0.0046 (13)	0.0015 (13)	0.0059 (13)
O6	0.0444 (16)	0.0295 (17)	0.0488 (16)	-0.0045 (13)	0.0005 (13)	-0.0060 (13)
N1	0.0282 (19)	0.061 (3)	0.070 (3)	-0.0053 (18)	0.0023 (17)	-0.004 (2)
N2	0.0266 (17)	0.030 (2)	0.045 (2)	-0.0025 (14)	0.0027 (15)	-0.0088 (15)
N3	0.0266 (17)	0.0263 (19)	0.052 (2)	-0.0036 (15)	0.0010 (14)	-0.0082 (17)
N4	0.047 (2)	0.030 (2)	0.062 (2)	-0.0042 (17)	-0.0051 (18)	-0.0024 (18)
N5	0.0257 (16)	0.0244 (19)	0.047 (2)	-0.0017 (14)	0.0003 (14)	-0.0068 (15)
N6	0.0396 (19)	0.0233 (19)	0.0373 (18)	-0.0021 (15)	0.0028 (14)	0.0028 (15)
C1	0.029 (2)	0.038 (2)	0.038 (2)	-0.0023 (17)	0.0013 (16)	-0.0056 (18)
C2	0.019 (2)	0.076 (4)	0.110 (4)	0.002 (2)	0.005 (2)	-0.043 (3)
C3	0.039 (3)	0.070 (4)	0.109 (4)	-0.003 (3)	0.000 (3)	-0.045 (3)
C4	0.030 (3)	0.089 (5)	0.115 (5)	0.005 (3)	0.016 (3)	-0.032 (4)
C5	0.034 (3)	0.067 (4)	0.101 (4)	-0.001 (2)	0.013 (3)	-0.039 (3)
C6	0.033 (2)	0.029 (2)	0.042 (2)	0.0000 (18)	0.0043 (18)	0.0011 (19)
C7	0.036 (2)	0.032 (2)	0.040 (2)	0.0027 (18)	-0.0041 (17)	0.0008 (19)
C8	0.050 (3)	0.035 (3)	0.050 (3)	-0.001 (2)	0.004 (2)	-0.003 (2)
C9	0.054 (3)	0.054 (3)	0.064 (3)	-0.014 (2)	0.007 (2)	0.007 (3)
C10	0.053 (3)	0.068 (4)	0.047 (3)	0.010 (3)	0.009 (2)	0.002 (3)
C11	0.064 (3)	0.056 (3)	0.046 (3)	0.010 (3)	0.002 (2)	-0.009 (2)
C12	0.050 (3)	0.042 (3)	0.044 (2)	0.001 (2)	-0.006 (2)	-0.004 (2)
C13	0.0307 (19)	0.025 (2)	0.034 (2)	-0.0031 (16)	0.0013 (16)	0.0014 (17)
C14	0.036 (2)	0.031 (2)	0.080 (3)	0.0004 (19)	0.017 (2)	-0.007 (2)
C15	0.039 (2)	0.036 (3)	0.091 (4)	-0.010 (2)	0.009 (2)	-0.004 (3)
C16	0.055 (3)	0.032 (3)	0.041 (2)	0.007 (2)	0.005 (2)	-0.0044 (19)
C17	0.038 (2)	0.031 (2)	0.040 (2)	-0.0023 (18)	0.0058 (18)	-0.0013 (18)
C18	0.029 (2)	0.025 (2)	0.031 (2)	-0.0024 (16)	0.0007 (16)	0.0026 (16)
C19	0.034 (2)	0.031 (2)	0.040 (2)	-0.0039 (18)	0.0096 (17)	-0.0066 (18)
C20	0.044 (3)	0.048 (3)	0.063 (3)	-0.003 (2)	0.008 (2)	0.004 (2)
C21	0.059 (3)	0.059 (4)	0.081 (4)	-0.021 (3)	0.019 (3)	-0.004 (3)
C22	0.037 (3)	0.068 (4)	0.079 (3)	-0.008 (3)	0.018 (2)	-0.023 (3)
C23	0.043 (3)	0.052 (3)	0.081 (3)	0.005 (2)	0.011 (2)	-0.017 (3)
C24	0.044 (3)	0.033 (3)	0.068 (3)	-0.001 (2)	0.015 (2)	-0.001 (2)

Geometric parameters (\AA , $^\circ$)

Br1—C10	1.893 (5)	C4—H4	0.9300
Br2—C22	1.891 (4)	C5—H5A	0.9300

S1—O2	1.426 (3)	C7—C8	1.376 (6)
S1—O3	1.427 (3)	C7—C12	1.392 (6)
S1—N3	1.661 (4)	C8—C9	1.386 (6)
S1—C7	1.758 (4)	C8—H8	0.9300
S2—O5	1.421 (3)	C9—C10	1.375 (7)
S2—O6	1.432 (3)	C9—H9	0.9300
S2—N6	1.654 (3)	C10—C11	1.370 (7)
S2—C19	1.760 (4)	C11—C12	1.369 (6)
O1—C6	1.213 (5)	C11—H11	0.9300
O4—C18	1.216 (4)	C12—H12	0.9300
N1—C3	1.306 (6)	C13—C17	1.375 (5)
N1—C4	1.313 (6)	C13—C14	1.387 (5)
N2—C6	1.343 (5)	C13—C18	1.495 (5)
N2—N3	1.409 (4)	C14—C15	1.375 (6)
N2—H2	0.75 (4)	C14—H14	0.9300
N3—H3	0.77 (4)	C15—H15	0.9300
N4—C15	1.322 (6)	C16—C17	1.379 (5)
N4—C16	1.323 (5)	C16—H16	0.9300
N5—C18	1.348 (5)	C17—H17	0.9300
N5—N6	1.403 (4)	C19—C20	1.375 (6)
N5—H5	0.83 (4)	C19—C24	1.386 (6)
N6—H6	0.83 (4)	C20—C21	1.379 (6)
C1—C5	1.351 (6)	C20—H20	0.9300
C1—C2	1.359 (6)	C21—C22	1.372 (7)
C1—C6	1.513 (5)	C21—H21	0.9300
C2—C3	1.377 (6)	C22—C23	1.376 (7)
C2—H2A	0.9300	C23—C24	1.376 (6)
C3—H3A	0.9300	C23—H23	0.9300
C4—C5	1.393 (7)	C24—H24	0.9300
O2—S1—O3	120.34 (19)	C9—C8—H8	120.6
O2—S1—N3	106.87 (17)	C10—C9—C8	119.3 (4)
O3—S1—N3	104.46 (18)	C10—C9—H9	120.3
O2—S1—C7	108.10 (19)	C8—C9—H9	120.3
O3—S1—C7	110.28 (19)	C11—C10—C9	121.9 (4)
N3—S1—C7	105.78 (17)	C11—C10—Br1	118.2 (4)
O5—S2—O6	120.20 (17)	C9—C10—Br1	119.9 (4)
O5—S2—N6	107.03 (17)	C12—C11—C10	119.4 (4)
O6—S2—N6	104.08 (17)	C12—C11—H11	120.3
O5—S2—C19	108.57 (18)	C10—C11—H11	120.3
O6—S2—C19	109.09 (18)	C11—C12—C7	119.3 (4)
N6—S2—C19	107.09 (17)	C11—C12—H12	120.4
C3—N1—C4	116.3 (4)	C7—C12—H12	120.4
C6—N2—N3	120.9 (3)	C17—C13—C14	117.6 (4)
C6—N2—H2	124 (3)	C17—C13—C18	118.3 (3)
N3—N2—H2	115 (3)	C14—C13—C18	124.1 (3)
N2—N3—S1	112.5 (3)	C15—C14—C13	118.7 (4)
N2—N3—H3	114 (3)	C15—C14—H14	120.6

S1—N3—H3	109 (3)	C13—C14—H14	120.6
C15—N4—C16	117.0 (4)	N4—C15—C14	123.9 (4)
C18—N5—N6	118.1 (3)	N4—C15—H15	118.1
C18—N5—H5	125 (3)	C14—C15—H15	118.1
N6—N5—H5	114 (3)	N4—C16—C17	123.5 (4)
N5—N6—S2	113.2 (2)	N4—C16—H16	118.2
N5—N6—H6	111 (3)	C17—C16—H16	118.2
S2—N6—H6	120 (3)	C13—C17—C16	119.2 (4)
C5—C1—C2	117.5 (4)	C13—C17—H17	120.4
C5—C1—C6	118.0 (4)	C16—C17—H17	120.4
C2—C1—C6	124.5 (4)	O4—C18—N5	123.8 (3)
C1—C2—C3	119.7 (4)	O4—C18—C13	121.0 (3)
C1—C2—H2A	120.2	N5—C18—C13	115.2 (3)
C3—C2—H2A	120.2	C20—C19—C24	121.0 (4)
N1—C3—C2	123.8 (5)	C20—C19—S2	120.1 (3)
N1—C3—H3A	118.1	C24—C19—S2	118.9 (3)
C2—C3—H3A	118.1	C19—C20—C21	119.2 (5)
N1—C4—C5	123.6 (4)	C19—C20—H20	120.4
N1—C4—H4	118.2	C21—C20—H20	120.4
C5—C4—H4	118.2	C22—C21—C20	120.1 (5)
C1—C5—C4	119.0 (5)	C22—C21—H21	120.0
C1—C5—H5A	120.5	C20—C21—H21	120.0
C4—C5—H5A	120.5	C21—C22—C23	120.7 (4)
O1—C6—N2	122.8 (4)	C21—C22—Br2	120.8 (4)
O1—C6—C1	121.0 (3)	C23—C22—Br2	118.5 (4)
N2—C6—C1	116.2 (3)	C24—C23—C22	119.8 (5)
C8—C7—C12	121.3 (4)	C24—C23—H23	120.1
C8—C7—S1	120.1 (3)	C22—C23—H23	120.1
C12—C7—S1	118.4 (3)	C23—C24—C19	119.2 (4)
C7—C8—C9	118.8 (4)	C23—C24—H24	120.4
C7—C8—H8	120.6	C19—C24—H24	120.4
C6—N2—N3—S1	97.7 (4)	Br1—C10—C11—C12	-180.0 (3)
O2—S1—N3—N2	60.2 (3)	C10—C11—C12—C7	-0.6 (6)
O3—S1—N3—N2	-171.2 (3)	C8—C7—C12—C11	-0.7 (6)
C7—S1—N3—N2	-54.8 (3)	S1—C7—C12—C11	173.8 (3)
C18—N5—N6—S2	103.9 (3)	C17—C13—C14—C15	1.0 (6)
O5—S2—N6—N5	-55.9 (3)	C18—C13—C14—C15	179.4 (4)
O6—S2—N6—N5	175.8 (2)	C16—N4—C15—C14	-0.1 (7)
C19—S2—N6—N5	60.3 (3)	C13—C14—C15—N4	-1.2 (7)
C5—C1—C2—C3	-1.4 (8)	C15—N4—C16—C17	1.7 (6)
C6—C1—C2—C3	177.8 (5)	C14—C13—C17—C16	0.4 (6)
C4—N1—C3—C2	1.2 (9)	C18—C13—C17—C16	-178.1 (3)
C1—C2—C3—N1	-0.1 (10)	N4—C16—C17—C13	-1.8 (6)
C3—N1—C4—C5	-0.8 (9)	N6—N5—C18—O4	3.7 (5)
C2—C1—C5—C4	1.7 (8)	N6—N5—C18—C13	-175.0 (3)
C6—C1—C5—C4	-177.5 (5)	C17—C13—C18—O4	16.8 (5)
N1—C4—C5—C1	-0.7 (9)	C14—C13—C18—O4	-161.5 (4)

N3—N2—C6—O1	4.6 (6)	C17—C13—C18—N5	-164.5 (3)
N3—N2—C6—C1	-174.9 (3)	C14—C13—C18—N5	17.2 (5)
C5—C1—C6—O1	3.2 (6)	O5—S2—C19—C20	10.3 (4)
C2—C1—C6—O1	-176.0 (5)	O6—S2—C19—C20	143.0 (3)
C5—C1—C6—N2	-177.3 (4)	N6—S2—C19—C20	-105.0 (4)
C2—C1—C6—N2	3.5 (6)	O5—S2—C19—C24	-170.3 (3)
O2—S1—C7—C8	-16.3 (4)	O6—S2—C19—C24	-37.6 (4)
O3—S1—C7—C8	-149.7 (3)	N6—S2—C19—C24	74.4 (4)
N3—S1—C7—C8	97.9 (3)	C24—C19—C20—C21	-0.7 (7)
O2—S1—C7—C12	169.1 (3)	S2—C19—C20—C21	178.7 (4)
O3—S1—C7—C12	35.7 (4)	C19—C20—C21—C22	0.0 (7)
N3—S1—C7—C12	-76.7 (3)	C20—C21—C22—C23	0.6 (8)
C12—C7—C8—C9	1.1 (6)	C20—C21—C22—Br2	-178.2 (4)
S1—C7—C8—C9	-173.4 (3)	C21—C22—C23—C24	-0.5 (8)
C7—C8—C9—C10	-0.1 (7)	Br2—C22—C23—C24	178.3 (4)
C8—C9—C10—C11	-1.2 (7)	C22—C23—C24—C19	-0.2 (7)
C8—C9—C10—Br1	-179.7 (3)	C20—C19—C24—C23	0.8 (6)
C9—C10—C11—C12	1.6 (7)	S2—C19—C24—C23	-178.6 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2 \cdots O4 ⁱ	0.76 (4)	2.15 (4)	2.882 (4)	165.09
N3—H3 \cdots N4 ⁱⁱ	0.78 (4)	2.10 (4)	2.868 (5)	168.21
N6—H6 \cdots O6 ⁱⁱⁱ	0.83 (4)	2.26 (4)	2.998 (4)	147.35
N5—H5 \cdots N1 ^{iv}	0.83 (4)	2.03 (4)	2.847 (4)	168.32
N3—H3 \cdots O1	0.78 (4)	2.49 (4)	2.732 (4)	100 (3)

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x+1, -y, -z+1$; (iv) $x+1, -y+1/2, z+1/2$.