

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[*(4-aminobenzoato) aqua-silver(I)*]- μ -hexamethylenetetramine]

 Jia-Jun Han,^{a*} Tian-Yun Zhang^a and Lin Geng^b
^aSchool of the Ocean, Harbin Institute of Technology, Weihai 264209, People's Republic of China, and ^bSchool of Materials Science and Engineering, Harbin Institute of Technology, Harbin 150001, People's Republic of China


Correspondence e-mail: hanjiajunhit@yahoo.com.cn

Received 13 December 2009; accepted 22 December 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.019; wR factor = 0.048; data-to-parameter ratio = 12.7.

In the title coordination polymer, $[\text{Ag}(\text{C}_7\text{H}_6\text{NO}_2)(\text{C}_6\text{H}_{12}\text{N}_4)(\text{H}_2\text{O})]_n$, the Ag^{I} ion is five-coordinated by two carboxylate O atoms from one 4-aminobenzoate anion (L), two N atoms from two different hexamethylenetetramine (hmt) ligands, and one water O atom in a distorted square-pyramidal geometry. The metal atom lies on a mirror plane and the L anion, hmt ligand and water molecule all lie across crystallographic mirror planes. Each hmt ligand bridges two neighboring Ag^{I} ions, resulting in the formation of a chain structure along the b axis. The chains are linked into a three-dimensional framework by $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

 For the applications and structures of silver(I) coordination polymers, see: Yang *et al.* (2007, 2008).


Experimental

Crystal data

 $[\text{Ag}(\text{C}_7\text{H}_6\text{NO}_2)(\text{C}_6\text{H}_{12}\text{N}_4)(\text{H}_2\text{O})]$
 $M_r = 402.21$

 Orthorhombic, $Pnma$
 $a = 19.8107$ (11) Å

 $b = 6.4877$ (3) Å

 $c = 11.3257$ (6) Å

 $V = 1455.65$ (13) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 1.41$ mm⁻¹
 $T = 293$ K

 $0.31 \times 0.27 \times 0.22$ mm

Data collection

Bruker APEX CCD area-detector diffractometer

 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

 $T_{\text{min}} = 0.66$, $T_{\text{max}} = 0.87$

7757 measured reflections

1557 independent reflections

 1373 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.019$
 $wR(F^2) = 0.048$
 $S = 1.08$

1557 reflections

123 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.37$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ag1—N1	2.3862 (17)	Ag1—O1	2.5413 (14)
Ag1—O1W	2.445 (2)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1W1 \cdots O1 ⁱ	0.81 (2)	1.95 (2)	2.742 (2)	168 (2)
N2—H2A \cdots O1 ⁱⁱ	0.84 (2)	2.27 (2)	3.072 (2)	159 (2)

 Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + \frac{1}{2}, -y + 1, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Harbin Institute of Technology for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2988).

References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yang, J., Ma, J.-F., Batten, S. R. & Su, Z.-M. (2008). *Chem. Commun.* pp. 2233–2235.
- Yang, G., Wang, Y.-L., Li, J.-P., Zhu, Y., Wang, S.-M., Hou, H.-W., Fan, Y.-T. & Ng, S. W. (2007). *Eur. J. Inorg. Chem.* pp. 714–719.

supporting information

Acta Cryst. (2010). E66, m111 [https://doi.org/10.1107/S1600536809055044]

catena-Poly[[*(*(4-aminobenzoato)aquasilver(I)*)]-μ*-hexamethylenetetramine]

Jia-Jun Han, Tian-Yun Zhang and Lin Geng

S1. Experimental

An aqueous solution (10 ml) of HL (0.104 g, 0.5 mmol) was added to solid Ag₂CO₃ (0.25 mmol) and stirred for several minutes until no further CO₂ was given off. A solution of hmt (0.5 mmol) in acetonitrile (10 ml) was then added and a white precipitate formed. The precipitate was dissolved by dropwise addition of an aqueous solution of NH₃ (14 M). Colourless blocks of the title compound were obtained by evaporation of the solution for several days at room temperature (33% yield).

S2. Refinement

Amino and water H-atoms were located in a difference Fourier map, and refined freely. The remaining H atoms were positioned geometrically (C-H = 0.93 Å) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

A view of the local coordination of the Ag^I centre in the title compound, showing the atom-labelling scheme.

Displacement ellipsoids are drawn at the 30% probability level. Symmetry codes: (i) $x, 3/2 - y, z$; (ii) $x, 1/2 - y, z$.




Figure 2
Part of the polymeric chain in the title compound.

catena-Poly[[4-aminobenzoato]aquasilver(I)]- μ -hexamethylenetetramine]

Crystal data

[Ag(C₇H₆NO₂)(C₆H₁₂N₄)(H₂O)]

$M_r = 402.21$

Orthorhombic, *Pnma*

Hall symbol: -P 2ac 2n

$a = 19.8107$ (11) Å

$b = 6.4877$ (3) Å

$c = 11.3257$ (6) Å

$V = 1455.65$ (13) Å³

$Z = 4$

$F(000) = 816$

$D_x = 1.835$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1557 reflections

$\theta = 3.0$ – 26.0°

$\mu = 1.41$ mm⁻¹

$T = 293$ K

Block, colourless

$0.31 \times 0.27 \times 0.22$ mm

Data collection

Bruker APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.66$, $T_{\max} = 0.87$

7757 measured reflections

1557 independent reflections

1373 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.1^\circ$

$h = -24 \rightarrow 23$

$k = -7 \rightarrow 7$

$l = -13 \rightarrow 9$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.019$

$wR(F^2) = 0.048$

$S = 1.08$

1557 reflections

123 parameters

0 restraints

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0211P)^2 + 0.5726P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ag1	0.425226 (10)	0.2500	0.638361 (18)	0.02218 (9)	
O1W	0.54862 (12)	0.2500	0.6362 (2)	0.0324 (5)	
H1W1	0.5643 (11)	0.352 (4)	0.606 (2)	0.036 (7)*	
O1	0.38093 (7)	0.4203 (2)	0.45173 (12)	0.0306 (4)	
N1	0.42059 (7)	0.5604 (3)	0.75114 (14)	0.0198 (4)	
N2	0.23019 (13)	0.2500	-0.0371 (2)	0.0261 (6)	
H2A	0.2075 (11)	0.358 (3)	-0.0497 (19)	0.033 (7)*	
N3	0.47738 (11)	0.7500	0.9101 (2)	0.0186 (5)	
N4	0.35383 (11)	0.7500	0.8979 (2)	0.0228 (5)	
C1	0.26785 (13)	0.2500	0.0657 (2)	0.0182 (6)	
C2	0.28812 (9)	0.4347 (3)	0.11800 (16)	0.0206 (4)	
H2	0.2781	0.5594	0.0814	0.025*	
C3	0.32294 (9)	0.4339 (3)	0.22365 (16)	0.0196 (4)	
H3	0.3357	0.5585	0.2576	0.024*	
C4	0.33924 (13)	0.2500	0.2803 (2)	0.0178 (6)	
C5	0.36952 (13)	0.2500	0.4018 (2)	0.0222 (6)	
C6	0.35759 (9)	0.5670 (3)	0.82288 (17)	0.0235 (5)	
H6A	0.3189	0.5649	0.7704	0.028*	
H6B	0.3554	0.4448	0.8721	0.028*	
C7	0.47854 (9)	0.5664 (3)	0.83436 (16)	0.0214 (4)	
H7A	0.4776	0.4440	0.8836	0.026*	
H7B	0.5203	0.5647	0.7897	0.026*	
C8	0.41317 (12)	0.7500	0.9761 (2)	0.0223 (6)	
H8A	0.4115	0.6292	1.0264	0.027*	0.50
H8B	0.4115	0.8708	1.0264	0.027*	0.50
C9	0.42303 (13)	0.7500	0.6783 (2)	0.0200 (6)	
H9A	0.4642	0.7500	0.6321	0.024*	
H9B	0.3852	0.7500	0.6240	0.024*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.02328 (13)	0.02679 (14)	0.01649 (13)	0.000	-0.00302 (9)	0.000
O1W	0.0240 (12)	0.0378 (15)	0.0352 (14)	0.000	0.0073 (10)	0.000
O1	0.0353 (8)	0.0378 (9)	0.0187 (7)	-0.0053 (8)	-0.0049 (6)	-0.0054 (7)
N1	0.0170 (8)	0.0292 (10)	0.0130 (8)	-0.0014 (7)	-0.0024 (6)	0.0004 (7)
N2	0.0244 (14)	0.0349 (17)	0.0189 (13)	0.000	-0.0072 (11)	0.000
N3	0.0137 (11)	0.0288 (14)	0.0134 (11)	0.000	0.0003 (9)	0.000
N4	0.0140 (11)	0.0378 (15)	0.0165 (12)	0.000	-0.0014 (9)	0.000
C1	0.0126 (12)	0.0283 (15)	0.0137 (14)	0.000	0.0039 (11)	0.000
C2	0.0208 (10)	0.0229 (11)	0.0179 (10)	0.0016 (8)	0.0010 (8)	0.0035 (8)
C3	0.0200 (10)	0.0210 (11)	0.0179 (10)	-0.0011 (8)	0.0041 (8)	-0.0020 (8)
C4	0.0112 (12)	0.0279 (16)	0.0143 (13)	0.000	0.0024 (10)	0.000
C5	0.0145 (13)	0.0355 (18)	0.0168 (14)	0.000	0.0022 (11)	0.000
C6	0.0159 (9)	0.0355 (13)	0.0192 (10)	-0.0040 (9)	-0.0024 (8)	0.0019 (9)
C7	0.0167 (10)	0.0308 (12)	0.0167 (10)	0.0015 (9)	-0.0025 (8)	0.0019 (8)
C8	0.0152 (14)	0.0377 (18)	0.0139 (14)	0.000	-0.0018 (11)	0.000
C9	0.0168 (13)	0.0301 (17)	0.0132 (13)	0.000	-0.0041 (11)	0.000

Geometric parameters (\AA , $^\circ$)

Ag1—N1	2.3862 (17)	C1—C2	1.396 (2)
Ag1—N1 ⁱ	2.3862 (17)	C1—C2 ⁱ	1.396 (2)
Ag1—O1W	2.445 (2)	C2—C3	1.381 (3)
Ag1—O1	2.5413 (14)	C2—H2	0.93
Ag1—O1 ⁱ	2.5413 (14)	C3—C4	1.393 (2)
O1W—H1W1	0.81 (2)	C3—H3	0.93
O1—C5	1.2615 (19)	C4—C3 ⁱ	1.393 (2)
N1—C9	1.482 (2)	C4—C5	1.500 (4)
N1—C7	1.486 (2)	C5—O1 ⁱ	1.2615 (19)
N1—C6	1.490 (2)	C6—H6A	0.97
N2—C1	1.383 (4)	C6—H6B	0.97
N2—H2A	0.84 (2)	C7—H7A	0.97
N3—C7	1.468 (2)	C7—H7B	0.97
N3—C7 ⁱⁱ	1.468 (2)	C8—H8A	0.97
N3—C8	1.475 (3)	C8—H8B	0.97
N4—C6	1.462 (2)	C9—N1 ⁱⁱ	1.482 (2)
N4—C6 ⁱⁱ	1.462 (2)	C9—H9A	0.97
N4—C8	1.472 (3)	C9—H9B	0.97
N1—Ag1—N1 ⁱ	115.09 (8)	C2—C3—H3	119.4
N1—Ag1—O1W	92.52 (5)	C4—C3—H3	119.4
N1 ⁱ —Ag1—O1W	92.52 (5)	C3 ⁱ —C4—C3	117.9 (2)
N1—Ag1—O1	93.74 (5)	C3 ⁱ —C4—C5	121.01 (12)
N1 ⁱ —Ag1—O1	143.00 (5)	C3—C4—C5	121.01 (12)
O1W—Ag1—O1	109.69 (6)	O1 ⁱ —C5—O1	122.2 (3)
N1—Ag1—O1 ⁱ	143.00 (5)	O1 ⁱ —C5—C4	118.87 (13)

N1 ⁱ —Ag1—O1 ⁱ	93.74 (5)	O1—C5—C4	118.87 (13)
O1W—Ag1—O1 ⁱ	109.69 (6)	N4—C6—N1	112.52 (17)
O1—Ag1—O1 ⁱ	51.53 (7)	N4—C6—H6A	109.1
Ag1—O1W—H1W1	112.9 (17)	N1—C6—H6A	109.1
C5—O1—Ag1	93.12 (14)	N4—C6—H6B	109.1
C9—N1—C7	107.80 (16)	N1—C6—H6B	109.1
C9—N1—C6	107.87 (17)	H6A—C6—H6B	107.8
C7—N1—C6	107.49 (15)	N3—C7—N1	112.34 (17)
C9—N1—Ag1	113.67 (12)	N3—C7—H7A	109.1
C7—N1—Ag1	109.39 (12)	N1—C7—H7A	109.1
C6—N1—Ag1	110.39 (12)	N3—C7—H7B	109.1
C1—N2—H2A	115.4 (16)	N1—C7—H7B	109.1
C7—N3—C7 ⁱⁱ	108.4 (2)	H7A—C7—H7B	107.9
C7—N3—C8	108.02 (14)	N4—C8—N3	112.6 (2)
C7 ⁱⁱ —N3—C8	108.02 (14)	N4—C8—H8A	109.1
C6—N4—C6 ⁱⁱ	108.6 (2)	N3—C8—H8A	109.1
C6—N4—C8	107.99 (14)	N4—C8—H8B	109.1
C6 ⁱⁱ —N4—C8	107.99 (14)	N3—C8—H8B	109.1
N2—C1—C2	120.83 (12)	H8A—C8—H8B	107.8
N2—C1—C2 ⁱ	120.83 (12)	N1 ⁱⁱ —C9—N1	112.3 (2)
C2—C1—C2 ⁱ	118.3 (2)	N1 ⁱⁱ —C9—H9A	109.2
C3—C2—C1	120.53 (19)	N1—C9—H9A	109.2
C3—C2—H2	119.7	N1 ⁱⁱ —C9—H9B	109.2
C1—C2—H2	119.7	N1—C9—H9B	109.2
C2—C3—C4	121.23 (19)	H9A—C9—H9B	107.9
N1—Ag1—O1—C5	-166.41 (14)	Ag1—O1—C5—C4	178.7 (2)
N1 ⁱ —Ag1—O1—C5	-23.97 (18)	C3 ⁱ —C4—C5—O1 ⁱ	0.9 (4)
O1W—Ag1—O1—C5	99.57 (14)	C3—C4—C5—O1 ⁱ	177.1 (2)
O1 ⁱ —Ag1—O1—C5	-0.38 (15)	C3 ⁱ —C4—C5—O1	-177.1 (2)
N1 ⁱ —Ag1—N1—C9	-179.54 (11)	C3—C4—C5—O1	-0.9 (4)
O1W—Ag1—N1—C9	86.50 (15)	C6 ⁱⁱ —N4—C6—N1	58.4 (3)
O1—Ag1—N1—C9	-23.43 (14)	C8—N4—C6—N1	-58.5 (2)
O1 ⁱ —Ag1—N1—C9	-41.73 (17)	C9—N1—C6—N4	-57.9 (2)
N1 ⁱ —Ag1—N1—C7	59.91 (14)	C7—N1—C6—N4	58.1 (2)
O1W—Ag1—N1—C7	-34.05 (13)	Ag1—N1—C6—N4	177.33 (13)
O1—Ag1—N1—C7	-143.98 (12)	C7 ⁱⁱ —N3—C7—N1	-58.6 (2)
O1 ⁱ —Ag1—N1—C7	-162.28 (10)	C8—N3—C7—N1	58.2 (2)
N1 ⁱ —Ag1—N1—C6	-58.17 (14)	C9—N1—C7—N3	58.2 (2)
O1W—Ag1—N1—C6	-152.14 (12)	C6—N1—C7—N3	-57.8 (2)
O1—Ag1—N1—C6	97.94 (12)	Ag1—N1—C7—N3	-177.73 (13)
O1 ⁱ —Ag1—N1—C6	79.64 (15)	C6—N4—C8—N3	58.64 (14)
N2—C1—C2—C3	176.9 (2)	C6 ⁱⁱ —N4—C8—N3	-58.64 (14)
C2 ⁱ —C1—C2—C3	-4.8 (4)	C7—N3—C8—N4	-58.54 (13)
C1—C2—C3—C4	0.5 (3)	C7 ⁱⁱ —N3—C8—N4	58.54 (13)
C2—C3—C4—C3 ⁱ	3.8 (4)	C7—N1—C9—N1 ⁱⁱ	-58.1 (2)

C2—C3—C4—C5	-172.5 (2)	C6—N1—C9—N1 ⁱⁱ	57.7 (2)
Ag1—O1—C5—O1 ⁱ	0.7 (3)	Ag1—N1—C9—N1 ⁱⁱ	-179.56 (11)

Symmetry codes: (i) $x, -y+1/2, z$; (ii) $x, -y+3/2, z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1 <i>W</i> —H1 <i>W</i> 1...O1 ⁱⁱⁱ	0.81 (2)	1.95 (2)	2.742 (2)	168 (2)
N2—H2 <i>A</i> ...O1 ^{iv}	0.84 (2)	2.27 (2)	3.072 (2)	159 (2)

Symmetry codes: (iii) $-x+1, -y+1, -z+1$; (iv) $-x+1/2, -y+1, z-1/2$.