

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(*N*-butyl-*N*-ethylthiocarbamate- κ^2S,S')nickel(II)

 Wan Nur Shazwani Wan Juhari,^a Ibrahim Baba,^a Yang Farina^a and Seik Weng Ng^{b*}
^aSchool of Chemical Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 21 February 2010; accepted 22 February 2010

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.034; wR factor = 0.098; data-to-parameter ratio = 23.2.

The dithiocarbamate anions in the title compound, $[\text{Ni}(\text{C}_7\text{H}_{14}\text{NS}_2)_2]$, chelate to the Ni^{II} atom, which is four-coordinate in a square-planar geometry. The Ni^{II} atom lies on a center of inversion.

Related literature

For nickel bis(diethylthiocarbamate) and nickel bis(di-n-butylthiocarbamate), see: Bonamico *et al.* (1965); Khan *et al.* (1987); Lokaj *et al.* (1984).



Experimental

Crystal data

 $[\text{Ni}(\text{C}_7\text{H}_{14}\text{NS}_2)_2]$
 $M_r = 411.33$

 Monoclinic, $P2_1/n$
 $a = 8.5641$ (9) Å
 $b = 8.6316$ (9) Å
 $c = 13.6047$ (14) Å
 $\beta = 94.753$ (2)°
 $V = 1002.23$ (18) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.38$ mm⁻¹
 $T = 293$ K
 $0.25 \times 0.25 \times 0.05$ mm

Data collection

 Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.724$, $T_{\text{max}} = 0.934$

 9338 measured reflections
 2295 independent reflections
 1628 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.098$
 $S = 1.03$
 2295 reflections
 99 parameters

 6 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XSEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank Universiti Kebangsaan Malaysia (UKM-GUP-NBT-08-27-111 and 06-01-02-SF0539) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2734).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bonamico, M., Dessy, G., Mariani, C., Vaciego, A. & Zambonelli, L. (1965). *Acta Cryst.* **19**, 619–626.
 Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Khan, M. N. I., Fackler, J. P., Murray, H. H., Heinrich, D. D. & Campana, C. (1987). *Acta Cryst.* **C43**, 1917–1919.
 Lokaj, J., Vrabec, V. & Kello, E. (1984). *Chem. Zvesti.* **38**, 313–317.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). publCIF. In preparation.

supporting information

Acta Cryst. (2010). E66, m339 [doi:10.1107/S1600536810006677]

Bis(*N*-butyl-*N*-ethyldithiocarbamato- κ^2S,S')nickel(II)

Wan Nur Shazwani Wan Juhari, Ibrahim Baba, Yang Farina and Seik Weng Ng

S1. Experimental

Nickel(II) chloride (10 mmol), butylethylamine (10 mmol) and carbon disulfide (10 mmol) were reacted in ethanol (50 ml) at 277 K to produce a brown solid. The mixture was stirred for an hour. The solid was collected and recrystallized from ethanol.

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.96 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2 to 1.5 $U(\text{C})$.




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $\text{Ni}(\text{C}_7\text{H}_{14}\text{NS}_2)_2$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Bis(*N*-butyl-*N*-ethyldithiocarbamato- κ^2S,S')nickel(II)*Crystal data*

$[\text{Ni}(\text{C}_7\text{H}_{14}\text{NS}_2)_2]$
 $M_r = 411.33$

Monoclinic, $P2_1/n$
Hall symbol: -P 2yn

$a = 8.5641$ (9) Å
 $b = 8.6316$ (9) Å
 $c = 13.6047$ (14) Å
 $\beta = 94.753$ (2)°
 $V = 1002.23$ (18) Å³
 $Z = 2$
 $F(000) = 436$
 $D_x = 1.363$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 2307 reflections
 $\theta = 2.4$ – 24.6 °
 $\mu = 1.38$ mm⁻¹
 $T = 293$ K
 Plate, brown
 $0.25 \times 0.25 \times 0.05$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.724$, $T_{\max} = 0.934$

9338 measured reflections
 2295 independent reflections
 1628 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$
 $\theta_{\max} = 27.5$ °, $\theta_{\min} = 2.7$ °
 $h = -10 \rightarrow 11$
 $k = -11 \rightarrow 11$
 $l = -17 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.098$
 $S = 1.03$
 2295 reflections
 99 parameters
 6 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0469P)^2 + 0.1755P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.21$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.5000	0.5000	0.05331 (16)
S1	0.75597 (8)	0.48176 (8)	0.53089 (5)	0.0629 (2)
S2	0.52687 (8)	0.56847 (9)	0.65651 (5)	0.0631 (2)
N1	0.8339 (3)	0.5601 (3)	0.72015 (17)	0.0647 (6)
C1	0.7236 (3)	0.5403 (3)	0.64766 (19)	0.0564 (6)
C2	0.7964 (3)	0.6124 (3)	0.81795 (18)	0.0658 (7)
H2A	0.7008	0.6732	0.8111	0.079*
H2B	0.8799	0.6791	0.8456	0.079*
C3	0.7752 (4)	0.4796 (3)	0.8887 (2)	0.0746 (8)
H3A	0.6875	0.4163	0.8630	0.090*
H3B	0.8684	0.4154	0.8928	0.090*
C4	0.7460 (4)	0.5353 (4)	0.9904 (2)	0.0836 (9)
H4A	0.6554	0.6032	0.9856	0.100*
H4B	0.8355	0.5955	1.0167	0.100*
C5	0.7180 (4)	0.4044 (4)	1.0615 (2)	0.1014 (11)
H5A	0.6962	0.4470	1.1241	0.152*
H5B	0.8098	0.3403	1.0699	0.152*
H5C	0.6305	0.3433	1.0356	0.152*

C6	1.0008 (4)	0.5276 (4)	0.7074 (2)	0.0805 (9)
H6A	1.0079	0.4559	0.6530	0.097*
H6B	1.0490	0.4791	0.7667	0.097*
C7	1.0871 (4)	0.6734 (4)	0.6871 (2)	0.0951 (10)
H7A	1.1957	0.6500	0.6818	0.143*
H7B	1.0780	0.7453	0.7401	0.143*
H7C	1.0432	0.7183	0.6265	0.143*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0586 (3)	0.0553 (3)	0.0464 (3)	−0.0031 (2)	0.0063 (2)	−0.00089 (19)
S1	0.0628 (4)	0.0763 (4)	0.0504 (4)	0.0013 (3)	0.0095 (3)	−0.0089 (3)
S2	0.0589 (4)	0.0785 (5)	0.0526 (4)	0.0006 (3)	0.0090 (3)	−0.0081 (3)
N1	0.0596 (13)	0.0780 (14)	0.0572 (13)	0.0057 (11)	0.0077 (11)	−0.0171 (12)
C1	0.0629 (16)	0.0556 (14)	0.0513 (15)	−0.0005 (11)	0.0083 (12)	−0.0052 (11)
C2	0.0670 (17)	0.0758 (17)	0.0542 (16)	0.0016 (13)	0.0023 (13)	−0.0198 (13)
C3	0.080 (2)	0.080 (2)	0.0629 (18)	0.0053 (14)	0.0010 (15)	−0.0130 (15)
C4	0.093 (2)	0.096 (2)	0.0621 (19)	0.0055 (17)	0.0047 (17)	−0.0136 (17)
C5	0.118 (3)	0.111 (3)	0.074 (2)	0.001 (2)	0.002 (2)	0.000 (2)
C6	0.0653 (18)	0.109 (3)	0.0660 (19)	0.0082 (17)	−0.0034 (15)	−0.0236 (17)
C7	0.076 (2)	0.125 (3)	0.086 (2)	0.008 (2)	0.0171 (17)	−0.006 (2)

Geometric parameters (Å, °)

Ni1—S1 ⁱ	2.2032 (8)	C3—H3B	0.9700
Ni1—S1	2.2032 (8)	C4—C5	1.519 (4)
Ni1—S2	2.2034 (7)	C4—H4A	0.9700
Ni1—S2 ⁱ	2.2034 (7)	C4—H4B	0.9700
S1—C1	1.712 (3)	C5—H5A	0.9600
S2—C1	1.716 (3)	C5—H5B	0.9600
N1—C1	1.319 (3)	C5—H5C	0.9600
N1—C2	1.466 (3)	C6—C7	1.497 (4)
N1—C6	1.481 (4)	C6—H6A	0.9700
C2—C3	1.517 (4)	C6—H6B	0.9700
C2—H2A	0.9700	C7—H7A	0.9600
C2—H2B	0.9700	C7—H7B	0.9600
C3—C4	1.505 (4)	C7—H7C	0.9600
C3—H3A	0.9700		
S1 ⁱ —Ni1—S1	180.0	H3A—C3—H3B	107.9
S1 ⁱ —Ni1—S2	100.82 (2)	C3—C4—C5	113.3 (3)
S1—Ni1—S2	79.18 (2)	C3—C4—H4A	108.9
S1 ⁱ —Ni1—S2 ⁱ	79.18 (2)	C5—C4—H4A	108.9
S1—Ni1—S2 ⁱ	100.82 (2)	C3—C4—H4B	108.9
S2—Ni1—S2 ⁱ	180.0	C5—C4—H4B	108.9
C1—S1—Ni1	85.45 (10)	H4A—C4—H4B	107.7
C1—S2—Ni1	85.34 (9)	C4—C5—H5A	109.5

C1—N1—C2	121.4 (2)	C4—C5—H5B	109.5
C1—N1—C6	121.7 (2)	H5A—C5—H5B	109.5
C2—N1—C6	116.8 (2)	C4—C5—H5C	109.5
N1—C1—S1	124.8 (2)	H5A—C5—H5C	109.5
N1—C1—S2	125.1 (2)	H5B—C5—H5C	109.5
S1—C1—S2	110.04 (16)	N1—C6—C7	111.0 (3)
N1—C2—C3	113.0 (2)	N1—C6—H6A	109.4
N1—C2—H2A	109.0	C7—C6—H6A	109.4
C3—C2—H2A	109.0	N1—C6—H6B	109.4
N1—C2—H2B	109.0	C7—C6—H6B	109.4
C3—C2—H2B	109.0	H6A—C6—H6B	108.0
H2A—C2—H2B	107.8	C6—C7—H7A	109.5
C4—C3—C2	112.3 (2)	C6—C7—H7B	109.5
C4—C3—H3A	109.1	H7A—C7—H7B	109.5
C2—C3—H3A	109.1	C6—C7—H7C	109.5
C4—C3—H3B	109.1	H7A—C7—H7C	109.5
C2—C3—H3B	109.1	H7B—C7—H7C	109.5
S2—Ni1—S1—C1	-0.24 (9)	Ni1—S1—C1—S2	0.32 (12)
S2 ⁱ —Ni1—S1—C1	179.76 (9)	Ni1—S2—C1—N1	180.0 (2)
S1 ⁱ —Ni1—S2—C1	-179.76 (9)	Ni1—S2—C1—S1	-0.32 (12)
S1—Ni1—S2—C1	0.24 (9)	C1—N1—C2—C3	94.2 (3)
C2—N1—C1—S1	179.3 (2)	C6—N1—C2—C3	-84.6 (3)
C6—N1—C1—S1	-2.0 (4)	N1—C2—C3—C4	176.7 (3)
C2—N1—C1—S2	-1.0 (4)	C2—C3—C4—C5	177.8 (3)
C6—N1—C1—S2	177.6 (2)	C1—N1—C6—C7	98.5 (3)
Ni1—S1—C1—N1	-180.0 (2)	C2—N1—C6—C7	-82.8 (3)

Symmetry code: (i) $-x+1, -y+1, -z+1$.