

Tris[2-(1*H*-imidazol-2-yl)imidazol-1-ido]-cobalt(III)

Qi Ma,^{a,b} Miaoli Zhu,^a Sisi Feng^a and Liping Lu^{a*}

^aInstitute of Molecular Science, Key Laboratory of Chemical Biology and Molecular Engineering of the Education Ministry, Shanxi University, Taiyuan, Shanxi 030006, People's Republic of China, and ^bCollege of Chemistry and Chemical Engineering, Shanxi Datong University, Datong, Shanxi 037009, People's Republic of China
Correspondence e-mail: luliping@sxu.edu.cn


Received 25 March 2010; accepted 29 March 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.041; wR factor = 0.138; data-to-parameter ratio = 13.3.

In the title compound, $[\text{Co}(\text{C}_6\text{H}_5\text{N}_4)_3]$, the Co^{III} atom adopts a distorted octahedral CoN_6 coordination geometry, arising from three *N,N'*-bidentate deprotonated 2,2'-biimidazole ligands. The dihedral angles between the five-membered rings of the ligands are 4.1 (2), 9.4 (2) and 10.5 (2) $^\circ$. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds, generating a layered network lying in $(11\bar{1})$.

Related literature

For related structures, see: Tadokoro & Nakasui (2000); Ye *et al.* (2005); Zhang *et al.* (2008).



Experimental

Crystal data

$[\text{Co}(\text{C}_6\text{H}_5\text{N}_4)_3]$

$M_r = 458.35$

Monoclinic, $P2_1/n$

$a = 12.299 (3)\text{ \AA}$

$b = 12.524 (3)\text{ \AA}$

$c = 12.932 (3)\text{ \AA}$

$\beta = 97.773 (4)^\circ$

$V = 1973.6 (8)\text{ \AA}^3$

$Z = 4$

$\text{Mo } K\alpha$ radiation

$\mu = 0.90\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.5 \times 0.4 \times 0.3\text{ mm}$

Data collection

Bruker SMART CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2000)

$T_{\min} = 0.654$, $T_{\max} = 0.762$

10212 measured reflections

3728 independent reflections

2358 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.138$

$S = 0.98$

3728 reflections

280 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.48\text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.49\text{ e } \text{\AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Co1—N4	1.917 (3)	Co1—N1	1.929 (3)
Co1—N3	1.922 (3)	Co1—N5	1.941 (3)
Co1—N6	1.926 (3)	Co1—N2	1.944 (3)
N4—Co1—N3	82.54 (12)	N1—Co1—N2	82.18 (12)
N6—Co1—N5	81.67 (13)		

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N11—H6 \cdots N12 ⁱ	0.86	1.95	2.808 (4)	172
N7—H11 \cdots N8 ⁱⁱ	0.86	1.99	2.814 (4)	159
N9—H19 \cdots N10 ⁱⁱⁱ	0.86	1.95	2.796 (4)	169
Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $-x, -y + 1, -z + 2$; (iii) $-x + 1, -y + 2, -z + 2$.				

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors acknowledge financial support from the National Natural Science Foundation of China (grant No. 20471033), the Province Natural Science Foundation of Shanxi Province of China (grant No. 20051013) and the Overseas Returned Scholar Foundation of Shanxi Province of China in 2008, as well as Doctor Startup Foundation of Shanxi University of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5377).

References

- Bruker (2000). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tadokoro, M. & Nakasui, K. (2000). *Coord. Chem. Rev.* **198**, 205–218.
- Ye, B. H., Ding, B. B., Weng, Y. Q. & Chen, X. M. (2005). *Cryst. Growth Des.* **5**, 801–806.
- Zhang, L.-C., Zhu, Z.-M., You, W.-S., Chang, S. & Wang, E.-B. (2008). *Acta Cryst. E* **64**, m308.

supporting information

Acta Cryst. (2010). E66, m500 [https://doi.org/10.1107/S1600536810011785]

Tris[2-(1*H*-imidazol-2-yl)imidazol-1-ido]cobalt(III)

Qi Ma, Miaoli Zhu, Sisi Feng and Liping Lu

S1. Comment

The neutral molecule 2,2'-biimidazole (H_2biim) and its monoanionic derivative(Hbiim^-) is a particular organic target for construction of hybrid materials. Its molecular moieties possess a double property. Namely they can be coordinated to metal centres and can act as a donor in hydrogen bonding interactions (Tadokoro & Nakasuji, 2000). The crystal structure of (I) is reported here.


The X-ray crystallographic analysis shows that the molecule of the compound (I) consists of three Hbiim^- and one Co^{3+} (Fig. 1). The Co^{3+} ion adopted octahedron coordination geometry, and coordinated with three Hbiim^- anion. Average bond distance $\text{Co}-\text{N}$ is 1.93 (3) Å, shorter than $\text{Co}-\text{N}$ bond distance found in related structures, i.e. 2.116 (2)-2.118 Å in $[\text{Co}(\text{H}_2\text{biim})_2(1,2-\text{bdc})]$ (Ye *et al.*, 2005), 2.1563 (18) Å, in diaquabis(2,2'-biimidazole)cobalt(II) dichloride (Zhang *et al.*, 2008). In the crystalline state, the neighboring molecules are linked furtherly by $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonding forming supermolecular structure(Fig. 2).

S2. Experimental

$\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.1904 g, 0.8 mmol), biimidazole (0.107 g, 1 mmol), and water (10 ml) were added to an aqueous solution (5 ml) containing NaN_3 (0.028 g,0.4 mmol). The resulting mixture was further stirred for 15 min in air, and then transferred and sealed in a 20 ml Teflon-lined reactor, which was heated at 423 K for 4 days and then cooled to room temperature at a rate of 5 K h^{-1} . Red blocks of (I) were obtained and washed with water.

S3. Refinement

H atoms attached to C and N atoms of (I) were placed in geometrically idealized positions ($\text{C}-\text{H} = 0.93\text{\AA}$, $\text{N}-\text{H} = 0.86\text{\AA}$ and constrained to ride on their parent atoms.

**Figure 1**

A view of the structure of (I) with displacement ellipsoids drawn at the 30% probability level.




Figure 2
hydrogen bond interaction in neighboring molecules.

Tris[2-(1*H*-imidazol-2-yl)imidazol-1-ido]cobalt(III)

Crystal data

$[\text{Co}(\text{C}_6\text{H}_5\text{N}_4)_3]$
 $M_r = 458.35$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 12.299 (3) \text{ \AA}$
 $b = 12.524 (3) \text{ \AA}$
 $c = 12.932 (3) \text{ \AA}$
 $\beta = 97.773 (4)^\circ$
 $V = 1973.6 (8) \text{ \AA}^3$
 $Z = 4$

$F(000) = 936$
 $D_x = 1.543 \text{ Mg m}^{-3}$
 $\text{Mo } K\alpha \text{ radiation, } \lambda = 0.71073 \text{ \AA}$
Cell parameters from 10212 reflections
 $\theta = 2.3\text{--}20.5^\circ$
 $\mu = 0.90 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
Block, red
 $0.5 \times 0.4 \times 0.3 \text{ mm}$

Data collection

Bruker SMART CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2000)
 $T_{\min} = 0.654$, $T_{\max} = 0.762$

10212 measured reflections
3728 independent reflections
2358 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.046$
 $\theta_{\max} = 25.7^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -14 \rightarrow 15$
 $k = -15 \rightarrow 13$
 $l = -15 \rightarrow 13$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.041$$

$$wR(F^2) = 0.138$$

$$S = 0.98$$

3728 reflections

280 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0798P)^2]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.48 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.49 \text{ e \AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.14643 (4)	0.83171 (4)	0.84581 (4)	0.03246 (19)
N1	0.1905 (2)	0.8669 (2)	0.7123 (2)	0.0325 (7)
N2	0.0080 (2)	0.8920 (2)	0.7814 (2)	0.0348 (7)
N3	0.2012 (2)	0.9650 (2)	0.9055 (2)	0.0324 (7)
N4	0.2936 (2)	0.7858 (2)	0.8922 (2)	0.0344 (7)
N5	0.1019 (2)	0.7861 (2)	0.9773 (2)	0.0348 (8)
N6	0.0917 (2)	0.6922 (2)	0.8032 (3)	0.0365 (8)
C5	0.2563 (3)	0.9108 (3)	0.5682 (3)	0.0400 (10)
H2	0.3051	0.9191	0.5197	0.048*
C6	0.2817 (3)	0.8656 (3)	0.6656 (3)	0.0396 (10)
H1	0.3498	0.8388	0.6939	0.048*
N12	0.1482 (2)	0.9419 (2)	0.5534 (2)	0.0360 (8)
C4	0.1135 (3)	0.9131 (3)	0.6429 (3)	0.0308 (8)
C3	0.0105 (3)	0.9241 (3)	0.6834 (3)	0.0343 (9)
N11	-0.0888 (2)	0.9582 (2)	0.6405 (3)	0.0433 (9)
H6	-0.1059	0.9829	0.5783	0.052*
C2	-0.1578 (3)	0.9462 (4)	0.7141 (4)	0.0516 (12)
H7	-0.2322	0.9627	0.7060	0.062*
C1	-0.0984 (3)	0.9061 (3)	0.8007 (3)	0.0435 (10)
H8	-0.1249	0.8905	0.8630	0.052*
C18	0.0637 (4)	0.6367 (3)	0.7131 (3)	0.0477 (11)
H9	0.0782	0.6578	0.6473	0.057*
C17	0.0115 (4)	0.5462 (3)	0.7345 (3)	0.0493 (11)
H10	-0.0153	0.4939	0.6866	0.059*
N7	0.0051 (3)	0.5451 (2)	0.8392 (3)	0.0397 (8)

H11	-0.0243	0.4963	0.8731	0.048*
C16	0.0534 (3)	0.6349 (3)	0.8788 (3)	0.0322 (9)
C15	0.0665 (3)	0.6837 (3)	0.9789 (3)	0.0306 (8)
N8	0.0546 (3)	0.6485 (2)	1.0734 (2)	0.0412 (8)
C14	0.0858 (3)	0.7350 (3)	1.1374 (3)	0.0477 (11)
H15	0.0868	0.7361	1.2094	0.057*
C13	0.1150 (3)	0.8185 (3)	1.0787 (3)	0.0392 (10)
H16	0.1393	0.8852	1.1036	0.047*
C12	0.3600 (3)	0.7001 (3)	0.8784 (3)	0.0429 (10)
H17	0.3368	0.6345	0.8496	0.051*
C11	0.4643 (3)	0.7274 (3)	0.9139 (3)	0.0469 (11)
H18	0.5256	0.6836	0.9151	0.056*
N9	0.4642 (3)	0.8312 (2)	0.9480 (3)	0.0413 (8)
H19	0.5204	0.8683	0.9734	0.050*
C10	0.3598 (3)	0.8634 (3)	0.9339 (3)	0.0331 (9)
C9	0.3067 (3)	0.9641 (3)	0.9483 (3)	0.0333 (9)
N10	0.3413 (2)	1.0555 (2)	0.9935 (3)	0.0368 (8)
C8	0.2507 (3)	1.1201 (3)	0.9770 (3)	0.0419 (10)
H23	0.2482	1.1907	0.9988	0.050*
C7	0.1643 (3)	1.0652 (3)	0.9236 (3)	0.0394 (9)
H24	0.0939	1.0913	0.9034	0.047*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0365 (3)	0.0314 (3)	0.0284 (3)	-0.0033 (2)	0.0010 (2)	0.0038 (2)
N1	0.0327 (16)	0.0320 (16)	0.0321 (19)	-0.0013 (13)	0.0023 (14)	0.0052 (14)
N2	0.0354 (17)	0.0352 (18)	0.033 (2)	-0.0026 (14)	0.0028 (14)	0.0041 (15)
N3	0.0359 (17)	0.0304 (17)	0.030 (2)	-0.0012 (13)	0.0013 (14)	0.0040 (13)
N4	0.0396 (17)	0.0262 (16)	0.036 (2)	-0.0012 (14)	-0.0009 (15)	0.0041 (14)
N5	0.0377 (17)	0.0316 (17)	0.035 (2)	-0.0041 (14)	0.0021 (15)	0.0008 (14)
N6	0.0437 (18)	0.0348 (17)	0.030 (2)	-0.0040 (14)	0.0017 (15)	0.0001 (14)
C5	0.040 (2)	0.046 (2)	0.035 (3)	-0.0064 (18)	0.0082 (19)	0.0010 (19)
C6	0.034 (2)	0.042 (2)	0.041 (3)	-0.0004 (17)	0.0028 (19)	0.0017 (19)
N12	0.0379 (18)	0.0370 (18)	0.032 (2)	-0.0054 (14)	0.0008 (15)	0.0026 (14)
C4	0.0361 (19)	0.0306 (19)	0.025 (2)	-0.0051 (16)	0.0013 (16)	0.0005 (16)
C3	0.038 (2)	0.033 (2)	0.031 (2)	-0.0034 (17)	0.0056 (18)	0.0003 (17)
N11	0.0374 (18)	0.055 (2)	0.036 (2)	0.0019 (15)	0.0008 (16)	0.0097 (16)
C2	0.034 (2)	0.071 (3)	0.051 (3)	0.004 (2)	0.009 (2)	0.010 (2)
C1	0.039 (2)	0.057 (3)	0.037 (3)	-0.0025 (19)	0.0129 (19)	0.006 (2)
C18	0.074 (3)	0.046 (2)	0.024 (2)	-0.006 (2)	0.010 (2)	-0.0032 (19)
C17	0.072 (3)	0.046 (3)	0.029 (3)	-0.013 (2)	0.003 (2)	-0.0085 (19)
N7	0.051 (2)	0.0336 (18)	0.034 (2)	-0.0088 (15)	0.0043 (16)	-0.0024 (15)
C16	0.040 (2)	0.033 (2)	0.024 (2)	-0.0040 (16)	0.0040 (17)	0.0020 (16)
C15	0.0329 (19)	0.030 (2)	0.028 (2)	-0.0031 (15)	0.0023 (16)	0.0038 (16)
N8	0.056 (2)	0.0400 (19)	0.027 (2)	-0.0081 (15)	0.0056 (17)	0.0011 (15)
C14	0.069 (3)	0.051 (3)	0.024 (2)	-0.007 (2)	0.010 (2)	-0.004 (2)
C13	0.051 (2)	0.037 (2)	0.029 (2)	-0.0044 (18)	0.0035 (19)	-0.0078 (18)

C12	0.056 (3)	0.030 (2)	0.041 (3)	0.0033 (18)	0.000 (2)	0.0027 (18)
C11	0.050 (2)	0.042 (3)	0.046 (3)	0.014 (2)	-0.003 (2)	0.006 (2)
N9	0.0407 (18)	0.0404 (19)	0.039 (2)	0.0016 (15)	-0.0083 (16)	-0.0009 (15)
C10	0.033 (2)	0.034 (2)	0.030 (2)	-0.0009 (16)	-0.0031 (17)	0.0073 (17)
C9	0.038 (2)	0.031 (2)	0.030 (2)	-0.0019 (16)	0.0032 (18)	0.0034 (16)
N10	0.0440 (18)	0.0334 (18)	0.034 (2)	-0.0023 (14)	0.0071 (15)	-0.0058 (14)
C8	0.053 (2)	0.033 (2)	0.043 (3)	-0.0008 (19)	0.016 (2)	-0.0025 (19)
C7	0.043 (2)	0.036 (2)	0.041 (3)	0.0060 (18)	0.0120 (19)	0.0044 (18)

Geometric parameters (\AA , $^\circ$)

Co1—N4	1.917 (3)	C2—C1	1.348 (5)
Co1—N3	1.922 (3)	C2—H7	0.9300
Co1—N6	1.926 (3)	C1—H8	0.9300
Co1—N1	1.929 (3)	C18—C17	1.350 (5)
Co1—N5	1.941 (3)	C18—H9	0.9300
Co1—N2	1.944 (3)	C17—N7	1.367 (5)
N1—C6	1.344 (4)	C17—H10	0.9300
N1—C4	1.345 (4)	N7—C16	1.341 (4)
N2—C3	1.333 (5)	N7—H11	0.8600
N2—C1	1.376 (4)	C16—C15	1.420 (5)
N3—C9	1.340 (4)	C15—N8	1.326 (5)
N3—C7	1.365 (4)	N8—C14	1.386 (5)
N4—C10	1.335 (5)	C14—C13	1.369 (5)
N4—C12	1.374 (4)	C14—H15	0.9300
N5—C15	1.356 (4)	C13—H16	0.9300
N5—C13	1.361 (5)	C12—C11	1.347 (5)
N6—C16	1.349 (5)	C12—H17	0.9300
N6—C18	1.361 (5)	C11—N9	1.373 (5)
C5—N12	1.373 (4)	C11—H18	0.9300
C5—C6	1.378 (5)	N9—C10	1.334 (5)
C5—H2	0.9300	N9—H19	0.8600
C6—H1	0.9300	C10—C9	1.443 (5)
N12—C4	1.336 (5)	C9—N10	1.329 (4)
C4—C3	1.442 (5)	N10—C8	1.370 (5)
C3—N11	1.341 (5)	C8—C7	1.372 (5)
N11—C2	1.367 (5)	C8—H23	0.9300
N11—H6	0.8600	C7—H24	0.9300
N4—Co1—N3	82.54 (12)	C1—C2—H7	126.2
N4—Co1—N6	95.52 (13)	N11—C2—H7	126.2
N3—Co1—N6	173.01 (13)	C2—C1—N2	108.6 (4)
N4—Co1—N1	88.88 (12)	C2—C1—H8	125.7
N3—Co1—N1	92.05 (13)	N2—C1—H8	125.7
N6—Co1—N1	94.62 (13)	C17—C18—N6	109.0 (4)
N4—Co1—N5	90.23 (13)	C17—C18—H9	125.5
N3—Co1—N5	91.61 (13)	N6—C18—H9	125.5
N6—Co1—N5	81.67 (13)	C18—C17—N7	107.7 (4)

N1—Co1—N5	176.08 (12)	C18—C17—H10	126.2
N4—Co1—N2	170.39 (13)	N7—C17—H10	126.2
N3—Co1—N2	94.22 (12)	C16—N7—C17	106.8 (3)
N6—Co1—N2	88.73 (13)	C16—N7—H11	126.6
N1—Co1—N2	82.18 (12)	C17—N7—H11	126.6
N5—Co1—N2	98.93 (12)	N7—C16—N6	110.4 (3)
C6—N1—C4	105.1 (3)	N7—C16—C15	134.4 (3)
C6—N1—Co1	138.8 (3)	N6—C16—C15	115.1 (3)
C4—N1—Co1	116.0 (2)	N8—C15—N5	113.9 (3)
C3—N2—C1	106.1 (3)	N8—C15—C16	133.1 (3)
C3—N2—Co1	113.1 (2)	N5—C15—C16	113.0 (3)
C1—N2—Co1	140.5 (3)	C15—N8—C14	103.5 (3)
C9—N3—C7	105.2 (3)	C13—C14—N8	109.8 (4)
C9—N3—Co1	115.2 (2)	C13—C14—H15	125.1
C7—N3—Co1	139.5 (3)	N8—C14—H15	125.1
C10—N4—C12	106.3 (3)	N5—C13—C14	107.5 (3)
C10—N4—Co1	114.0 (2)	N5—C13—H16	126.2
C12—N4—Co1	138.2 (3)	C14—C13—H16	126.2
C15—N5—C13	105.3 (3)	C11—C12—N4	108.2 (3)
C15—N5—Co1	114.9 (3)	C11—C12—H17	125.9
C13—N5—Co1	138.2 (3)	N4—C12—H17	125.9
C16—N6—C18	106.1 (3)	C12—C11—N9	107.9 (3)
C16—N6—Co1	114.7 (3)	C12—C11—H18	126.1
C18—N6—Co1	138.3 (3)	N9—C11—H18	126.1
N12—C5—C6	109.9 (3)	C10—N9—C11	106.6 (3)
N12—C5—H2	125.1	C10—N9—H19	126.7
C6—C5—H2	125.1	C11—N9—H19	126.7
N1—C6—C5	107.8 (3)	N9—C10—N4	111.0 (3)
N1—C6—H1	126.1	N9—C10—C9	133.7 (3)
C5—C6—H1	126.1	N4—C10—C9	115.2 (3)
C4—N12—C5	102.7 (3)	N10—C9—N3	114.1 (3)
N12—C4—N1	114.6 (3)	N10—C9—C10	133.3 (3)
N12—C4—C3	133.6 (3)	N3—C9—C10	112.6 (3)
N1—C4—C3	111.8 (3)	C9—N10—C8	103.6 (3)
N2—C3—N11	110.7 (3)	N10—C8—C7	109.9 (3)
N2—C3—C4	116.8 (3)	N10—C8—H23	125.0
N11—C3—C4	132.5 (4)	C7—C8—H23	125.0
C3—N11—C2	107.0 (3)	N3—C7—C8	107.1 (3)
C3—N11—H6	126.5	N3—C7—H24	126.4
C2—N11—H6	126.5	C8—C7—H24	126.4
C1—C2—N11	107.6 (3)		
N4—Co1—N1—C6	-1.6 (4)	C1—N2—C3—C4	-177.6 (3)
N3—Co1—N1—C6	80.9 (4)	Co1—N2—C3—C4	-2.2 (4)
N6—Co1—N1—C6	-97.0 (4)	N12—C4—C3—N2	-175.1 (4)
N2—Co1—N1—C6	174.9 (4)	N1—C4—C3—N2	2.5 (5)
N4—Co1—N1—C4	-176.0 (3)	N12—C4—C3—N11	7.4 (7)
N3—Co1—N1—C4	-93.5 (3)	N1—C4—C3—N11	-174.9 (4)

N6—Co1—N1—C4	88.5 (3)	N2—C3—N11—C2	-0.7 (4)
N2—Co1—N1—C4	0.4 (2)	C4—C3—N11—C2	176.9 (4)
N3—Co1—N2—C3	92.5 (3)	C3—N11—C2—C1	0.7 (5)
N6—Co1—N2—C3	-93.8 (3)	N11—C2—C1—N2	-0.4 (5)
N1—Co1—N2—C3	1.0 (2)	C3—N2—C1—C2	0.0 (5)
N5—Co1—N2—C3	-175.2 (2)	Co1—N2—C1—C2	-173.3 (3)
N3—Co1—N2—C1	-94.5 (4)	C16—N6—C18—C17	-1.3 (5)
N6—Co1—N2—C1	79.2 (4)	Co1—N6—C18—C17	-169.1 (3)
N1—Co1—N2—C1	174.0 (4)	N6—C18—C17—N7	0.8 (5)
N5—Co1—N2—C1	-2.2 (4)	C18—C17—N7—C16	0.1 (5)
N4—Co1—N3—C9	-0.3 (3)	C17—N7—C16—N6	-0.9 (4)
N1—Co1—N3—C9	-88.9 (3)	C17—N7—C16—C15	174.7 (4)
N5—Co1—N3—C9	89.8 (3)	C18—N6—C16—N7	1.3 (4)
N2—Co1—N3—C9	-171.2 (3)	Co1—N6—C16—N7	172.5 (2)
N4—Co1—N3—C7	-176.5 (4)	C18—N6—C16—C15	-175.2 (3)
N1—Co1—N3—C7	94.9 (4)	Co1—N6—C16—C15	-4.0 (4)
N5—Co1—N3—C7	-86.5 (4)	C13—N5—C15—N8	0.8 (4)
N2—Co1—N3—C7	12.6 (4)	Co1—N5—C15—N8	169.0 (2)
N3—Co1—N4—C10	-3.9 (3)	C13—N5—C15—C16	-177.1 (3)
N6—Co1—N4—C10	-177.1 (3)	Co1—N5—C15—C16	-8.9 (4)
N1—Co1—N4—C10	88.3 (3)	N7—C16—C15—N8	15.6 (7)
N5—Co1—N4—C10	-95.5 (3)	N6—C16—C15—N8	-169.0 (4)
N3—Co1—N4—C12	-167.7 (4)	N7—C16—C15—N5	-167.0 (4)
N6—Co1—N4—C12	19.0 (4)	N6—C16—C15—N5	8.5 (5)
N1—Co1—N4—C12	-75.5 (4)	N5—C15—N8—C14	-0.6 (4)
N5—Co1—N4—C12	100.7 (4)	C16—C15—N8—C14	176.8 (4)
N4—Co1—N5—C15	-90.1 (3)	C15—N8—C14—C13	0.1 (5)
N3—Co1—N5—C15	-172.6 (3)	C15—N5—C13—C14	-0.7 (4)
N6—Co1—N5—C15	5.5 (2)	Co1—N5—C13—C14	-164.5 (3)
N2—Co1—N5—C15	92.8 (3)	N8—C14—C13—N5	0.4 (5)
N4—Co1—N5—C13	72.7 (4)	C10—N4—C12—C11	1.1 (4)
N3—Co1—N5—C13	-9.9 (4)	Co1—N4—C12—C11	165.8 (3)
N6—Co1—N5—C13	168.2 (4)	N4—C12—C11—N9	-1.3 (5)
N2—Co1—N5—C13	-104.4 (4)	C12—C11—N9—C10	1.0 (4)
N4—Co1—N6—C16	88.8 (3)	C11—N9—C10—N4	-0.3 (4)
N1—Co1—N6—C16	178.1 (3)	C11—N9—C10—C9	-175.9 (4)
N5—Co1—N6—C16	-0.7 (3)	C12—N4—C10—N9	-0.5 (4)
N2—Co1—N6—C16	-99.9 (3)	Co1—N4—C10—N9	-169.3 (3)
N4—Co1—N6—C18	-104.1 (4)	C12—N4—C10—C9	176.0 (3)
N1—Co1—N6—C18	-14.8 (4)	Co1—N4—C10—C9	7.1 (4)
N5—Co1—N6—C18	166.5 (4)	C7—N3—C9—N10	0.6 (4)
N2—Co1—N6—C18	67.3 (4)	Co1—N3—C9—N10	-176.9 (2)
C4—N1—C6—C5	-0.4 (4)	C7—N3—C9—C10	-178.5 (3)
Co1—N1—C6—C5	-175.3 (3)	Co1—N3—C9—C10	4.0 (4)
N12—C5—C6—N1	0.5 (4)	N9—C10—C9—N10	-10.7 (8)
C6—C5—N12—C4	-0.3 (4)	N4—C10—C9—N10	173.8 (4)
C5—N12—C4—N1	0.1 (4)	N9—C10—C9—N3	168.2 (4)
C5—N12—C4—C3	177.6 (4)	N4—C10—C9—N3	-7.3 (5)

C6—N1—C4—N12	0.2 (4)	N3—C9—N10—C8	-0.8 (4)
Co1—N1—C4—N12	176.4 (2)	C10—C9—N10—C8	178.1 (4)
C6—N1—C4—C3	-177.9 (3)	C9—N10—C8—C7	0.7 (4)
Co1—N1—C4—C3	-1.7 (4)	C9—N3—C7—C8	-0.1 (4)
C1—N2—C3—N11	0.4 (4)	Co1—N3—C7—C8	176.4 (3)
Co1—N2—C3—N11	175.8 (2)	N10—C8—C7—N3	-0.4 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N11—H6···N12 ⁱ	0.86	1.95	2.808 (4)	172
N7—H11···N8 ⁱⁱ	0.86	1.99	2.814 (4)	159
N9—H19···N10 ⁱⁱⁱ	0.86	1.95	2.796 (4)	169

Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $-x, -y+1, -z+2$; (iii) $-x+1, -y+2, -z+2$.