

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(4-Methylbenzoyl)benzene-sulfonamide

P. A. Suchetan,^a B. Thimme Gowda,^{a*} Sabine Foro^b and Hartmut Fuess^b

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 18 June 2010; accepted 20 June 2010

Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.109; data-to-parameter ratio = 15.0.

In the title compound, $\text{C}_{14}\text{H}_{13}\text{NO}_3\text{S}$, the conformation of the N—H bond in the C—SO₂—NH—C(O) segment is *anti* to the C=O bond. The dihedral angle between the sulfonyl benzene ring and the S—N—C—O segment (r.m.s. deviation = 0.039 Å) is 77.1 (1)° and that between the sulfonyl and benzoyl benzene rings is 71.9 (1)°.

Related literature

For background to our study of the effect of ring and side-chain substituents on the crystal structures of *N*-aromatic sulfonamides and for related structures, see: Gowda *et al.* (2009); Suchetan *et al.* (2009, 2010).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{13}\text{NO}_3\text{S}$	$\gamma = 81.949$ (9)°
$M_r = 275.31$	$V = 638.36$ (11) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 5.5519$ (6) Å	Mo $K\alpha$ radiation
$b = 10.541$ (1) Å	$\mu = 0.26$ mm ⁻¹
$c = 11.105$ (1) Å	$T = 299$ K
$\alpha = 85.654$ (9)°	$0.40 \times 0.32 \times 0.16$ mm
$\beta = 83.667$ (9)°	

Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector	Diffraction, 2009)
Absorption correction: multi-scan (<i>CrysAlis RED</i> ; Oxford)	$T_{\min} = 0.904$, $T_{\max} = 0.960$
	4293 measured reflections
	2595 independent reflections
	2207 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	173 parameters
$wR(F^2) = 0.109$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.24$ e Å ⁻³
2595 reflections	$\Delta\rho_{\text{min}} = -0.30$ e Å ⁻³

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

PAS thanks the Council of Scientific and Industrial Research (CSIR), Government of India, New Delhi, for the award of a research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5108).

References

- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2009). *Acta Cryst.* **E65**, o2516.
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2009). *Acta Cryst.* **E65**, o3156.
 Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2010). *Acta Cryst.* **E66**, o1039.

supporting information

Acta Cryst. (2010). E66, o1772 [doi:10.1107/S1600536810023974]

***N*-(4-Methylbenzoyl)benzenesulfonamide**

P. A. Suchetan, B. Thimme Gowda, Sabine Foro and Hartmut Fues

S1. Comment

As a part of studying the effect of ring and the side chain substituents on the crystal structures of *N*-aromatic sulfonamides (Gowda *et al.*, 2009; Suchetan *et al.*, 2009, 2010), the crystal structure of *N*-(4-methylbenzoyl)benzenesulfonamide has been determined (Fig. 1). The conformation of the N—H bond in the C—SO₂—NH—C(O) segment is *anti* to the C=O bond, similar to those observed in *N*-(benzoyl)benzenesulfonamide (II) (Gowda *et al.*, 2009), *N*-(benzoyl)-4-methylbenzenesulfonamide (III) (Suchetan *et al.*, 2010) and *N*-(4-chlorobenzoyl)-benzenesulfonamide (IV) (Suchetan *et al.*, 2009).

The molecules are twisted at the S—N bonds with the C—SO₂—NH—C torsional angle of 67.4 (1)°, compared to the values of -66.9 (3)° in (II), 73.2 (2)° in (III) and 69.4 (2)° in (IV).

The dihedral angle between the sulfonyl-bound benzene ring and the S—N—C—O segment (r.m.s. deviation 0.039 Å) is 77.1 (1)°, compared to the values of 86.5 (1)° in (II), 76.5 (1)° in (III) and 75.7 (1)° in (IV).


The dihedral angle between the sulfonyl and the benzoyl benzene rings is 71.9 (1)°, compared to the values of 80.3 (1)° in (II), 79.4 (1)° in (III), and 68.6 (1)° in (IV).

S2. Experimental

The title compound was prepared by refluxing a mixture of 4-methylbenzoic acid, benzenesulfonamide and phosphorous oxy chloride for 3 h on a water bath. The resultant mixture was cooled and poured into ice cold water. The solid obtained was filtered, washed thoroughly with water and then dissolved in a sodium bicarbonate solution. The compound was later reprecipitated by acidifying the filtered solution with dilute HCl. It was filtered, dried and recrystallized. Colourless needle-shaped single crystals of the title compound used in X-ray diffraction studies were obtained by slow evaporation of its toluene solution at room temperature.

S3. Refinement

H atoms were positioned with idealized geometry using a riding model with N—H = 0.86 Å, C—H = 0.93–0.96 Å and were refined with isotropic displacement parameters (set to 1.2 times of the U_{eq} of the parent atom).

**Figure 1**

Molecular structure of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.

***N*-(4-Methylbenzoyl)benzenesulfonamide**

Crystal data

$C_{14}H_{13}NO_3S$

$M_r = 275.31$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.5519$ (6) Å

$b = 10.541$ (1) Å

$c = 11.105$ (1) Å

$\alpha = 85.654$ (9)°

$\beta = 83.667$ (9)°

$\gamma = 81.949$ (9)°

$V = 638.36$ (11) Å³

$Z = 2$

$F(000) = 288$

$D_x = 1.432$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2679 reflections

$\theta = 2.6$ – 27.7 °

$\mu = 0.26$ mm⁻¹

$T = 299$ K

Needle, colourless

$0.40 \times 0.32 \times 0.16$ mm

Data collection

Oxford Diffraction Xcalibur

diffractometer with a Sapphire CCD detector

Radiation source: fine-focus sealed tube

Graphite monochromator

Rotation method data acquisition using ω and φ scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.904$, $T_{\max} = 0.960$

4293 measured reflections

2595 independent reflections

2207 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 26.4$ °, $\theta_{\min} = 2.6$ °

$h = -6 \rightarrow 6$

$k = -13 \rightarrow 11$

$l = -13 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.109$

$S = 1.08$

2595 reflections

173 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0564P)^2 + 0.1768P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.020$

$$\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0535 (3)	0.17087 (15)	0.37945 (14)	0.0370 (3)
C2	0.2512 (3)	0.07861 (18)	0.39763 (17)	0.0491 (4)
H2	0.3709	0.0944	0.4448	0.059*
C3	0.2687 (4)	-0.03734 (19)	0.34486 (19)	0.0557 (5)
H3	0.4008	-0.1001	0.3567	0.067*
C4	0.0919 (4)	-0.06031 (17)	0.27507 (18)	0.0514 (5)
H4	0.1050	-0.1383	0.2394	0.062*
C5	-0.1047 (4)	0.03194 (19)	0.25778 (17)	0.0517 (5)
H5	-0.2241	0.0157	0.2106	0.062*
C6	-0.1262 (3)	0.14869 (17)	0.31001 (16)	0.0439 (4)
H6	-0.2592	0.2110	0.2985	0.053*
C7	0.1508 (3)	0.45512 (14)	0.24545 (14)	0.0362 (3)
C8	0.2982 (3)	0.55415 (14)	0.18611 (13)	0.0345 (3)
C9	0.4790 (3)	0.60283 (16)	0.23892 (15)	0.0393 (4)
H9	0.5113	0.5748	0.3177	0.047*
C10	0.6111 (3)	0.69266 (17)	0.17506 (15)	0.0429 (4)
H10	0.7328	0.7235	0.2113	0.051*
C11	0.5653 (3)	0.73765 (16)	0.05771 (15)	0.0403 (4)
C12	0.3825 (3)	0.69012 (18)	0.00681 (16)	0.0471 (4)
H12	0.3477	0.7197	-0.0712	0.056*
C13	0.2508 (3)	0.60001 (17)	0.06897 (15)	0.0451 (4)
H13	0.1291	0.5695	0.0324	0.054*
C14	0.7134 (4)	0.83301 (19)	-0.01353 (19)	0.0553 (5)
H14A	0.8409	0.7882	-0.0661	0.066*
H14B	0.7844	0.8795	0.0415	0.066*
H14C	0.6090	0.8920	-0.0613	0.066*
N1	0.2026 (3)	0.40757 (13)	0.36175 (13)	0.0466 (4)
H1N	0.3319	0.4262	0.3887	0.056*
O1	0.1439 (3)	0.29602 (14)	0.55935 (12)	0.0705 (5)
O2	-0.2229 (3)	0.37520 (13)	0.45410 (13)	0.0618 (4)
O3	-0.0051 (2)	0.41356 (12)	0.19791 (11)	0.0493 (3)
S1	0.02709 (9)	0.31703 (4)	0.45017 (4)	0.04709 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0466 (9)	0.0344 (8)	0.0314 (8)	-0.0135 (7)	-0.0018 (6)	0.0008 (6)
C2	0.0481 (10)	0.0518 (10)	0.0501 (10)	-0.0113 (8)	-0.0125 (8)	-0.0001 (8)
C3	0.0556 (11)	0.0452 (10)	0.0633 (12)	-0.0003 (8)	-0.0039 (9)	0.0008 (9)
C4	0.0641 (12)	0.0386 (9)	0.0524 (11)	-0.0164 (8)	0.0059 (9)	-0.0079 (8)
C5	0.0555 (11)	0.0545 (11)	0.0510 (11)	-0.0219 (9)	-0.0081 (8)	-0.0105 (8)
C6	0.0453 (9)	0.0427 (9)	0.0453 (9)	-0.0096 (7)	-0.0076 (7)	-0.0014 (7)
C7	0.0448 (9)	0.0310 (8)	0.0330 (8)	-0.0035 (6)	-0.0065 (6)	-0.0027 (6)
C8	0.0391 (8)	0.0319 (8)	0.0313 (8)	-0.0004 (6)	-0.0035 (6)	-0.0018 (6)
C9	0.0429 (9)	0.0439 (9)	0.0313 (8)	-0.0054 (7)	-0.0071 (6)	0.0016 (6)
C10	0.0417 (9)	0.0476 (9)	0.0408 (9)	-0.0092 (7)	-0.0061 (7)	-0.0030 (7)
C11	0.0419 (9)	0.0359 (8)	0.0397 (8)	0.0000 (7)	0.0026 (7)	-0.0005 (6)
C12	0.0571 (11)	0.0495 (10)	0.0342 (8)	-0.0070 (8)	-0.0097 (7)	0.0074 (7)
C13	0.0524 (10)	0.0494 (10)	0.0366 (9)	-0.0127 (8)	-0.0145 (7)	0.0030 (7)
C14	0.0585 (12)	0.0500 (11)	0.0552 (11)	-0.0114 (9)	0.0036 (9)	0.0053 (9)
N1	0.0680 (10)	0.0420 (8)	0.0359 (7)	-0.0246 (7)	-0.0154 (7)	0.0053 (6)
O1	0.1269 (14)	0.0621 (9)	0.0333 (7)	-0.0439 (9)	-0.0218 (7)	0.0059 (6)
O2	0.0745 (10)	0.0493 (8)	0.0565 (8)	-0.0036 (7)	0.0148 (7)	-0.0111 (6)
O3	0.0544 (7)	0.0519 (7)	0.0459 (7)	-0.0172 (6)	-0.0166 (6)	0.0056 (5)
S1	0.0756 (3)	0.0393 (2)	0.0295 (2)	-0.0204 (2)	-0.00330 (19)	-0.00178 (16)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.381 (2)	C9—C10	1.384 (2)
C1—C2	1.382 (3)	C9—H9	0.93
C1—S1	1.7626 (16)	C10—C11	1.391 (2)
C2—C3	1.382 (3)	C10—H10	0.93
C2—H2	0.93	C11—C12	1.383 (2)
C3—C4	1.373 (3)	C11—C14	1.510 (2)
C3—H3	0.93	C12—C13	1.377 (2)
C4—C5	1.376 (3)	C12—H12	0.93
C4—H4	0.93	C13—H13	0.93
C5—C6	1.385 (3)	C14—H14A	0.96
C5—H5	0.93	C14—H14B	0.96
C6—H6	0.93	C14—H14C	0.96
C7—O3	1.209 (2)	N1—S1	1.6529 (15)
C7—N1	1.395 (2)	N1—H1N	0.86
C7—C8	1.487 (2)	O1—S1	1.4257 (14)
C8—C9	1.391 (2)	O2—S1	1.4340 (15)
C8—C13	1.393 (2)		
C6—C1—C2	121.09 (16)	C9—C10—C11	121.20 (16)
C6—C1—S1	119.75 (13)	C9—C10—H10	119.4
C2—C1—S1	119.13 (13)	C11—C10—H10	119.4
C3—C2—C1	119.19 (17)	C12—C11—C10	117.86 (15)
C3—C2—H2	120.4	C12—C11—C14	120.74 (16)

C1—C2—H2	120.4	C10—C11—C14	121.39 (16)
C4—C3—C2	120.29 (18)	C13—C12—C11	121.56 (16)
C4—C3—H3	119.9	C13—C12—H12	119.2
C2—C3—H3	119.9	C11—C12—H12	119.2
C3—C4—C5	120.12 (17)	C12—C13—C8	120.55 (16)
C3—C4—H4	119.9	C12—C13—H13	119.7
C5—C4—H4	119.9	C8—C13—H13	119.7
C4—C5—C6	120.56 (17)	C11—C14—H14A	109.5
C4—C5—H5	119.7	C11—C14—H14B	109.5
C6—C5—H5	119.7	H14A—C14—H14B	109.5
C1—C6—C5	118.75 (17)	C11—C14—H14C	109.5
C1—C6—H6	120.6	H14A—C14—H14C	109.5
C5—C6—H6	120.6	H14B—C14—H14C	109.5
O3—C7—N1	119.51 (15)	C7—N1—S1	123.07 (12)
O3—C7—C8	123.77 (14)	C7—N1—H1N	118.5
N1—C7—C8	116.70 (14)	S1—N1—H1N	118.5
C9—C8—C13	118.37 (15)	O1—S1—O2	119.65 (10)
C9—C8—C7	124.69 (14)	O1—S1—N1	103.47 (8)
C13—C8—C7	116.93 (14)	O2—S1—N1	109.71 (8)
C10—C9—C8	120.45 (15)	O1—S1—C1	109.18 (9)
C10—C9—H9	119.8	O2—S1—C1	108.24 (8)
C8—C9—H9	119.8	N1—S1—C1	105.73 (8)
<hr/>			
C6—C1—C2—C3	0.2 (3)	C10—C11—C12—C13	-0.7 (3)
S1—C1—C2—C3	178.30 (14)	C14—C11—C12—C13	177.89 (17)
C1—C2—C3—C4	0.1 (3)	C11—C12—C13—C8	0.2 (3)
C2—C3—C4—C5	-0.3 (3)	C9—C8—C13—C12	0.8 (3)
C3—C4—C5—C6	0.2 (3)	C7—C8—C13—C12	-178.80 (16)
C2—C1—C6—C5	-0.4 (3)	O3—C7—N1—S1	-12.2 (2)
S1—C1—C6—C5	-178.42 (13)	C8—C7—N1—S1	168.97 (11)
C4—C5—C6—C1	0.2 (3)	C7—N1—S1—O1	-177.83 (14)
O3—C7—C8—C9	-179.72 (16)	C7—N1—S1—O2	-49.07 (16)
N1—C7—C8—C9	-0.9 (2)	C7—N1—S1—C1	67.44 (15)
O3—C7—C8—C13	-0.1 (2)	C6—C1—S1—O1	149.92 (14)
N1—C7—C8—C13	178.70 (15)	C2—C1—S1—O1	-28.17 (17)
C13—C8—C9—C10	-1.3 (2)	C6—C1—S1—O2	18.16 (16)
C7—C8—C9—C10	178.29 (15)	C2—C1—S1—O2	-159.93 (14)
C8—C9—C10—C11	0.8 (3)	C6—C1—S1—N1	-99.34 (14)
C9—C10—C11—C12	0.2 (3)	C2—C1—S1—N1	82.58 (15)
C9—C10—C11—C14	-178.37 (16)		
