

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (2E)-1-(2-Bromophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one

 Jerry P. Jasinski,<sup>a\*</sup> Ray J. Butcher,<sup>b</sup> K. Veena,<sup>c</sup>  
 B. Narayana<sup>c</sup> and H. S. Yathirajan<sup>d</sup>

<sup>a</sup>Department of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, <sup>b</sup>Department of Chemistry, Howard University, 525 College Street NW, Washington DC 20059, USA, <sup>c</sup>Department of Studies in Chemistry, Mangalore University, Mangalagangothri, 574 199, India, and <sup>d</sup>Department of Studies in Chemistry, University of Mysore, Manasagangothri, Mysore 570 006, India  
 Correspondence e-mail: jjasinski@keene.edu


Received 3 June 2010; accepted 9 June 2010

Key indicators: single-crystal X-ray study;  $T = 110$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.034;  $wR$  factor = 0.091; data-to-parameter ratio = 9.5.

In the title compound,  $\text{C}_{16}\text{H}_{13}\text{BrO}_2$ , two benzene rings form a dihedral angle of  $44.3(9)^\circ$ . In the crystal, weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules into chains propagating in  $[010]$ . The crystal packing also exhibits short  $\text{Br}\cdots\text{Br}$  contacts of  $3.4787(8)$  Å. A comparison of the DFT-optimized gas-phase molecular geometry with that in the crystal structure revealed only small differences.

## Related literature

For the radical quenching properties of included phenol groups, see: Dhar (1981). For related structures, see: Arai *et al.* (1994); Li *et al.* (1992); Patil *et al.* (2007); Shettigar *et al.* (2006). For standard bond lengths, see Allen *et al.* (1987). For density functional theory, see: Schmidt & Polik (2007); Hehre *et al.* (1986).



## Experimental

## Crystal data

$\text{C}_{16}\text{H}_{13}\text{BrO}_2$   $V = 651.68(7)$  Å<sup>3</sup>  
 $M_r = 317.17$   $Z = 2$   
 Monoclinic,  $P2_1$   $\text{Cu K}\alpha$  radiation  
 $a = 12.7300(8)$  Å  $\mu = 4.25$  mm<sup>-1</sup>  
 $b = 4.0061(3)$  Å  $T = 110$  K  
 $c = 13.0035(6)$  Å  $0.48 \times 0.41 \times 0.28$  mm  
 $\beta = 100.671(5)^\circ$

## Data collection

Oxford Diffraction Xcalibur diffractometer with Ruby (Gemini Cu) detector  
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2007)  
 $T_{\min} = 0.423$ ,  $T_{\max} = 1.000$   
 2117 measured reflections  
 1641 independent reflections  
 1621 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.018$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.091$   
 $S = 1.07$   
 1641 reflections  
 173 parameters  
 1 restraint  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.71$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.65$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983),  
 133 Friedel pairs  
 Flack parameter: 0.00 (3)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C12}-\text{H12A}\cdots\text{O1}^{\dagger}$	0.95	2.59	3.541 (5)	174

 Symmetry code: (i)  $-x + 1, y - \frac{1}{2}, -z$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2007); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

KV thanks the UGC for the sanction of a Junior Research Fellowship and for a SAP Chemical grant. HSY thanks the UOM for sabbatical leave. RJB acknowledges the NSF MRI program (grant No. CHE-0619278) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2728).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
 Arai, H., Higashigaki, Y., Goto, M. & Yano, S. (1994). *Jpn. J. Appl. Phys.* **33**, 5755–5758.  
 Dhar, D. N. (1981). *The Chemistry of Chalcones and Related Compounds*. New York: John Wiley.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Hehre, W. J., Randon, L., Schleyer, P. R. & Pople, J. A. (1986). *Ab Initio Molecular Orbital Theory*. New York: John Wiley.  
 Li, Z.-D., Huang, L.-R., Su, G.-B. & Wang, H.-J. (1992). *Jiegou Huaxue*, **11**, 1–4.  
 Oxford Diffraction (2007). *CrysAlis PRO* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.  
 Patil, P. S., Chantrapromma, S., Fun, H.-K., Dharmaparakash, S. M. & Babu, H. B. R. (2007). *Acta Cryst.* **E63**, o2612.  
 Schmidt, J. R. & Polik, W. F. (2007). *WebMO Pro*. WebMO, LLC, Holland, MI, USA, available from <http://www.webmo.net>.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Shettigar, V., Rosli, M. M., Fun, H.-K., Razak, I. A., Patil, P. S. & Dharmaparakash, S. M. (2006). *Acta Cryst.* **E62**, o4128–o4129.

## supporting information

*Acta Cryst.* (2010). E66, o1661 [doi:10.1107/S160053681002218X]

**(2E)-1-(2-Bromophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one**

Jerry P. Jasinski, Ray J. Butcher, K. Veena, B. Narayana and H. S. Yathirajan

**S1. Comment**

Chalcones, or 1,3-diaryl-2-propen-1-ones, belong to the flavonoid family. Chemically they consist of open-chain flavonoids in which the two aromatic rings are joined by a three-carbon  $\alpha,\beta$ -unsaturated carbonyl system. A vast number of naturally occurring chalcones are polyhydroxylated in the aryl rings. The radical quenching properties of the phenolic groups present in many chalcones have raised interest in using the compounds or chalcone rich plant extracts as drugs or food preservatives (Dhar, 1981). The crystal structures of some closely related chalcones, viz., 1-(4-bromophenyl)-3-(3-methoxy-phenyl)prop-2-en-1-one (Patil *et al.*, 2007); 4-bromo-4'-methoxy-chalcone (Li *et al.*, 1992), 4-bromo-4'-methoxychalcone (Arai *et al.*, 1994) and 1-(4-bromophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one (Shettigar *et al.*, 2006) have been reported. Hence in continuation with the synthesis and crystal structure determination and also owing to the importance of these flavonoid analogs, this new bromo chalcone, (I), C<sub>16</sub>H<sub>13</sub>BrO<sub>2</sub>, is synthesized and its crystal structure is reported.

The title compound, (I), C<sub>16</sub>H<sub>13</sub>BrO<sub>2</sub>, is a chalcone with 4-methoxyphenyl and 2-bromophenyl rings bonded at opposite rings of a propene group (Fig. 1). The dihedral angle between mean planes of the *para*-methoxy and *ortho*-bromo substituted benzene rings is 44.3 (9)°. The angles between the mean plane of the prop-2-ene-1-one group and the mean planes of the 4-methoxyphenyl and 2-bromophenyl rings are 6.3 (1)° and 44.6(36)°, respectively. Bond distances and angles are in normal ranges (Allen *et al.*, 1987). While no classical hydrogen bonds are present, a weak intermolecular C12—H12A...O1 interaction (Table 1) is observed which contributes to the stability of crystal packing.


A density functional theory (DFT) geometry optimization molecular orbital calculation (Schmidt & Polik, 2007) was performed on (I) with the B3LYP 6-31-G(*d*) basis set (Hehre *et al.*, 1986). The dihedral angle between mean planes of the *para*-methoxy and *ortho*-bromo substituted benzene rings becomes 45.98°, an increase of 1.59°. The angles between the mean plane of the prop-2-ene-1-one group and the mean planes of the 4-methoxyphenyl and 2-bromophenyl rings become 3.65° and 42.40°, changes of -2.65° and +0.74°, respectively. These observations suggest that the weak intermolecular C12—H12A...O1 interaction produces a small effect on crystal stability.

**S2. Experimental**

A 50% KOH solution was added to a mixture of 2-bromo acetophenone (0.01 mol, 1.99 g) and 4-methoxy benzaldehyde (0.01 mol, 1.36 g) in 25 ml of ethanol (Fig. 2). The mixture was stirred for an hour at room temperature and the precipitate was collected by filtration and purified by recrystallization from ethanol. The single-crystal was grown from ethyl acetate by slow evaporation method and yield of the compound was 50% (m.p.336–338 K). Analytical data, composition (%): found (calculated): C: 60.52 (60.59%); H: 4.10 (4.13%).


### S3. Refinement

The H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C–H distances = 0.95–0.98Å and with  $U_{\text{iso}}(\text{H}) = 1.17\text{--}1.47 U_{\text{eq}}(\text{C})$ .



**Figure 1**

Molecular structure of (I) showing the atom labeling scheme and 50% probability displacement ellipsoids.

**Figure 2**

Scheme for the synthesis of (I).

**(2E)-1-(2-Bromophenyl)-3-(4-methoxyphenyl)prop-2-en-1-one***Crystal data* $C_{16}H_{13}BrO_2$  $M_r = 317.17$ Monoclinic,  $P2_1$ Hall symbol:  $P\ 2_1yb$  $a = 12.7300\ (8)\ \text{\AA}$  $b = 4.0061\ (3)\ \text{\AA}$  $c = 13.0035\ (6)\ \text{\AA}$  $\beta = 100.671\ (5)^\circ$  $V = 651.68\ (7)\ \text{\AA}^3$  $Z = 2$  $F(000) = 320$  $D_x = 1.616\ \text{Mg m}^{-3}$ Cu  $K\alpha$  radiation,  $\lambda = 1.54184\ \text{\AA}$ 

Cell parameters from 1981 reflections

 $\theta = 4.5\text{--}74.1^\circ$  $\mu = 4.25\ \text{mm}^{-1}$  $T = 110\ \text{K}$ 

Chunk, colourless

 $0.48 \times 0.41 \times 0.28\ \text{mm}$

*Data collection*

Oxford Diffraction Xcalibur  
diffractometer with Ruby (Gemini Cu) detector  
Radiation source: Enhance (Cu) X-ray Source  
Graphite monochromator  
Detector resolution: 10.5081 pixels mm<sup>-1</sup>  
 $\omega$  scans  
Absorption correction: multi-scan  
(*CrysAlis RED*; Oxford Diffraction, 2007)  
 $T_{\min} = 0.423$ ,  $T_{\max} = 1.000$

2117 measured reflections  
1641 independent reflections  
1621 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.018$   
 $\theta_{\max} = 74.1^\circ$ ,  $\theta_{\min} = 4.5^\circ$   
 $h = -9 \rightarrow 15$   
 $k = -2 \rightarrow 4$   
 $l = -15 \rightarrow 16$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.091$   
 $S = 1.07$   
1641 reflections  
173 parameters  
1 restraint  
Primary atom site location: structure-invariant  
direct methods  
Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0718P)^2 + 0.5775P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.71 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.65 \text{ e } \text{\AA}^{-3}$   
Absolute structure: Flack (1983), 133 Friedel  
pairs  
Absolute structure parameter: 0.00 (3)

*Special details*

**Experimental.** IR data (KBr)  $\nu \text{ cm}^{-1}$ : 2998  $\text{cm}^{-1}$ , 2937  $\text{cm}^{-1}$ , 2839  $\text{cm}^{-1}$  (C—H al.str), 3058  $\text{cm}^{-1}$  (C—H ar. str) 1646  $\text{cm}^{-1}$  (C=O), 1580  $\text{cm}^{-1}$  (C=C); 1245  $\text{cm}^{-1}$  (C—O—C).

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.38764 (3)	0.18817 (18)	0.46303 (2)	0.01538 (15)
O1	0.2178 (2)	0.2427 (10)	0.1406 (2)	0.0211 (8)
O2	0.8860 (2)	0.2977 (9)	0.1659 (2)	0.0209 (7)
C1	0.2190 (3)	0.4727 (13)	0.3077 (3)	0.0149 (9)
C2	0.2577 (3)	0.4233 (12)	0.4142 (3)	0.0119 (8)
C3	0.2022 (3)	0.5312 (12)	0.4896 (3)	0.0166 (9)
H3A	0.2299	0.4917	0.5616	0.020*
C4	0.1060 (3)	0.6971 (19)	0.4599 (3)	0.0223 (8)
H4A	0.0679	0.7740	0.5116	0.027*
C5	0.0651 (3)	0.7513 (12)	0.3545 (3)	0.0211 (11)
H5A	-0.0008	0.8660	0.3342	0.025*
C6	0.1203 (3)	0.6385 (14)	0.2797 (3)	0.0187 (10)

H6A	0.0913	0.6732	0.2078	0.022*
C7	0.2730 (3)	0.3590 (12)	0.2195 (3)	0.0173 (9)
C8	0.3891 (3)	0.4058 (13)	0.2303 (3)	0.0160 (9)
H8A	0.4255	0.5526	0.2824	0.019*
C9	0.4441 (3)	0.2431 (12)	0.1673 (3)	0.0161 (10)
H9A	0.4037	0.1037	0.1154	0.019*
C10	0.5595 (3)	0.2564 (10)	0.1697 (3)	0.0145 (11)
C11	0.5999 (3)	0.1064 (11)	0.0878 (3)	0.0164 (10)
H11A	0.5519	-0.0064	0.0344	0.020*
C12	0.7083 (3)	0.1171 (11)	0.0822 (3)	0.0155 (10)
H12A	0.7334	0.0191	0.0248	0.019*
C13	0.7787 (3)	0.2733 (11)	0.1619 (3)	0.0156 (10)
C14	0.7411 (4)	0.4183 (13)	0.2470 (3)	0.0186 (9)
H14A	0.7897	0.5209	0.3022	0.022*
C15	0.6327 (3)	0.4102 (13)	0.2495 (3)	0.0175 (9)
H15A	0.6075	0.5107	0.3065	0.021*
C16	0.9272 (3)	0.1512 (18)	0.0810 (3)	0.0223 (11)
H16A	0.9121	-0.0888	0.0783	0.033*
H16B	1.0046	0.1870	0.0916	0.033*
H16C	0.8930	0.2553	0.0151	0.033*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br	0.0159 (2)	0.0153 (2)	0.0148 (2)	-0.0005 (2)	0.00224 (13)	0.00195 (19)
O1	0.0204 (13)	0.028 (2)	0.0151 (11)	-0.0044 (15)	0.0031 (10)	-0.0034 (14)
O2	0.0169 (14)	0.027 (2)	0.0207 (14)	-0.0012 (13)	0.0070 (12)	-0.0027 (13)
C1	0.0154 (18)	0.016 (2)	0.0154 (17)	-0.0033 (18)	0.0081 (15)	-0.0001 (17)
C2	0.0083 (16)	0.008 (2)	0.0179 (17)	0.0027 (16)	-0.0026 (13)	0.0032 (17)
C3	0.021 (2)	0.014 (2)	0.0160 (18)	-0.0060 (18)	0.0067 (15)	-0.0023 (16)
C4	0.0242 (18)	0.018 (2)	0.0292 (19)	-0.004 (3)	0.0173 (15)	-0.006 (3)
C5	0.0136 (17)	0.016 (3)	0.034 (2)	0.0014 (18)	0.0047 (16)	0.0002 (19)
C6	0.0165 (17)	0.016 (3)	0.0228 (17)	-0.003 (2)	0.0023 (14)	0.0024 (19)
C7	0.023 (2)	0.015 (2)	0.0152 (18)	-0.0028 (19)	0.0061 (16)	0.0040 (17)
C8	0.0176 (19)	0.015 (3)	0.0155 (17)	-0.0006 (18)	0.0036 (14)	0.0011 (17)
C9	0.0189 (17)	0.016 (3)	0.0131 (15)	-0.0011 (18)	0.0025 (13)	0.0027 (18)
C10	0.0197 (18)	0.014 (3)	0.0113 (15)	0.0016 (17)	0.0058 (14)	0.0018 (15)
C11	0.0208 (19)	0.015 (3)	0.0135 (16)	0.0001 (17)	0.0037 (14)	0.0000 (15)
C12	0.0200 (18)	0.015 (3)	0.0124 (15)	0.0016 (17)	0.0062 (14)	-0.0020 (15)
C13	0.0179 (18)	0.014 (3)	0.0159 (17)	0.0019 (16)	0.0068 (14)	0.0034 (16)
C14	0.024 (2)	0.019 (2)	0.0122 (16)	-0.001 (2)	0.0024 (15)	-0.0014 (18)
C15	0.022 (2)	0.020 (3)	0.0116 (16)	0.002 (2)	0.0071 (15)	0.0001 (18)
C16	0.0181 (17)	0.025 (3)	0.0261 (18)	0.000 (2)	0.0101 (14)	-0.004 (2)

*Geometric parameters (Å, °)*

Br—C2	1.907 (4)	C8—H8A	0.9500
O1—C7	1.224 (5)	C9—C10	1.464 (5)

O2—C13	1.361 (5)	C9—H9A	0.9500
O2—C16	1.433 (5)	C10—C11	1.401 (6)
C1—C2	1.396 (5)	C10—C15	1.403 (6)
C1—C6	1.408 (6)	C11—C12	1.396 (6)
C1—C7	1.512 (5)	C11—H11A	0.9500
C2—C3	1.380 (6)	C12—C13	1.387 (6)
C3—C4	1.384 (7)	C12—H12A	0.9500
C3—H3A	0.9500	C13—C14	1.410 (6)
C4—C5	1.390 (6)	C14—C15	1.387 (6)
C4—H4A	0.9500	C14—H14A	0.9500
C5—C6	1.379 (6)	C15—H15A	0.9500
C5—H5A	0.9500	C16—H16A	0.9800
C6—H6A	0.9500	C16—H16B	0.9800
C7—C8	1.471 (6)	C16—H16C	0.9800
C8—C9	1.341 (6)		
C13—O2—C16	116.7 (3)	C8—C9—H9A	116.4
C2—C1—C6	117.3 (4)	C10—C9—H9A	116.4
C2—C1—C7	125.7 (4)	C11—C10—C15	117.6 (4)
C6—C1—C7	117.1 (4)	C11—C10—C9	118.5 (4)
C3—C2—C1	121.9 (4)	C15—C10—C9	123.9 (4)
C3—C2—Br	116.4 (3)	C12—C11—C10	122.2 (4)
C1—C2—Br	121.7 (3)	C12—C11—H11A	118.9
C2—C3—C4	119.7 (4)	C10—C11—H11A	118.9
C2—C3—H3A	120.2	C13—C12—C11	118.9 (4)
C4—C3—H3A	120.2	C13—C12—H12A	120.6
C3—C4—C5	120.1 (4)	C11—C12—H12A	120.6
C3—C4—H4A	120.0	O2—C13—C12	124.5 (4)
C5—C4—H4A	120.0	O2—C13—C14	115.1 (4)
C6—C5—C4	119.9 (4)	C12—C13—C14	120.4 (4)
C6—C5—H5A	120.1	C15—C14—C13	119.5 (4)
C4—C5—H5A	120.1	C15—C14—H14A	120.2
C5—C6—C1	121.3 (4)	C13—C14—H14A	120.2
C5—C6—H6A	119.4	C14—C15—C10	121.3 (4)
C1—C6—H6A	119.4	C14—C15—H15A	119.3
O1—C7—C8	122.6 (4)	C10—C15—H15A	119.3
O1—C7—C1	118.7 (4)	O2—C16—H16A	109.5
C8—C7—C1	118.6 (4)	O2—C16—H16B	109.5
C9—C8—C7	120.5 (4)	H16A—C16—H16B	109.5
C9—C8—H8A	119.7	O2—C16—H16C	109.5
C7—C8—H8A	119.7	H16A—C16—H16C	109.5
C8—C9—C10	127.2 (4)	H16B—C16—H16C	109.5
C6—C1—C2—C3	0.1 (7)	C1—C7—C8—C9	164.4 (4)
C7—C1—C2—C3	-179.0 (4)	C7—C8—C9—C10	-178.4 (4)
C6—C1—C2—Br	177.7 (4)	C8—C9—C10—C11	-170.9 (4)
C7—C1—C2—Br	-1.3 (7)	C8—C9—C10—C15	9.4 (7)
C1—C2—C3—C4	-0.8 (7)	C15—C10—C11—C12	-2.4 (6)

Br—C2—C3—C4	-178.6 (4)	C9—C10—C11—C12	177.9 (4)
C2—C3—C4—C5	0.7 (8)	C10—C11—C12—C13	1.9 (6)
C3—C4—C5—C6	0.2 (9)	C16—O2—C13—C12	0.2 (7)
C4—C5—C6—C1	-1.1 (8)	C16—O2—C13—C14	179.6 (5)
C2—C1—C6—C5	0.9 (7)	C11—C12—C13—O2	179.4 (4)
C7—C1—C6—C5	180.0 (4)	C11—C12—C13—C14	0.0 (6)
C2—C1—C7—O1	139.4 (5)	O2—C13—C14—C15	179.2 (4)
C6—C1—C7—O1	-39.6 (7)	C12—C13—C14—C15	-1.4 (7)
C2—C1—C7—C8	-43.2 (7)	C13—C14—C15—C10	0.8 (7)
C6—C1—C7—C8	137.8 (5)	C11—C10—C15—C14	1.0 (7)
O1—C7—C8—C9	-18.2 (7)	C9—C10—C15—C14	-179.3 (4)

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C12—H12A...O1 <sup>i</sup>	0.95	2.59	3.541 (5)	174

Symmetry code: (i)  $-x+1, y-1/2, -z$ .