

**Di-*tert*-butyl 2,2'-[*(biphenyl-4,4'*-diyl)-dioxy]diacetate**

**Qamar Ali,<sup>a</sup> Sammer Yousuf,<sup>a</sup> Muhammad Raza Shah<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>**

<sup>a</sup>H.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and

<sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 15 June 2010; accepted 16 June 2010

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.120; data-to-parameter ratio = 18.9.

The complete molecule of the title compound,  $C_{24}H_{30}O_6$ , is generated by a crystallographic inversion centre. In the unique part of the molecule, the four-atom  $-\text{O}-\text{CH}_2-\text{C}(=\text{O})-\text{O}-$  chain between the benzene ring and the *tert*-butyl group assumes a zigzag conformation [ $\text{O}-\text{C}-\text{C}-\text{O}$  torsion angle =  $-162.3(1)^\circ$ ].

**Related literature**

For a related structure, see: Shah *et al.* (2010).

**Experimental***Crystal data*

$C_{24}H_{30}O_6$   
 $M_r = 414.48$   
Monoclinic,  $P2_1/c$   
 $a = 9.9390(7)\text{ \AA}$   
 $b = 12.6247(8)\text{ \AA}$   
 $c = 9.8458(7)\text{ \AA}$   
 $\beta = 114.645(1)^\circ$

$V = 1122.88(13)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.09\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.4 \times 0.3 \times 0.2\text{ mm}$

*Data collection*

Bruker SMART APEX  
diffractometer  
7450 measured reflections

2572 independent reflections  
2029 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.024$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.120$   
 $S = 1.02$   
2572 reflections

136 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.16\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.14\text{ e \AA}^{-3}$

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the Higher Education Commission of Pakistan and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5072).

**References**

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2002). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Shah, K., Yousuf, S., Raza Shah, M. & Ng, S. W. (2010). *Acta Cryst. E* **66**, o1705.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *publCIF*. In preparation.

# supporting information

*Acta Cryst.* (2010). E66, o1739 [doi:10.1107/S1600536810023391]

## Di-*tert*-butyl 2,2'-(biphenyl-4,4'-diyl)dioxy]diacetate

**Qamar Ali, Sammer Yousuf, Muhammad Raza Shah and Seik Weng Ng**

### S1. Comment


We are interested in the solid-state structures of *V*-shaped molecules; recently we reported the crystal structure of 9,9-bis-[4-(*tert*-butoxycarbonylmethoxy)phenyl]fluorene (Shah *et al.*, 2010). For such a shape, the number of carbon atoms making up the kink must be an odd number. In the present compound, the two aromatic rings are directly connected; the molecule lies on a center of symmetry (Fig. 1). The four-atom  $\text{O}-\text{CH}_2-\text{C}(=\text{O})-\text{O}-$  chain between the aromatic ring and the *tert*-butyl group assumes a zigzag conformation [ $\text{O}-\text{C}-\text{C}-\text{O}$  torsion angle 162.3 (1)  $^{\circ}$ ].

### S2. Experimental

4,4'-Dihydroxybiphenyl (1 g, 2.4 mmol) was dissolved in acetone (25 ml). To the solution was added potassium carbonate (0.67 g, 4.8 mmol) and *t*-butyl bromoacetate (0.75 ml, 4.8 mmol). The mixture was stirred at room temperature for 3 h. The solvent was evaporated under reduced pressure and the residue was dissolved in a mixture of water (50 ml) and dichloromethane (50 ml). The aqueous layer was extracted three times with dichloromethane. The combined organic phases were evaporated under reduced pressure and the solid material was recrystallized from *n*-hexane.

### S3. Refinement

H-atoms were placed in calculated positions [C–H 0.93–0.97  $\text{\AA}$ ,  $U(\text{H})$  1.2–1.5  $U(\text{C})$ ] and were included in the refinement in the riding model approximation.



**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{24}\text{H}_{30}\text{O}_6$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Symmetry code: (i) = 1 -  $x$ , 1 -  $y$ , 1 -  $z$ .

**Di-*tert*-butyl 2,2'-[**(biphenyl-4,4'-diyl)dioxy**]diacetate***Crystal data*

$C_{24}H_{36}O_6$   
 $M_r = 414.48$   
Monoclinic,  $P2_1/c$   
Hall symbol: -P 2ybc  
 $a = 9.9390 (7)$  Å  
 $b = 12.6247 (8)$  Å  
 $c = 9.8458 (7)$  Å  
 $\beta = 114.645 (1)^\circ$   
 $V = 1122.88 (13)$  Å<sup>3</sup>  
 $Z = 2$

$F(000) = 444$   
 $D_x = 1.226 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 2683 reflections  
 $\theta = 2.8\text{--}28.4^\circ$   
 $\mu = 0.09 \text{ mm}^{-1}$   
 $T = 293$  K  
Block, colorless  
 $0.4 \times 0.3 \times 0.2$  mm

*Data collection*

Bruker SMART APEX  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
7450 measured reflections  
2572 independent reflections

2029 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.024$   
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.3^\circ$   
 $h = -9 \rightarrow 12$   
 $k = -15 \rightarrow 16$   
 $l = -12 \rightarrow 12$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.120$   
 $S = 1.02$   
2572 reflections  
136 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0597P)^2 + 0.1713P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.16 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.14 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.16964 (10)	1.03985 (7)	0.42919 (10)	0.0453 (2)
O2	0.24672 (13)	0.91302 (9)	0.31464 (12)	0.0643 (3)
O3	0.49943 (10)	0.88751 (7)	0.57432 (12)	0.0538 (3)
C1	0.03883 (16)	1.07454 (11)	0.29402 (15)	0.0507 (4)
C2	0.0877 (2)	1.11440 (17)	0.1761 (2)	0.0784 (5)
H2A	0.1256	1.0563	0.1397	0.118*
H2B	0.1636	1.1669	0.2193	0.118*
H2C	0.0047	1.1452	0.0949	0.118*
C3	-0.0216 (2)	1.16473 (14)	0.3541 (2)	0.0726 (5)
H3A	-0.0548	1.1376	0.4260	0.109*
H3B	-0.1032	1.1973	0.2733	0.109*
H3C	0.0548	1.2163	0.4010	0.109*
C4	-0.07095 (18)	0.98421 (15)	0.2398 (2)	0.0708 (5)
H4A	-0.0985	0.9629	0.3184	0.106*

H4B	-0.0266	0.9255	0.2119	0.106*
H4C	-0.1574	1.0070	0.1549	0.106*
C5	0.25888 (15)	0.96225 (10)	0.42366 (15)	0.0434 (3)
C6	0.37853 (15)	0.94408 (10)	0.57882 (16)	0.0474 (3)
H6A	0.4129	1.0118	0.6274	0.057*
H6B	0.3377	0.9046	0.6374	0.057*
C7	0.48857 (14)	0.77893 (10)	0.55146 (14)	0.0430 (3)
C8	0.40501 (16)	0.71184 (11)	0.59620 (17)	0.0501 (3)
H8	0.3454	0.7391	0.6398	0.060*
C9	0.41070 (15)	0.60337 (10)	0.57559 (15)	0.0465 (3)
H9	0.3543	0.5589	0.6066	0.056*
C10	0.49740 (13)	0.55858 (9)	0.51049 (13)	0.0372 (3)
C11	0.58025 (16)	0.62869 (11)	0.46692 (16)	0.0485 (3)
H11	0.6399	0.6020	0.4230	0.058*
C12	0.57629 (16)	0.73697 (11)	0.48722 (17)	0.0513 (4)
H12	0.6332	0.7817	0.4573	0.062*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0429 (5)	0.0464 (5)	0.0447 (5)	0.0080 (4)	0.0165 (4)	-0.0013 (4)
O2	0.0713 (7)	0.0657 (7)	0.0547 (6)	0.0123 (5)	0.0250 (5)	-0.0130 (5)
O3	0.0412 (5)	0.0355 (5)	0.0814 (7)	0.0021 (4)	0.0224 (5)	-0.0019 (4)
C1	0.0454 (7)	0.0560 (8)	0.0483 (7)	0.0088 (6)	0.0171 (6)	0.0094 (6)
C2	0.0807 (12)	0.0946 (14)	0.0670 (11)	0.0121 (10)	0.0378 (10)	0.0271 (9)
C3	0.0677 (11)	0.0677 (11)	0.0821 (11)	0.0265 (9)	0.0309 (9)	0.0127 (9)
C4	0.0509 (9)	0.0787 (11)	0.0664 (10)	-0.0030 (8)	0.0082 (8)	0.0053 (8)
C5	0.0447 (7)	0.0381 (6)	0.0503 (7)	-0.0006 (5)	0.0227 (6)	-0.0037 (5)
C6	0.0470 (7)	0.0381 (6)	0.0538 (8)	0.0051 (5)	0.0177 (6)	-0.0026 (5)
C7	0.0361 (6)	0.0364 (6)	0.0503 (7)	0.0024 (5)	0.0117 (5)	0.0010 (5)
C8	0.0493 (8)	0.0448 (7)	0.0637 (8)	-0.0005 (6)	0.0312 (7)	-0.0061 (6)
C9	0.0489 (7)	0.0417 (7)	0.0549 (8)	-0.0054 (6)	0.0277 (6)	-0.0014 (5)
C10	0.0341 (6)	0.0379 (6)	0.0359 (6)	0.0010 (5)	0.0109 (5)	0.0033 (5)
C11	0.0487 (7)	0.0415 (7)	0.0645 (8)	0.0046 (6)	0.0327 (7)	0.0058 (6)
C12	0.0485 (8)	0.0399 (7)	0.0715 (9)	0.0004 (6)	0.0310 (7)	0.0085 (6)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C5	1.3379 (15)	C4—H4C	0.9600
O1—C1	1.4867 (16)	C5—C6	1.5118 (19)
O2—C5	1.2020 (16)	C6—H6A	0.9700
O3—C7	1.3861 (15)	C6—H6B	0.9700
O3—C6	1.4143 (16)	C7—C12	1.3787 (19)
C1—C4	1.513 (2)	C7—C8	1.3807 (19)
C1—C2	1.519 (2)	C8—C9	1.3890 (19)
C1—C3	1.518 (2)	C8—H8	0.9300
C2—H2A	0.9600	C9—C10	1.3905 (18)
C2—H2B	0.9600	C9—H9	0.9300

C2—H2C	0.9600	C10—C11	1.3929 (18)
C3—H3A	0.9600	C10—C10 <sup>i</sup>	1.497 (2)
C3—H3B	0.9600	C11—C12	1.3845 (19)
C3—H3C	0.9600	C11—H11	0.9300
C4—H4A	0.9600	C12—H12	0.9300
C4—H4B	0.9600		
C5—O1—C1	121.76 (10)	O2—C5—C6	124.59 (12)
C7—O3—C6	119.66 (10)	O1—C5—C6	108.99 (10)
O1—C1—C4	109.05 (11)	O3—C6—C5	111.44 (11)
O1—C1—C2	110.05 (12)	O3—C6—H6A	109.3
C4—C1—C2	113.18 (14)	C5—C6—H6A	109.3
O1—C1—C3	102.28 (12)	O3—C6—H6B	109.3
C4—C1—C3	110.99 (14)	C5—C6—H6B	109.3
C2—C1—C3	110.73 (14)	H6A—C6—H6B	108.0
C1—C2—H2A	109.5	C12—C7—C8	119.36 (12)
C1—C2—H2B	109.5	C12—C7—O3	115.67 (12)
H2A—C2—H2B	109.5	C8—C7—O3	124.85 (12)
C1—C2—H2C	109.5	C7—C8—C9	119.45 (13)
H2A—C2—H2C	109.5	C7—C8—H8	120.3
H2B—C2—H2C	109.5	C9—C8—H8	120.3
C1—C3—H3A	109.5	C8—C9—C10	122.62 (12)
C1—C3—H3B	109.5	C8—C9—H9	118.7
H3A—C3—H3B	109.5	C10—C9—H9	118.7
C1—C3—H3C	109.5	C9—C10—C11	116.31 (11)
H3A—C3—H3C	109.5	C9—C10—C10 <sup>i</sup>	121.94 (14)
H3B—C3—H3C	109.5	C11—C10—C10 <sup>i</sup>	121.75 (14)
C1—C4—H4A	109.5	C12—C11—C10	121.79 (13)
C1—C4—H4B	109.5	C12—C11—H11	119.1
H4A—C4—H4B	109.5	C10—C11—H11	119.1
C1—C4—H4C	109.5	C7—C12—C11	120.47 (13)
H4A—C4—H4C	109.5	C7—C12—H12	119.8
H4B—C4—H4C	109.5	C11—C12—H12	119.8
O2—C5—O1	126.41 (13)		
C5—O1—C1—C4	64.56 (16)	C12—C7—C8—C9	-0.1 (2)
C5—O1—C1—C2	-60.14 (17)	O3—C7—C8—C9	-175.79 (13)
C5—O1—C1—C3	-177.86 (12)	C7—C8—C9—C10	-0.3 (2)
C1—O1—C5—O2	0.4 (2)	C8—C9—C10—C11	0.3 (2)
C1—O1—C5—C6	-178.70 (11)	C8—C9—C10—C10 <sup>i</sup>	179.82 (14)
C7—O3—C6—C5	-77.86 (15)	C9—C10—C11—C12	-0.1 (2)
O2—C5—C6—O3	18.60 (19)	C10 <sup>i</sup> —C10—C11—C12	-179.55 (14)
O1—C5—C6—O3	-162.30 (10)	C8—C7—C12—C11	0.3 (2)
C6—O3—C7—C12	153.70 (12)	O3—C7—C12—C11	176.43 (12)
C6—O3—C7—C8	-30.43 (19)	C10—C11—C12—C7	-0.3 (2)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .