

Bis(1-benzylpiperazine-1,4-dinium) hexachloridocadmate(II) dihydrate

Meher El Glaoui,^a Matthias Zeller,^b Erwann Jeanneau^c and Cherif Ben Nasr^{a*}

^aLaboratoire de Chimie des Matériaux, Faculté des Sciences de Bizerte, 7021 Zarzouna, Tunisia, ^bYoungstown State University, Department of Chemistry, One University Plaza, Youngstown, Ohio 44553-3663, USA, and ^cUniversité Lyon1, Centre de Diffractométrie Henri Longchambon, 43 Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France
Correspondence e-mail: cherif_bennasr@yahoo.fr


Received 30 June 2010; accepted 1 July 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.002 \text{ \AA}$; R factor = 0.022; wR factor = 0.057; data-to-parameter ratio = 25.8.

The asymmetric unit of the title compound, $(C_{11}H_{18}N_2)_2[CdCl_6] \cdot 2H_2O$, consists of one 1-benzylpiperazine-1,4-dinium cation, one water molecule and one-half of a $[CdCl_6]^{4-}$ anion, located on an inversion centre. The crystal packing is governed by an extensive three-dimensional network of intermolecular O—H···Cl, C—H···Cl, N—H···O and N—H···Cl hydrogen bonds, two of them bifurcated.

Related literature

For *meta*-chlorido complexes, see: El Glaoui, Jeanneau, *et al.* (2009); El Glaoui, Kefi *et al.* (2009). For the role of C—H···Cl hydrogen bonds, see: Janiak & Scharmann (2003). For a discussion of Cd—Cl distances and Cl—Cd—Cl bond angles, see: Bala *et al.* (2006).



Experimental

Crystal data

$(C_{11}H_{18}N_2)_2[CdCl_6] \cdot 2H_2O$

$M_r = 717.68$

Monoclinic, $P2_1/c$

$a = 12.734 (2) \text{ \AA}$

$b = 9.1686 (14) \text{ \AA}$

$c = 13.216 (2) \text{ \AA}$

$\beta = 103.249 (3)^\circ$

$V = 1502.0 (4) \text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 1.29 \text{ mm}^{-1}$

$T = 100 \text{ K}$

$0.55 \times 0.45 \times 0.25 \text{ mm}$

Data collection

Bruker SMART APEX CCD

diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2009)

$T_{\min} = 0.622$, $T_{\max} = 0.746$

11244 measured reflections

4446 independent reflections

4123 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.022$

$wR(F^2) = 0.057$

$S = 1.07$

4446 reflections

172 parameters

3 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.83 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N2—H2A···Cl1 ⁱ	0.92	2.58	3.3383 (11)	140
N2—H2A···Cl2 ⁱ	0.92	2.59	3.2672 (11)	131
N2—H2B···Cl2 ⁱⁱ	0.92	2.47	3.1846 (11)	135
N2—H2B···Cl3 ⁱⁱ	0.92	2.58	3.2799 (12)	133
N1—H1···O1	0.89 (1)	1.92 (1)	2.7945 (16)	170 (2)
O1—H1A···Cl1	0.84 (2)	2.39 (2)	3.1678 (11)	155 (2)
O1—H1B···Cl3 ⁱⁱⁱ	0.83 (2)	2.42 (2)	3.2152 (11)	161 (2)
C9—H9A···Cl3 ⁱⁱ	0.99	2.83	3.331 (2)	112
C9—H9B···Cl3 ⁱⁱⁱ	0.99	2.85	3.659 (3)	139
C10—H10A···Cl3 ⁱⁱⁱ	0.99	2.73	3.565 (2)	143
C10—H10B···Cl2 ⁱⁱ	0.99	2.84	3.340 (4)	112
C11—H11A···Cl1 ⁱⁱ	0.99	2.71	3.626 (1)	154
C11—H11B···Cl1	0.99	2.72	3.587 (1)	146

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $x, y + 1, z$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXTL*.

We would like to acknowledge the support provided by the Secretary of State for Scientific Research and Technology of Tunisia. The diffractometer was funded by NSF grant 0087210, by Ohio Board of Regents grant CAP-491, and by YSU.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5541).

References

- Bala, R., Sharma, R. P., Sharma, U. & Ferretti, V. (2006). *Acta Cryst. C* **62**, m628–m631.
- Brandenburg, K. (1998). *DIAMOND*. Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison Wisconsin, USA.
- El Glaoui, M., Jeanneau, E., Lefebvre, F. & Ben Nasr, C. (2009). *Can. J. Anal. Sci. Spectr.* **54**, 70–81.
- El Glaoui, M., Kefi, R., Jeanneau, E., Lefebvre, F. & Ben Nasr, C. (2009). *Can. J. Anal. Sci. Spectr.* **54**, 281–291.
- Janiak, C. & Scharmann, T. G. (2003). *Polyhedron*, **22**, 1123–1133.
- Sheldrick, G. M. (2008). *Acta Cryst. A*, **64**, 112–122.

supporting information

Acta Cryst. (2010). E66, m895 [https://doi.org/10.1107/S1600536810026073]

Bis(1-benzylpiperazine-1,4-dium) hexachloridocadmate(II) dihydrate

Meher El Glaoui, Matthias Zeller, Erwann Jeanneau and Cherif Ben Nasr

S1. Comment

As a part of our ongoing investigations in molecular salts containing *meta*-chlorido complexes (El Glaoui, Jeanneau, *et al.*, 2009; El Glaoui, Kefi *et al.*, 2009), we present here the crystal structure of one such compound, $(C_{11}H_{18}N_2)_2CdCl_6 \cdot 2H_2O$, (Fig. 1). The asymmetric unit of its structure consists of one 1-benzylpiperazine-1,4-dium dication doubly protonated at the N1 and N2 nitrogen atoms, one water molecule and one-half of a $CdCl_6^{4-}$ anion (located on a crystallographic inversion centre) (Fig. 1). The atomic arrangement of $(C_{11}H_{18}N_2)_2CdCl_6 \cdot 2H_2O$ can be described as built up by inorganic chains of $CdCl_6$ octahedra and water molecules extending along the *b* direction held together by O—H \cdots Cl hydrogen bonds (Fig. 2, Table 1). Two such chains cross the unit cell at $z = 0$, $z = 1/2$ and $x = 1/2$ (Fig. 3). The organic groups are located between these chains and connect to them through N—H \cdots Cl, C—H \cdots Cl and N—H \cdots O hydrogen bonds to form a three dimensional infinite network (Fig. 3, Table 1). All the chloride ions are involved in hydrogen bonding. It should be pointed out at this point that the C—H \cdots Cl hydrogen bonds do usually not play a large role in stabilizing a structure (Janiak & Scharmann, 2003), but due to the large number of these interactions in the title compound they seem to substantially contribute to the choice of packing observed in the structure of the title compound. Among all the hydrogen bonds, two are bifurcated: N2—H2A \cdots (Cl1, Cl2) and N2—H2B \cdots (Cl2, Cl3). The H1 hydrogen atom attached to the N1 nitrogen atom is bonded only to the water molecule, *via* the N1—H1 \cdots O1 hydrogen bond, and not to the $CdCl_6^{4-}$ anion.


The Cd II ion is in an octahedral coordination environment composed of six chloride anions as to form a hexachlorocadmate (II) ion. In this kind of anion, the Cd—Cl bond lengths and Cl—Cd—Cl bond angles are generally not equal to one another but vary with the environment around the Cl atoms (Bala *et al.*, 2006). In the title compound, the values of the Cd—Cl bond lengths vary between 2.5528 (5) and 2.7055 (4) Å. The Cl—Cd—Cl angles range from 87.354 (11) to 92.646 (11)°. These geometrical parameters agree with those found in $[Co(NH_3)_6]_4 [CdCl_6] [CdCl_4(SCN)(H_2O)]_2Cl_2 \cdot 2H_2O$ where the Cd—Cl distances are between 2.5937 (9) and 2.691 (1) Å and the Cl—Cd—Cl angles ranging from 89.23 (3) to 95.50 (3)° (Bala *et al.*, 2006). Owing to the obvious differences of Cd—Cl distances and Cl—Cd—Cl angles in $(C_{11}H_{18}N_2)_2CdCl_6 \cdot 2H_2O$, the coordination geometry of the Cd atom could be regarded as a slightly distorted octahedron which is in full agreement with the literature data (Bala, *et al.*, 2006).

S2. Experimental

1-Benzylpiperazine (2 mmol, 0.352 g) and $CdCl_2$ (1 mmol, 0.183 g), were dissolved in dilute HCl (10 ml, 1 M) and the resultant solution was slowly evaporated at room temperature. A crystal of the title compound, which remained stable under normal conditions of temperature and humidity, was isolated after several days and subjected to X-ray diffraction analysis (yield 55%).

S3. Refinement

C—H and NH_2^+ hydrogen atoms were placed in calculated positions with C—H in the range 0.93–0.97 and N—H equal to 0.92 Å. The N—H⁺ and the water hydrogen atom positions were refined with N—H and O—H distance restraints of 0.91 (2) and 0.84 (2) Å. The $U_{\text{iso}}(\text{H})$ values of all H atoms were constrained to 1.2 or 1.5 times U_{eq} of the respective parent atom.

**Figure 1**

A view of the title compound, showing 50% probability displacement ellipsoids, arbitrary spheres for the H atoms, and the atom numbering scheme.





Figure 2


Projection along the a axis of the inorganic chains in $(\text{C}_{11}\text{H}_{18}\text{N}_2)_2\text{CdCl}_6 \cdot 2\text{H}_2\text{O}$. Hydrogen bonds are denoted by dotted lines.

**Figure 3**

The packing of $(C_{11}H_{18}N_2)_2[CdCl_6] \cdot 2H_2O$, viewed down the b axis. Hydrogen bonds are denoted by dotted lines.

Bis(1-benzylpiperazine-1,4-dium) hexachloridocadmate(II) dihydrate

Crystal data



$M_r = 717.68$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.734(2)$ Å

$b = 9.1686(14)$ Å

$c = 13.216(2)$ Å

$\beta = 103.249(3)^\circ$

$V = 1502.0(4)$ Å³

$Z = 2$

$F(000) = 732$

$D_x = 1.587$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2788 reflections

$\theta = 2.7\text{--}31.0^\circ$

$\mu = 1.29$ mm⁻¹

$T = 100$ K

Plate, colourless

$0.55 \times 0.45 \times 0.25$ mm

Data collection

Bruker SMART APEX CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2009)

$T_{\min} = 0.622$, $T_{\max} = 0.746$

11244 measured reflections

4446 independent reflections

4123 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 31.3^\circ$, $\theta_{\min} = 1.6^\circ$

$h = -17 \rightarrow 18$

$k = -12 \rightarrow 13$

$l = -18 \rightarrow 19$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.022$$

$$wR(F^2) = 0.057$$

$$S = 1.07$$

4446 reflections

172 parameters

3 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0269P)^2 + 0.6306P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.51 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.83 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$
C1	0.03492 (12)	0.81561 (17)	0.11848 (12)	0.0242 (3)
H1C	0.0759	0.7794	0.0722	0.029*
C2	-0.07139 (13)	0.85867 (19)	0.08024 (14)	0.0312 (3)
H2	-0.1032	0.8503	0.0081	0.037*
C3	-0.13120 (12)	0.91374 (16)	0.14687 (15)	0.0292 (3)
H3	-0.2032	0.9457	0.1202	0.035*
C4	-0.08597 (12)	0.92209 (16)	0.25191 (14)	0.0269 (3)
H4	-0.1272	0.9588	0.2978	0.032*
C5	0.01972 (11)	0.87702 (15)	0.29110 (12)	0.0213 (3)
H5	0.0500	0.8814	0.3637	0.026*
C6	0.08142 (10)	0.82535 (13)	0.22417 (11)	0.0164 (2)
C7	0.19878 (10)	0.78999 (13)	0.26364 (11)	0.0156 (2)
H7A	0.2267	0.7431	0.2076	0.019*
H7B	0.2074	0.7203	0.3222	0.019*
C8	0.23911 (10)	1.04936 (12)	0.22283 (10)	0.0125 (2)
H8A	0.1609	1.0712	0.2067	0.015*
H8B	0.2591	1.0205	0.1575	0.015*
C9	0.30187 (10)	1.18282 (12)	0.26720 (10)	0.0122 (2)
H9A	0.2858	1.2635	0.2162	0.015*
H9B	0.2795	1.2138	0.3309	0.015*
C10	0.44650 (10)	1.02741 (13)	0.36618 (10)	0.0120 (2)
H10A	0.4307	1.0540	0.4337	0.014*
H10B	0.5244	1.0051	0.3782	0.014*
C11	0.38168 (9)	0.89435 (12)	0.32287 (10)	0.0116 (2)

H11A	0.4022	0.8627	0.2584	0.014*
H11B	0.3981	0.8137	0.3738	0.014*
Cd1	0.5000	0.5000	0.5000	0.01071 (4)
Cl1	0.36321 (2)	0.69672 (3)	0.55240 (2)	0.01362 (6)
Cl2	0.59831 (3)	0.49308 (3)	0.69137 (2)	0.01174 (6)
Cl3	0.37573 (2)	0.28368 (3)	0.54229 (2)	0.01263 (6)
N1	0.26320 (8)	0.92695 (11)	0.29971 (8)	0.01118 (18)
N2	0.41977 (8)	1.15258 (10)	0.29270 (8)	0.01172 (19)
H2A	0.4565	1.2342	0.3221	0.014*
H2B	0.4416	1.1317	0.2326	0.014*
O1	0.23289 (9)	0.99157 (10)	0.49769 (9)	0.0180 (2)
H1	0.2483 (14)	0.95558 (19)	0.3591 (12)	0.018 (4)*
H1A	0.2690 (17)	0.924 (2)	0.5315 (17)	0.044 (6)*
H1B	0.2655 (18)	1.066 (2)	0.5229 (18)	0.047 (7)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0181 (7)	0.0284 (7)	0.0253 (7)	-0.0036 (5)	0.0031 (6)	-0.0025 (6)
C2	0.0203 (7)	0.0358 (8)	0.0323 (9)	-0.0051 (6)	-0.0049 (6)	0.0040 (7)
C3	0.0131 (6)	0.0210 (6)	0.0504 (10)	-0.0028 (5)	0.0008 (6)	0.0050 (6)
C4	0.0169 (7)	0.0201 (6)	0.0458 (10)	-0.0028 (5)	0.0117 (6)	-0.0048 (6)
C5	0.0166 (6)	0.0208 (6)	0.0274 (7)	-0.0038 (5)	0.0068 (5)	-0.0034 (5)
C6	0.0113 (6)	0.0134 (5)	0.0241 (7)	-0.0034 (4)	0.0033 (5)	-0.0013 (5)
C7	0.0132 (6)	0.0106 (5)	0.0223 (6)	-0.0020 (4)	0.0027 (5)	-0.0012 (4)
C8	0.0126 (5)	0.0112 (5)	0.0129 (6)	0.0007 (4)	0.0013 (4)	0.0025 (4)
C9	0.0122 (5)	0.0106 (5)	0.0144 (5)	0.0007 (4)	0.0042 (4)	0.0005 (4)
C10	0.0114 (5)	0.0129 (5)	0.0111 (6)	0.0004 (4)	0.0014 (4)	0.0004 (4)
C11	0.0096 (5)	0.0116 (5)	0.0136 (5)	0.0017 (4)	0.0028 (4)	0.0007 (4)
Cd1	0.01320 (7)	0.00972 (6)	0.00955 (7)	-0.00012 (4)	0.00332 (5)	0.00002 (4)
Cl1	0.01548 (14)	0.01289 (12)	0.01327 (13)	0.00202 (10)	0.00489 (11)	0.00058 (10)
Cl2	0.01277 (14)	0.01112 (12)	0.01138 (14)	-0.00071 (8)	0.00289 (11)	0.00010 (9)
Cl3	0.01428 (13)	0.01192 (12)	0.01209 (13)	-0.00170 (9)	0.00382 (10)	-0.00078 (9)
N1	0.0112 (5)	0.0100 (4)	0.0127 (5)	-0.0004 (3)	0.0032 (4)	0.0003 (4)
N2	0.0119 (5)	0.0110 (4)	0.0131 (5)	-0.0008 (3)	0.0046 (4)	-0.0005 (4)
O1	0.0185 (5)	0.0165 (4)	0.0190 (5)	0.0001 (3)	0.0046 (4)	-0.0001 (3)

Geometric parameters (\AA , ^\circ)

C1—C2	1.389 (2)	C9—H9A	0.9900
C1—C6	1.390 (2)	C9—H9B	0.9900
C1—H1C	0.9500	C10—N2	1.4915 (16)
C2—C3	1.385 (3)	C10—C11	1.5102 (17)
C2—H2	0.9500	C10—H10A	0.9900
C3—C4	1.378 (3)	C10—H10B	0.9900
C3—H3	0.9500	C11—N1	1.4992 (15)
C4—C5	1.390 (2)	C11—H11A	0.9900
C4—H4	0.9500	C11—H11B	0.9900

C5—C6	1.3938 (19)	Cd1—Cl2 ⁱ	2.5528 (5)
C5—H5	0.9500	Cd1—Cl2	2.5528 (5)
C6—C7	1.5012 (18)	Cd1—Cl3	2.6751 (4)
C7—N1	1.5158 (15)	Cd1—Cl3 ⁱ	2.6751 (4)
C7—H7A	0.9900	Cd1—Cl1 ⁱ	2.7055 (4)
C7—H7B	0.9900	Cd1—Cl1	2.7055 (4)
C8—N1	1.4978 (15)	N1—H1	0.889 (14)
C8—C9	1.5051 (16)	N2—H2A	0.9200
C8—H8A	0.9900	N2—H2B	0.9200
C8—H8B	0.9900	O1—H1A	0.837 (16)
C9—N2	1.4875 (15)	O1—H1B	0.830 (16)
C2—C1—C6	120.17 (15)	C11—C10—H10A	109.5
C2—C1—H1C	119.9	N2—C10—H10B	109.5
C6—C1—H1C	119.9	C11—C10—H10B	109.5
C3—C2—C1	120.28 (16)	H10A—C10—H10B	108.1
C3—C2—H2	119.9	N1—C11—C10	110.70 (9)
C1—C2—H2	119.9	N1—C11—H11A	109.5
C4—C3—C2	119.81 (14)	C10—C11—H11A	109.5
C4—C3—H3	120.1	N1—C11—H11B	109.5
C2—C3—H3	120.1	C10—C11—H11B	109.5
C3—C4—C5	120.30 (15)	H11A—C11—H11B	108.1
C3—C4—H4	119.8	Cl2 ⁱ —Cd1—Cl2	180.0
C5—C4—H4	119.8	Cl2 ⁱ —Cd1—Cl3	92.646 (11)
C4—C5—C6	120.22 (14)	Cl2—Cd1—Cl3	87.354 (11)
C4—C5—H5	119.9	Cl2 ⁱ —Cd1—Cl3 ⁱ	87.354 (11)
C6—C5—H5	119.9	Cl2—Cd1—Cl3 ⁱ	92.646 (11)
C1—C6—C5	119.18 (13)	Cl3—Cd1—Cl3 ⁱ	180.0
C1—C6—C7	119.76 (12)	Cl2 ⁱ —Cd1—Cl1 ⁱ	87.784 (12)
C5—C6—C7	120.93 (13)	Cl2—Cd1—Cl1 ⁱ	92.218 (12)
C6—C7—N1	110.75 (10)	Cl3—Cd1—Cl1 ⁱ	90.322 (15)
C6—C7—H7A	109.5	Cl3 ⁱ —Cd1—Cl1 ⁱ	89.680 (15)
N1—C7—H7A	109.5	Cl2 ⁱ —Cd1—Cl1	92.215 (12)
C6—C7—H7B	109.5	Cl2—Cd1—Cl1	87.783 (11)
N1—C7—H7B	109.5	Cl3—Cd1—Cl1	89.678 (14)
H7A—C7—H7B	108.1	Cl3 ⁱ —Cd1—Cl1	90.321 (14)
N1—C8—C9	109.69 (10)	Cl1 ⁱ —Cd1—Cl1	180.0
N1—C8—H8A	109.7	C8—N1—C11	109.14 (9)
C9—C8—H8A	109.7	C8—N1—C7	113.30 (10)
N1—C8—H8B	109.7	C11—N1—C7	110.23 (9)
C9—C8—H8B	109.7	C8—N1—H1	108.9 (11)
H8A—C8—H8B	108.2	C11—N1—H1	106.7 (11)
N2—C9—C8	110.79 (9)	C7—N1—H1	108.3 (12)
N2—C9—H9A	109.5	C9—N2—C10	111.02 (9)
C8—C9—H9A	109.5	C9—N2—H2A	109.4
N2—C9—H9B	109.5	C10—N2—H2A	109.4
C8—C9—H9B	109.5	C9—N2—H2B	109.4
H9A—C9—H9B	108.1	C10—N2—H2B	109.4

N2—C10—C11	110.58 (10)	H2A—N2—H2B	108.0
N2—C10—H10A	109.5	H1A—O1—H1B	103 (3)
C6—C1—C2—C3	1.0 (2)	N1—C8—C9—N2	59.09 (13)
C1—C2—C3—C4	-1.8 (2)	N2—C10—C11—N1	-56.82 (13)
C2—C3—C4—C5	0.7 (2)	C9—C8—N1—C11	-59.78 (12)
C3—C4—C5—C6	1.1 (2)	C9—C8—N1—C7	176.99 (10)
C2—C1—C6—C5	0.8 (2)	C10—C11—N1—C8	59.02 (13)
C2—C1—C6—C7	-175.11 (13)	C10—C11—N1—C7	-175.95 (10)
C4—C5—C6—C1	-1.8 (2)	C6—C7—N1—C8	-48.38 (14)
C4—C5—C6—C7	173.99 (12)	C6—C7—N1—C11	-171.01 (10)
C1—C6—C7—N1	109.33 (14)	C8—C9—N2—C10	-56.81 (13)
C5—C6—C7—N1	-66.48 (15)	C11—C10—N2—C9	55.34 (12)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N2—H2A \cdots Cl1 ⁱⁱ	0.92	2.58	3.3383 (11)	140
N2—H2A \cdots Cl2 ⁱⁱ	0.92	2.59	3.2672 (11)	131
N2—H2B \cdots Cl2 ⁱⁱⁱ	0.92	2.47	3.1846 (11)	135
N2—H2B \cdots Cl3 ⁱⁱⁱ	0.92	2.58	3.2799 (12)	133
N1—H1 \cdots O1	0.89 (1)	1.92 (1)	2.7945 (16)	170 (2)
O1—H1A \cdots Cl1	0.84 (2)	2.39 (2)	3.1678 (11)	155 (2)
O1—H1B \cdots Cl3 ^{iv}	0.83 (2)	2.42 (2)	3.2152 (11)	161 (2)
C9—H9A \cdots Cl3 ⁱⁱⁱ	0.99	2.83	3.331 (2)	112
C9—H9B \cdots Cl3 ^{iv}	0.99	2.85	3.659 (3)	139
C10—H10A \cdots Cl3 ^{iv}	0.99	2.73	3.565 (2)	143
C10—H10B \cdots Cl2 ⁱⁱⁱ	0.99	2.84	3.340 (4)	112
C11—H11A \cdots Cl1 ⁱⁱⁱ	0.99	2.71	3.626 (1)	154
C11—H11B \cdots Cl1	0.99	2.72	3.587 (1)	146

Symmetry codes: (ii) $-x+1, -y+2, -z+1$; (iii) $x, -y+3/2, z-1/2$; (iv) $x, y+1, z$.