

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dicyanidobis(*N,N'*-dimethylthiourea- κ S)-mercury(II)Muhammad Riaz Malik,^a Saqib Ali,^a Saeed Ahmad,^{b*} Muhammad Altaf^c and Helen Stoeckli-Evans^c^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad, Pakistan,^bDepartment of Chemistry, University of Engineering and Technology, Lahore 54890, Pakistan, and ^cInstitute of Physics, University of Neuchâtel, rue Emile-Argand 11, CH-2009 Neuchâtel, Switzerland

Correspondence e-mail: saeed_a786@hotmail.com


Received 29 July 2010; accepted 30 July 2010

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{N}-\text{C}) = 0.008$ Å; R factor = 0.017; wR factor = 0.036; data-to-parameter ratio = 16.5.

In the title complex, $[\text{Hg}(\text{CN})_2(\text{C}_3\text{H}_8\text{N}_2\text{S})_2]$, the Hg^{II} atom is located on a twofold rotation axis. It is four-coordinate having an irregular tetrahedral geometry composed of two cyanide C atoms [$\text{Hg}-\text{C} = 2.090$ (6) Å] and two thione S atoms of *N,N'*-dimethylthiourea (dmu) [$\text{Hg}-\text{S} = 2.7114$ (9) Å]. The $\text{NC}-\text{Hg}-\text{CN}$ bond angle of 148.83 (13°) has the greatest deviation from the ideal tetrahedral geometry. The molecular structure is stabilized by intramolecular $\text{N}-\text{H}\cdots\text{S}$ interactions involving dmu units related by the twofold symmetry. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{N}(\text{CN})$ hydrogen-bonding interactions link symmetry-related molecules into a two-dimensional network in (110).

Related literature

For the biological applications of mercury(II) complexes of thiones, see: Akrivos (2001); Bell *et al.* (2001); Popovic *et al.* (2000). For background to mercury(II) complexes of thiourea and its derivatives, see: Ahmad *et al.* (2009); Jiang *et al.* (2001); Lobana *et al.* (2008); Mufakkar *et al.* (2010); Nawaz *et al.* (2010); Popovic *et al.* (2000); Wu *et al.* (2004). For the crystal structures of cyanide complexes of d^{10} metals, see: Ahmad *et al.* (2009); Altaf *et al.* (2010); Fettouhi *et al.* (2010); Hanif *et al.* (2007).



Experimental

Crystal data

$[\text{Hg}(\text{CN})_2(\text{C}_3\text{H}_8\text{N}_2\text{S})_2]$
 $M_r = 460.98$
 Monoclinic, $C2/c$
 $a = 18.1161$ (11) Å
 $b = 7.7533$ (5) Å
 $c = 14.0553$ (8) Å
 $\beta = 128.533$ (3°)

$V = 1544.32$ (16) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 10.23$ mm⁻¹
 $T = 173$ K
 $0.40 \times 0.31 \times 0.25$ mm

Data collection

Stoe IPDS 2 diffractometer
 Absorption correction: multi-scan
 (*MULscanABS* embedded in
PLATON; Spek, 2009)
 $T_{\text{min}} = 0.270$, $T_{\text{max}} = 1.000$

8116 measured reflections
 1451 independent reflections
 1411 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.036$
 $S = 1.14$
 1451 reflections
 88 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.68$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.97$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{S1}^i$	0.80 (6)	2.67 (5)	3.415 (4)	157 (4)
$\text{N2}-\text{H2N}\cdots\text{N3}^{ii}$	0.79 (5)	2.21 (6)	2.951 (7)	155 (4)

Symmetry codes: (i) $-x + 1, y, -z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *X-Area* (Stoe & Cie, 2009); cell refinement: *X-Area*; data reduction: *X-RED32* (Stoe & Cie, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and *PLATON*.

We thank the staff of the X-ray Application Lab, CSEM, Neuchâtel, for access to the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2389).

References

- Ahmad, S., Sadaf, H., Akkurt, M., Sharif, S. & Khan, I. U. (2009). *Acta Cryst.* **E65**, m1191–m1192.
 Akrivos, P. D. (2001). *Coord. Chem. Rev.* **213**, 181–210.
 Altaf, M., Stoeckli-Evans, H., Ahmad, S., Isab, A. A., Al-Arfaj, A. R., Malik, M. R. & S. Ali, (2010). *J. Chem. Crystallogr.* **40**. In the press.
 Bell, N. A., Branston, T. N., Clegg, W., Parker, L., Raper, E. S., Sammon, C. & Constable, C. P. (2001). *Inorg. Chim. Acta*, **319**, 130–136.
 Fettouhi, M., Riaz Malik, M., Ali, S. A., Isab, A. & Ahmad, S. (2010). *Acta Cryst.* **E66**, m997.
 Hanif, M., Ahmad, S., Altaf, M. & Stoeckli-Evans, H. (2007). *Acta Cryst.* **E63**, m2594.
 Jiang, X. N., Xu, D., Yuan, D. R. & Yu, W. T. (2001). *Chin. Chem. Lett.* **12**, 279–282.
 Lobana, T. S., Sharma, R., Sharma, R., Sultana, R. & Butcher, R. J. (2008). *Z. Anorg. Allg. Chem.* **634**, 718–723.
 Mufakkar, M., Tahir, M. N., Sadaf, H., Ahmad, S. & Waheed, A. (2010). *Acta Cryst.* **E66**, m1001–m1002.

- Nawaz, S., Sadaf, H., Fettouhi, M., Fazal, A. & Ahmad, S. (2010). *Acta Cryst.* **E66**, m952.
- Popovic, Z., Pavlovic, G., Matkovic-Calogovic, D., Soldin, Z., M. Rajic, M. Vikić-Topić, D., Kovacek, D. (2000). *Inorg. Chim. Acta*, **306**, 142–152.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stoe & Cie (2009). *X-AREA* and *X-RED32*. Stoe & Cie GmbH, Darmstadt, Germany.
- Wu, Z.-Y., Xu, D.-J. & Hung, C.-H. (2004). *J. Coord. Chem.* **57**, 791–796.

supporting information

Acta Cryst. (2010). E66, m1060–m1061 [https://doi.org/10.1107/S1600536810030424]

Dicyanidobis(*N,N'*-dimethylthiourea- κ S)mercury(II)

Muhammad Riaz Malik, Saqib Ali, Saeed Ahmad, Muhammad Altaf and Helen Stoeckli-Evans

S1. Comment

The structural characterization of mercury(II) complexes of thioamides is an important aspect of inorganic chemistry because such complexes can be used as models for metal-sulfur interactions in biological systems (Akrivos, 2001; Bell *et al.*, 2001; Popovic *et al.*, 2000). Several crystallographic reports about mercury(II) complexes of the type, L_2HgX_2 (L = thiourea or its derivatives) reveal that these complexes usually consist of discrete monomeric molecules with tetrahedral (somewhat distorted) coordination environments around mercury(II) (Ahmad *et al.*, 2009; Bell *et al.*, 2001; Jiang *et al.*, 2001; Lobana *et al.*, 2008; Mufakkar *et al.*, 2010; Nawaz *et al.*, 2010; Popovic *et al.*, 2000; Wu *et al.*, 2004). Recently, we have reported the crystal structures of a number of cyanido complexes of d^{10} metal ions with L -type ligands, including the crystal structure of a trinuclear complex, [$\{(tmtu)_2Hg(CN)_2\}_2Hg(CN)_2$] ($tmtu$ = tetramethylthiourea), which presents a unique example of a $Hg(CN)_2$ bridged mercury(II)-thione complex (Ahmad *et al.*, 2009; Altaf *et al.*, 2010; Fettouhi *et al.*, 2010; Hanif *et al.*, 2007). Herein, we report on the crystal structure of the title mercury cyanide complex of *N,N'*-dimethylthiourea, [$Hg(dmtu)_2(CN)_2$].

The title monomeric complex is composed of an $Hg(CN)_2$ unit with two *N,N'*-dimethylthiourea ($dmtu$) ligands coordinated to the Hg atom *via* the S atom (Fig. 1). The four-coordinate mercury atom is located on a two-fold rotation axis and adopts a severely distorted tetrahedral geometry, the bond angles being in the range of 94.31 (3) - 148.83 (13)°. The molecular structure is stabilized by intramolecular N-H...S interactions involving $dmtu$ units related by the two-fold symmetry (Fig. 1, Table 1). The bond distances and bond angles are in agreement with those reported for related compounds (Ahmad *et al.*, 2009; Altaf *et al.*, 2010; Jiang *et al.*, 2001; Lobana *et al.*, 2008; Mufakkar *et al.*, 2010; Nawaz *et al.*, 2010; Popovic *et al.*, 2000; Wu *et al.*, 2004). The SCN_2 moiety of $dmtu$ is planar [to within 0.002 (1) Å] with the C—N and C—S bond lengths corresponding to the values intermediate between single and double bonds. The Hg-C≡N unit is nearly linear with a bond angle of 175.3 (3)°. The compound is closely related with [$Hg(N,N'$ -dibutylthiourea) $_2(CN)_2$] (Ahmad *et al.*, 2009).

In the crystal packing of the title complex, symmetry-related molecules are connected *via* intermolecular N—H...N hydrogen bonds, involving the thiourea NH atoms and the N atom of the CN^- anions (Fig. 2, Table 1). This gives rise to the formation of a two-dimensional network in (110). This is the same arrangement as observed previously for the dibutylthiourea compound mentioned above.

S2. Experimental

To 0.25 g (1.0 mmol) mercury(II) cyanide in 10 ml methanol was added 2 equivalents of *N,N'*-dimethylthiourea in methanol. On mixing, a clear solution was obtained. It was then stirred for 30 minutes after which it was filtered and the filtrate kept at RT for crystallization by slow evaporation of the solvent. As a result, colourless block-like crystals, suitable for X-ray diffraction analysis, were obtained.

S3. Refinement

The NH H-atoms were located in difference electron-density maps and were freely refined: N—H = 0.80 (6) & 0.79 (5) Å. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.98 Å, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{parent C-atom})$.




Figure 1

The molecular structure of the title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. The intramolecular N—H...S interactions are shown as double dashed lines (see Table 1 for details). [Symmetry code (a) $-x+1, y, -z+1/2$.]




Figure 2

A crystal packing diagram of the title complex showing the N—H...S and N—H...N hydrogen bonding interactions (dashed lines; see Table 1 for details).

Dicyanidobis(*N,N'*-dimethylthiourea- κ S)mercury(II)*Crystal data*[Hg(CN)₂(C₃H₈N₂S)₂] $M_r = 460.98$ Monoclinic, *C2/c*Hall symbol: $-C\ 2yc$ $a = 18.1161(11)$ Å $b = 7.7533(5)$ Å $c = 14.0553(8)$ Å $\beta = 128.533(3)^\circ$ $V = 1544.32(16)$ Å³ $Z = 4$ $F(000) = 872$ $D_x = 1.983$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 12331 reflections
 $\theta = 1.9\text{--}26.1^\circ$
 $\mu = 10.23 \text{ mm}^{-1}$

$T = 173 \text{ K}$
 Block, colourless
 $0.40 \times 0.31 \times 0.25 \text{ mm}$

Data collection

Stoe IPDS 2
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ - + ω - scans
 Absorption correction: multi-scan
 (MULscanABS embedded in PLATON; Spek,
 2009)
 $T_{\min} = 0.270$, $T_{\max} = 1.000$

8116 measured reflections
 1451 independent reflections
 1411 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$
 $\theta_{\max} = 25.6^\circ$, $\theta_{\min} = 2.9^\circ$
 $h = -21 \rightarrow 21$
 $k = -9 \rightarrow 9$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.036$
 $S = 1.14$
 1451 reflections
 88 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0129P)^2 + 2.6833P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.68 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.97 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.50000	-0.21587 (2)	0.25000	0.0231 (1)
S1	0.37952 (6)	0.02195 (11)	0.08082 (7)	0.0255 (3)
N1	0.3941 (2)	0.1650 (4)	0.2640 (3)	0.0277 (9)
N2	0.2450 (2)	0.0856 (4)	0.1004 (2)	0.0241 (8)
N3	0.3776 (3)	-0.3173 (4)	0.3370 (3)	0.0441 (12)
C1	0.3355 (2)	0.0948 (4)	0.1535 (3)	0.0210 (9)
C2	0.3657 (3)	0.2174 (5)	0.3367 (3)	0.0386 (13)
C3	0.1748 (2)	0.0052 (5)	-0.0168 (3)	0.0306 (11)
C4	0.4229 (3)	-0.2883 (4)	0.3084 (3)	0.0295 (10)
H1N	0.449 (3)	0.161 (5)	0.296 (3)	0.024 (10)*
H2A	0.31970	0.31150	0.29540	0.0580*
H2B	0.42120	0.25690	0.41680	0.0580*

H2C	0.33710	0.11910	0.34680	0.0580*
H2N	0.227 (3)	0.128 (5)	0.134 (3)	0.030 (10)*
H3A	0.17870	0.05510	-0.07770	0.0460*
H3B	0.11180	0.02530	-0.04110	0.0460*
H3C	0.18680	-0.11920	-0.01060	0.0460*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.0209 (1)	0.0274 (1)	0.0294 (1)	0.0000	0.0197 (1)	0.0000
S1	0.0249 (5)	0.0344 (4)	0.0242 (4)	0.0067 (3)	0.0188 (4)	0.0020 (3)
N1	0.0244 (18)	0.0359 (17)	0.0275 (13)	0.0008 (14)	0.0185 (14)	-0.0045 (12)
N2	0.0225 (16)	0.0279 (15)	0.0273 (13)	0.0015 (11)	0.0181 (12)	-0.0035 (11)
N3	0.040 (2)	0.058 (2)	0.054 (2)	-0.0070 (17)	0.0389 (19)	0.0028 (16)
C1	0.0236 (18)	0.0192 (15)	0.0255 (14)	0.0042 (12)	0.0179 (14)	0.0031 (11)
C2	0.041 (3)	0.050 (2)	0.0352 (18)	-0.0019 (18)	0.0288 (19)	-0.0128 (16)
C3	0.022 (2)	0.0341 (19)	0.0336 (16)	-0.0025 (15)	0.0163 (16)	-0.0047 (14)
C4	0.028 (2)	0.0288 (17)	0.0344 (16)	-0.0018 (15)	0.0207 (16)	0.0009 (14)

Geometric parameters (Å, °)

Hg1—S1	2.7114 (9)	N3—C4	1.139 (8)
Hg1—C4	2.090 (6)	N1—H1N	0.80 (6)
Hg1—S1 ⁱ	2.7114 (9)	N2—H2N	0.79 (5)
Hg1—C4 ⁱ	2.090 (6)	C2—H2A	0.9800
S1—C1	1.736 (4)	C2—H2B	0.9800
N1—C1	1.335 (5)	C2—H2C	0.9800
N1—C2	1.459 (7)	C3—H3A	0.9800
N2—C1	1.314 (6)	C3—H3B	0.9800
N2—C3	1.452 (4)	C3—H3C	0.9800
S1—Hg1—C4	99.05 (11)	S1—C1—N1	119.6 (3)
S1—Hg1—S1 ⁱ	94.31 (3)	Hg1—C4—N3	175.3 (3)
S1—Hg1—C4 ⁱ	102.01 (9)	N1—C2—H2A	109.00
S1 ⁱ —Hg1—C4	102.01 (9)	N1—C2—H2B	110.00
C4—Hg1—C4 ⁱ	148.83 (13)	N1—C2—H2C	109.00
S1 ⁱ —Hg1—C4 ⁱ	99.05 (11)	H2A—C2—H2B	109.00
Hg1—S1—C1	96.84 (11)	H2A—C2—H2C	109.00
C1—N1—C2	123.8 (4)	H2B—C2—H2C	109.00
C1—N2—C3	124.7 (3)	N2—C3—H3A	109.00
C1—N1—H1N	117 (3)	N2—C3—H3B	110.00
C2—N1—H1N	118 (3)	N2—C3—H3C	109.00
C3—N2—H2N	117 (3)	H3A—C3—H3B	110.00
C1—N2—H2N	118 (3)	H3A—C3—H3C	109.00
S1—C1—N2	121.1 (3)	H3B—C3—H3C	109.00
N1—C1—N2	119.3 (4)		
C4—Hg1—S1—C1	32.52 (15)	C2—N1—C1—S1	-174.9 (3)

S1 ⁱ —Hg1—S1—C1	-70.39 (13)	C2—N1—C1—N2	6.6 (5)
C4 ⁱ —Hg1—S1—C1	-170.60 (16)	C3—N2—C1—S1	4.6 (5)
Hg1—S1—C1—N1	60.6 (3)	C3—N2—C1—N1	-177.0 (3)
Hg1—S1—C1—N2	-121.0 (3)		

Symmetry code: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N1—H1N···S1 ⁱ	0.80 (6)	2.67 (5)	3.415 (4)	157 (4)
N2—H2N···N3 ⁱⁱ	0.79 (5)	2.21 (6)	2.951 (7)	155 (4)

Symmetry codes: (i) $-x+1, y, -z+1/2$; (ii) $-x+1/2, y+1/2, -z+1/2$.