

$[\mu\text{-}N,N,N',N'\text{-Tetrakis(2-pyridylmethyl)-butane-1,4-diamine}]_{\text{bis}}[\text{dinitrato-cadmium(II)}]$

 Mark Bartholomä,[‡] Hoi Cheung and Jon Zubieta*

Department of Chemistry, Syracuse University, Syracuse, New York 13244, USA

Correspondence e-mail: jazubiet@syr.edu


Received 29 July 2010; accepted 26 August 2010

 Key indicators: single-crystal X-ray study; $T = 90\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.030; wR factor = 0.069; data-to-parameter ratio = 17.2.

The title dinuclear cadmium complex, $[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{28}\text{H}_{32}\text{N}_6)]$, is located on an inversion center. The unique Cd^{II} ion displays a 5 + 2 coordination. A distorted square-pyramidal geometry is formed by the dipicolylamine group of the ligand *via* the N atoms in a meridional fashion and two O atoms of the nitrate ligands with short Cd–O distances. The coordination is completed by two loosely bound O atoms of the nitrate ligands.

Related literature

For crystallographic data of tetrakis(pyridin-2-yl-methyl)-alkyl-diamine compounds, see: Fujihara *et al.* (2004); Mambanda *et al.* (2007). For the superoxide dismutase activity of iron complexes, see: Tamura *et al.* (2000). For dinuclear Pt complexes of similar ligands, see: Ertürk *et al.* (2007). For the use of the dipicolylamine moiety for binding of the $M(\text{CO})_3$ core ($M = \text{Re}, {}^{99\text{m}}\text{Tc}$), see: Bartholomä *et al.* (2009). For crystal structures closely related to the title compound, see: Bartholomä *et al.* (2010a,b,c,d)



[‡] Current address: Harvard Medical School, Children's Hospital Boston, Department of Radiology, 300 Longwood Ave, Enders 4, Boston, MA 02115, USA.

Experimental

Crystal data

$[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{28}\text{H}_{32}\text{N}_6)]$
 $M_r = 925.44$
 Triclinic, $P\bar{1}$
 $a = 8.0548(8)\text{ \AA}$
 $b = 8.7010(8)\text{ \AA}$
 $c = 13.2566(13)\text{ \AA}$
 $\alpha = 107.488(2)^\circ$
 $\beta = 96.767(2)^\circ$

$\gamma = 104.631(2)^\circ$
 $V = 838.30(14)\text{ \AA}^3$
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 1.35\text{ mm}^{-1}$
 $T = 90\text{ K}$
 $0.48 \times 0.30 \times 0.08\text{ mm}$

Data collection

Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 1998)
 $T_{\text{min}} = 0.564$, $T_{\text{max}} = 0.900$

8382 measured reflections
 4043 independent reflections
 3912 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.069$
 $S = 1.15$
 4043 reflections

235 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.01\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.50\text{ e \AA}^{-3}$

Table 1

Selected bond lengths (\AA).

Cd1–N2	2.250 (2)	Cd1–O2	2.588 (2)
Cd1–N3	2.251 (2)	Cd1–O4	2.687 (2)
Cd1–O5	2.279 (2)	Cd1–N1	2.427 (2)
Cd1–O1	2.322 (2)		

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 1999); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

This work was supported by funding from Syracuse University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5105).

References

- Bartholomä, M., Cheung, H., Darling, K. & Zubieta, J. (2010d). *Acta Cryst.* E66, m1201–m1202.
- Bartholomä, M., Cheung, H. & Zubieta, J. (2010a). *Acta Cryst.* E66, m1195–m1196.
- Bartholomä, M., Cheung, H. & Zubieta, J. (2010b). *Acta Cryst.* E66, m1198.
- Bartholomä, M., Cheung, H. & Zubieta, J. (2010c). *Acta Cryst.* E66, m1199–m1200.
- Bartholomä, M., Valliant, J., Maresca, K. P., Babich, J. & Zubieta, J. (2009). *Chem. Commun.* 5, 473–604.
- Brandenburg, K. & Putz, H. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (1998). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ertürk, H., Hofmann, A., Puchta, R. & van Eldik, R. (2007). *Dalton Trans.* pp. 2295–2301.
- Fujihara, T., Saito, M. & Nagasawa, A. (2004). *Acta Cryst.* E60, o1126–o1128.
- Mambanda, A., Jaganyi, D. & Munro, O. Q. (2007). *Acta Cryst.* C63, o676–o680.
- Sheldrick, G. M. (2008). *Acta Cryst.* A64, 112–122.
- Tamura, M., Urano, Y., Kikuchi, K., Higuchi, T., Hirobe, M. & Nagano, T. (2000). *J. Organomet. Chem.* 611, 586–592.

supporting information

Acta Cryst. (2010). E66, m1197 [doi:10.1107/S1600536810034549]

[μ -*N,N,N',N'*-Tetrakis(2-pyridylmethyl)butane-1,4-diamine]bis-[dinitratocadmium(II)]

Mark Bartholomä, Hoi Cheung and Jon Zubieta

S1. Comment

The described ligand *N^l,N^l,N^l,N^l*-tetrakis(pyridin-2-ylmethyl)butane-1,4-diamine has been used as starting material in the hydrothermal synthesis of metal-organic transition metal/molybdateoxide frameworks in the principal author's laboratory (Bartholomä, unpublished results). The title complex was prepared as part of a series with different cadmium and copper salts to study the coordination properties of the ligand with these metals without the interaction of metaloxide clusters (Bartholomä, 2010*b,c,d*).

Another crystalline species of a monomeric dinuclear cadmium complex using the corresponding acetate salt as metal source has been reported by our group (Bartholomä, 2010*a*). In the cadmium acetate structure, the Cd—N distances of the pyridine N atoms are slightly longer [2.313 (3) Å and 2.379 (3) Å] whereas the distance of the tertiary nitrogen atom is with marginally shorter [2.405 (3)] Å when compared to the distances of the title complex.

The dipicolylamine moiety has originally been developed in our laboratory as metal chelating entity for binding of the *M*(CO)₃ core (*M* = Re, ^{99m}Tc) for radiopharmaceutical purposes. However, a different coordination mode has been observed for the *M*(CO)₃ core in which the dipicolylamine metal chelate is coordinated in a facial manner (Bartholomä, 2009).

Crystal structures of the ligands *N^l,N^l,N³,N³*-tetrakis(2-pyridiniomethyl)-1,3-diaminopropane and *N^l,N^l,N^l,N^l*-tetrakis(pyridin-2-ylmethyl)butane-1,4-diamine have been described recently (Fujihara, 2004; Mambanda, 2007). Superoxide dismutase activity of iron(II) complexes of *N^l,N^l,N³,N³*-tetrakis(2-pyridiniomethyl)-1,3-diaminopropane and related ligands has been investigated by Tamura *et al.* (2000). Studies on the thermodynamic and kinetic behaviour of the reaction of platinum(II) complexes of higher ligand homologues with chloride have been performed by Ertürk *et al.* (2007).

S2. Experimental

***N^l,N^l,N^l,N^l*-tetrakis(pyridin-2-ylmethyl)butane-1,4-diamine.** An amount of 1.00 g (11.34 mmol) 1,4-diaminobutane was dissolved in 30 ml anhydrous dichloroethane under an inert atmosphere (argon) followed by the addition of 4.55 ml (47.65 mmol) pyridine-2-carboxaldehyde. The mixture was stirred for 15 min at r.t. and then cooled with an ice bath prior to the portionwise addition of 14.43 g (68.06 mmol) sodium triacetoxyborohydride (gas evolution, exothermic reaction). The reaction was stirred overnight allowing the temperature slowly to rise to room temperature. The reaction was quenched by the dropwise addition of saturated sodium bicarbonate solution and stirring was continued until the gas evolution ceased. The mixture was separated and the organic layer was further washed with saturated sodium bicarbonate solution, water and brine. The organic phase was dried with anhydrous sodium sulfate, filtered and the solvent removed under reduced pressure. The crude reaction mixture was then purified by silica gel column chromatography starting with chloroform and increasing gradient to chloroform:methanol 10:1 (v/v). Yield: 4.02 g (78%). ¹H NMR (CDCl₃): δ = 8.40

(m, 4H), 7.51 (m, 4H), 7.39 (d, $J = 7.81$ Hz, 4H), 7.02 (m, 4H), 3.67 (s, 8H), 2.39 (m, 4H), 1.42 (m, 4H) p.p.m..

Synthesis of metal complex. To 2 ml of an aqueous solution of cadmium nitrate, two equivalents (50 mg, 0.11 mmol) of N^1,N^1,N^4,N^4 -tetrakis(pyridin-2-ylmethyl)butane-1,4-diamine in 2 ml methanol were added followed by the addition of 2 ml N,N -dimethylformamide. Single crystals were obtained after a week by slow evaporation of the solvents at room temperature.

S3. Refinement

All the H atoms were placed in idealized positions and refined using the riding model approximation with $C-H_{\text{aryl}} = 0.95$ Å, $C-H_{\text{methyl}} = 0.98$ Å and $C-H_{\text{methylene}} = 0.99$ Å and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ and $1.2U_{\text{eq}}(\text{C}_{\text{methylene/aryl}})$.




Figure 1

The crystal structure of the title complex. The displacement ellipsoids are drawn at 50% probability level. Hydrogen atoms are omitted for clarity. Unlabeled atoms are related by the symmetry code $(-x, -y + 2, -z + 1)$.

$[\mu\text{-}N,N,N/i>'<,N'$ -Tetrakis(2-pyridylmethyl)butane- 1,4-diamine]bis[dinitratocadmium(II)]

Crystal data

$[\text{Cd}_2(\text{NO}_3)_4(\text{C}_{28}\text{H}_{32}\text{N}_6)]$

$M_r = 925.44$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 8.0548$ (8) Å

$b = 8.7010$ (8) Å

$c = 13.2566$ (13) Å

$\alpha = 107.488$ (2)°

$\beta = 96.767$ (2)°

$\gamma = 104.631$ (2)°

$V = 838.30$ (14) Å³

$Z = 1$

$F(000) = 462$
 $D_x = 1.833 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 5752 reflections
 $\theta = 2.5\text{--}28.4^\circ$

$\mu = 1.35 \text{ mm}^{-1}$
 $T = 90 \text{ K}$
 Plate, colourless
 $0.48 \times 0.30 \times 0.08 \text{ mm}$

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 512 pixels mm^{-1}
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 1998)
 $T_{\min} = 0.564$, $T_{\max} = 0.900$

8382 measured reflections
 4043 independent reflections
 3912 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 28.1^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -10 \rightarrow 10$
 $k = -11 \rightarrow 11$
 $l = -16 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.069$
 $S = 1.15$
 4043 reflections
 235 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0286P)^2 + 0.8284P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.01 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.50 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.37059 (2)	0.85996 (2)	0.212593 (14)	0.01675 (6)
O1	0.4785 (3)	0.7640 (3)	0.05839 (16)	0.0335 (5)
O2	0.2104 (3)	0.6092 (3)	0.03615 (16)	0.0270 (4)
O3	0.3515 (3)	0.5592 (3)	-0.09358 (19)	0.0511 (7)
O4	0.5248 (3)	1.1197 (3)	0.39807 (18)	0.0347 (5)
O5	0.6551 (3)	1.0276 (3)	0.27055 (17)	0.0347 (5)
O6	0.7985 (3)	1.2505 (3)	0.41010 (18)	0.0356 (5)
N1	0.0990 (3)	0.8167 (2)	0.27756 (16)	0.0158 (4)
N2	0.2297 (3)	1.0238 (3)	0.15892 (16)	0.0186 (4)
N3	0.3643 (3)	0.6590 (3)	0.28768 (16)	0.0189 (4)
N4	0.3458 (3)	0.6419 (3)	-0.00139 (19)	0.0269 (5)

N5	0.6610 (3)	1.1361 (3)	0.36162 (18)	0.0221 (4)
C1	0.0530 (3)	0.6391 (3)	0.2732 (2)	0.0207 (5)
H1A	-0.0030	0.5639	0.1979	0.025*
H1B	-0.0335	0.6213	0.3190	0.025*
C2	0.2108 (3)	0.5907 (3)	0.31117 (19)	0.0194 (5)
C3	0.1947 (4)	0.4684 (3)	0.3606 (2)	0.0245 (5)
H3A	0.0869	0.4226	0.3787	0.029*
C4	0.3379 (4)	0.4149 (3)	0.3827 (2)	0.0276 (6)
H4A	0.3289	0.3310	0.4158	0.033*
C5	0.4951 (4)	0.4839 (3)	0.3566 (2)	0.0273 (6)
H5	0.5944	0.4475	0.3706	0.033*
C6	0.5034 (4)	0.6068 (3)	0.3099 (2)	0.0238 (5)
H6A	0.6109	0.6562	0.2928	0.029*
C7	-0.0241 (3)	0.8446 (3)	0.19778 (19)	0.0192 (5)
H7A	-0.1265	0.8629	0.2287	0.023*
H7B	-0.0672	0.7419	0.1322	0.023*
C8	0.0606 (3)	0.9957 (3)	0.16622 (18)	0.0175 (5)
C9	-0.0369 (3)	1.0924 (3)	0.13757 (19)	0.0203 (5)
H9	-0.1557	1.0729	0.1446	0.024*
C10	0.0413 (4)	1.2182 (3)	0.0984 (2)	0.0229 (5)
H10	-0.0244	1.2841	0.0772	0.028*
C11	0.2153 (4)	1.2464 (3)	0.0906 (2)	0.0225 (5)
H11	0.2712	1.3318	0.0641	0.027*
C12	0.3059 (3)	1.1477 (3)	0.1222 (2)	0.0208 (5)
H12	0.4260	1.1675	0.1180	0.025*
C13	0.1306 (3)	0.9396 (3)	0.38864 (18)	0.0167 (4)
H13A	0.1702	1.0550	0.3854	0.020*
H13B	0.2283	0.9253	0.4341	0.020*
C14	-0.0244 (3)	0.9281 (3)	0.44558 (19)	0.0194 (5)
H14A	-0.1265	0.9347	0.3994	0.023*
H14B	-0.0579	0.8180	0.4571	0.023*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.01544 (9)	0.01864 (10)	0.01966 (10)	0.00647 (6)	0.00744 (6)	0.00900 (7)
O1	0.0278 (10)	0.0318 (11)	0.0282 (10)	-0.0031 (8)	0.0117 (8)	0.0001 (8)
O2	0.0228 (9)	0.0307 (10)	0.0304 (10)	0.0077 (8)	0.0088 (8)	0.0137 (8)
O3	0.0489 (14)	0.0507 (15)	0.0297 (12)	-0.0048 (12)	0.0180 (10)	-0.0073 (10)
O4	0.0253 (10)	0.0371 (11)	0.0373 (11)	0.0056 (9)	0.0163 (9)	0.0063 (9)
O5	0.0217 (10)	0.0412 (12)	0.0302 (11)	0.0043 (9)	0.0097 (8)	-0.0004 (9)
O6	0.0222 (10)	0.0322 (11)	0.0377 (12)	-0.0033 (8)	0.0008 (8)	0.0031 (9)
N1	0.0168 (9)	0.0142 (9)	0.0159 (9)	0.0048 (7)	0.0058 (7)	0.0036 (7)
N2	0.0203 (10)	0.0182 (10)	0.0184 (10)	0.0058 (8)	0.0056 (8)	0.0073 (8)
N3	0.0200 (10)	0.0176 (10)	0.0196 (10)	0.0061 (8)	0.0052 (8)	0.0063 (8)
N4	0.0271 (12)	0.0264 (11)	0.0255 (11)	0.0053 (9)	0.0100 (9)	0.0072 (9)
N5	0.0207 (10)	0.0242 (11)	0.0249 (11)	0.0093 (9)	0.0069 (8)	0.0105 (9)
C1	0.0185 (11)	0.0171 (11)	0.0258 (12)	0.0041 (9)	0.0082 (9)	0.0065 (9)

C2	0.0253 (12)	0.0146 (11)	0.0150 (11)	0.0036 (9)	0.0053 (9)	0.0018 (9)
C3	0.0313 (14)	0.0178 (12)	0.0194 (12)	0.0017 (10)	0.0049 (10)	0.0046 (9)
C4	0.0402 (16)	0.0172 (12)	0.0217 (12)	0.0043 (11)	-0.0005 (11)	0.0077 (10)
C5	0.0318 (14)	0.0210 (12)	0.0268 (13)	0.0094 (11)	-0.0017 (11)	0.0068 (10)
C6	0.0228 (12)	0.0215 (12)	0.0246 (12)	0.0059 (10)	0.0021 (10)	0.0062 (10)
C7	0.0161 (11)	0.0226 (12)	0.0180 (11)	0.0046 (9)	0.0046 (9)	0.0067 (9)
C8	0.0190 (11)	0.0190 (11)	0.0137 (10)	0.0065 (9)	0.0043 (8)	0.0035 (9)
C9	0.0194 (11)	0.0241 (12)	0.0181 (11)	0.0107 (9)	0.0047 (9)	0.0047 (9)
C10	0.0312 (14)	0.0218 (12)	0.0167 (11)	0.0142 (10)	0.0018 (10)	0.0039 (9)
C11	0.0302 (13)	0.0173 (11)	0.0192 (12)	0.0049 (10)	0.0045 (10)	0.0071 (9)
C12	0.0215 (12)	0.0201 (12)	0.0201 (12)	0.0040 (9)	0.0048 (9)	0.0078 (9)
C13	0.0159 (11)	0.0167 (10)	0.0152 (10)	0.0023 (8)	0.0048 (8)	0.0036 (8)
C14	0.0184 (11)	0.0199 (11)	0.0196 (12)	0.0059 (9)	0.0081 (9)	0.0047 (10)

Geometric parameters (Å, °)

Cd1—N2	2.250 (2)	C3—C4	1.381 (4)
Cd1—N3	2.251 (2)	C3—H3A	0.9500
Cd1—O5	2.279 (2)	C4—C5	1.389 (4)
Cd1—O1	2.322 (2)	C4—H4A	0.9500
Cd1—O2	2.588 (2)	C5—C6	1.379 (4)
Cd1—O4	2.687 (2)	C5—H5	0.9500
Cd1—N1	2.427 (2)	C6—H6A	0.9500
O1—N4	1.273 (3)	C7—C8	1.521 (3)
O2—N4	1.254 (3)	C7—H7A	0.9900
O3—N4	1.231 (3)	C7—H7B	0.9900
O4—N5	1.243 (3)	C8—C9	1.387 (3)
O5—N5	1.276 (3)	C9—C10	1.391 (4)
O6—N5	1.233 (3)	C9—H9	0.9500
N1—C1	1.477 (3)	C10—C11	1.382 (4)
N1—C7	1.479 (3)	C10—H10	0.9500
N1—C13	1.485 (3)	C11—C12	1.381 (4)
N2—C8	1.342 (3)	C11—H11	0.9500
N2—C12	1.351 (3)	C12—H12	0.9500
N3—C6	1.344 (3)	C13—C14	1.530 (3)
N3—C2	1.345 (3)	C13—H13A	0.9900
C1—C2	1.514 (4)	C13—H13B	0.9900
C1—H1A	0.9900	C14—C14 ⁱ	1.528 (5)
C1—H1B	0.9900	C14—H14A	0.9900
C2—C3	1.396 (4)	C14—H14B	0.9900
N2—Cd1—N3	148.10 (8)	C3—C4—C5	119.9 (2)
N2—Cd1—O5	103.49 (8)	C3—C4—H4A	120.1
N3—Cd1—O5	103.16 (8)	C5—C4—H4A	120.1
N2—Cd1—O1	98.56 (8)	C6—C5—C4	118.3 (3)
N3—Cd1—O1	103.09 (8)	C6—C5—H5	120.9
O5—Cd1—O1	79.90 (7)	C4—C5—H5	120.9
N2—Cd1—N1	74.12 (7)	N3—C6—C5	122.2 (3)

N3—Cd1—N1	74.27 (7)	N3—C6—H6A	118.9
O5—Cd1—N1	139.43 (7)	C5—C6—H6A	118.9
O1—Cd1—N1	140.62 (7)	N1—C7—C8	112.38 (19)
N4—O1—Cd1	100.89 (15)	N1—C7—H7A	109.1
N5—O5—Cd1	105.38 (15)	C8—C7—H7A	109.1
C1—N1—C7	112.91 (19)	N1—C7—H7B	109.1
C1—N1—C13	113.20 (19)	C8—C7—H7B	109.1
C7—N1—C13	113.10 (19)	H7A—C7—H7B	107.9
C1—N1—Cd1	104.04 (14)	N2—C8—C9	121.2 (2)
C7—N1—Cd1	103.73 (14)	N2—C8—C7	117.4 (2)
C13—N1—Cd1	108.92 (13)	C9—C8—C7	121.2 (2)
C8—N2—C12	119.4 (2)	C8—C9—C10	119.3 (2)
C8—N2—Cd1	116.61 (16)	C8—C9—H9	120.4
C12—N2—Cd1	123.99 (17)	C10—C9—H9	120.4
C6—N3—C2	119.9 (2)	C11—C10—C9	119.4 (2)
C6—N3—Cd1	123.75 (17)	C11—C10—H10	120.3
C2—N3—Cd1	116.38 (17)	C9—C10—H10	120.3
O3—N4—O2	121.4 (2)	C12—C11—C10	118.4 (2)
O3—N4—O1	120.3 (2)	C12—C11—H11	120.8
O2—N4—O1	118.3 (2)	C10—C11—H11	120.8
O6—N5—O4	122.7 (2)	N2—C12—C11	122.3 (2)
O6—N5—O5	120.0 (2)	N2—C12—H12	118.8
O4—N5—O5	117.3 (2)	C11—C12—H12	118.8
N1—C1—C2	112.7 (2)	N1—C13—C14	116.85 (19)
N1—C1—H1A	109.0	N1—C13—H13A	108.1
C2—C1—H1A	109.0	C14—C13—H13A	108.1
N1—C1—H1B	109.0	N1—C13—H13B	108.1
C2—C1—H1B	109.0	C14—C13—H13B	108.1
H1A—C1—H1B	107.8	H13A—C13—H13B	107.3
N3—C2—C3	120.8 (2)	C14 ⁱ —C14—C13	110.4 (2)
N3—C2—C1	117.8 (2)	C14 ⁱ —C14—H14A	109.6
C3—C2—C1	121.2 (2)	C13—C14—H14A	109.6
C4—C3—C2	119.0 (3)	C14 ⁱ —C14—H14B	109.6
C4—C3—H3A	120.5	C13—C14—H14B	109.6
C2—C3—H3A	120.5	H14A—C14—H14B	108.1
N2—Cd1—O1—N4	-82.99 (18)	Cd1—O5—N5—O6	-176.1 (2)
N3—Cd1—O1—N4	73.42 (18)	Cd1—O5—N5—O4	4.4 (3)
O5—Cd1—O1—N4	174.73 (19)	C7—N1—C1—C2	152.9 (2)
N1—Cd1—O1—N4	-7.6 (2)	C13—N1—C1—C2	-77.0 (3)
N2—Cd1—O5—N5	74.64 (18)	Cd1—N1—C1—C2	41.1 (2)
N3—Cd1—O5—N5	-87.67 (18)	C6—N3—C2—C3	1.3 (3)
O1—Cd1—O5—N5	171.11 (18)	Cd1—N3—C2—C3	-178.25 (17)
N1—Cd1—O5—N5	-6.6 (2)	C6—N3—C2—C1	-173.2 (2)
N2—Cd1—N1—C1	147.56 (15)	Cd1—N3—C2—C1	7.2 (3)
N3—Cd1—N1—C1	-28.13 (14)	N1—C1—C2—N3	-35.6 (3)
O5—Cd1—N1—C1	-120.12 (16)	N1—C1—C2—C3	149.8 (2)
O1—Cd1—N1—C1	63.46 (19)	N3—C2—C3—C4	-1.6 (4)

N2—Cd1—N1—C7	29.25 (14)	C1—C2—C3—C4	172.8 (2)
N3—Cd1—N1—C7	-146.43 (15)	C2—C3—C4—C5	0.6 (4)
O5—Cd1—N1—C7	121.57 (16)	C3—C4—C5—C6	0.7 (4)
O1—Cd1—N1—C7	-54.85 (19)	C2—N3—C6—C5	0.0 (4)
N2—Cd1—N1—C13	-91.45 (15)	Cd1—N3—C6—C5	179.57 (19)
N3—Cd1—N1—C13	92.86 (15)	C4—C5—C6—N3	-1.0 (4)
O5—Cd1—N1—C13	0.9 (2)	C1—N1—C7—C8	-154.2 (2)
O1—Cd1—N1—C13	-175.55 (14)	C13—N1—C7—C8	75.6 (2)
N3—Cd1—N2—C8	-5.2 (3)	Cd1—N1—C7—C8	-42.3 (2)
O5—Cd1—N2—C8	-151.19 (16)	C12—N2—C8—C9	-0.5 (3)
O1—Cd1—N2—C8	127.22 (17)	Cd1—N2—C8—C9	178.50 (17)
N1—Cd1—N2—C8	-13.12 (16)	C12—N2—C8—C7	174.4 (2)
N3—Cd1—N2—C12	173.73 (17)	Cd1—N2—C8—C7	-6.6 (3)
O5—Cd1—N2—C12	27.8 (2)	N1—C7—C8—N2	36.1 (3)
O1—Cd1—N2—C12	-53.8 (2)	N1—C7—C8—C9	-149.0 (2)
N1—Cd1—N2—C12	165.9 (2)	N2—C8—C9—C10	1.4 (4)
N2—Cd1—N3—C6	-175.28 (17)	C7—C8—C9—C10	-173.3 (2)
O5—Cd1—N3—C6	-29.3 (2)	C8—C9—C10—C11	-1.2 (4)
O1—Cd1—N3—C6	53.2 (2)	C9—C10—C11—C12	0.1 (4)
N1—Cd1—N3—C6	-167.4 (2)	C8—N2—C12—C11	-0.7 (4)
N2—Cd1—N3—C2	4.3 (3)	Cd1—N2—C12—C11	-179.60 (18)
O5—Cd1—N3—C2	150.30 (16)	C10—C11—C12—N2	0.9 (4)
O1—Cd1—N3—C2	-127.20 (17)	C1—N1—C13—C14	-62.6 (3)
N1—Cd1—N3—C2	12.17 (16)	C7—N1—C13—C14	67.4 (3)
Cd1—O1—N4—O3	178.6 (2)	Cd1—N1—C13—C14	-177.82 (17)
Cd1—O1—N4—O2	-1.2 (3)	N1—C13—C14—C14 ⁱ	-175.1 (2)

Symmetry code: (i) $-x, -y+2, -z+1$.