

## 2-Dicyclohexylamino-3-phenyl-5,6-di-hydro-8H-thiopyrano[4',3':4,5]thieno-[2,3-d]pyrimidin-4(3H)-one

Xizhen Liang<sup>a,b\*</sup> and Yueming Zhou<sup>a</sup>

<sup>a</sup>Faculty of Biology, Chemistry and Material Science, East China Institute of Technology, Nanchang 330013, Jiangxi, People's Republic of China, and <sup>b</sup>College of Chemistry and Chemical Engineering, Central South University, Changsha 410083, Hunan, People's Republic of China

Correspondence e-mail: lxzhen11@gmail.com


Received 18 February 2011; accepted 10 March 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.039;  $wR$  factor = 0.097; data-to-parameter ratio = 14.6.

In the title compound,  $\text{C}_{27}\text{H}_{33}\text{N}_3\text{OS}_2$ , the dihedral angle between the two fused rings of the thieno[3,2-*d*]pyrimidine system is  $3.73(9)^\circ$ . The phenyl ring is twisted with respect to the pyrimidine ring [dihedral angle =  $71.60(10)^\circ$ ] and the thiopyran ring shows an envelope conformation with the S atom as the flap. An intramolecular C—H···O interaction occurs. In the crystal, inversion dimers linked by pairs of C—H···O hydrogen bonds occur.

### Related literature

For the properties of compounds containing the thiopyrimidine system, see: Santagati *et al.* (2002); Kikuchi *et al.* (2006). For related crystal structures, see: Hu *et al.* (2007); Xie *et al.* (2008).



### Experimental

#### Crystal data

$\text{C}_{27}\text{H}_{33}\text{N}_3\text{OS}_2$	$V = 2486.1(4)\text{ \AA}^3$
$M_r = 479.70$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 10.7118(10)\text{ \AA}$	$\mu = 0.24\text{ mm}^{-1}$
$b = 22.601(2)\text{ \AA}$	$T = 293\text{ K}$
$c = 10.924(1)\text{ \AA}$	$0.30 \times 0.25 \times 0.18\text{ mm}$
$\beta = 109.939(1)^\circ$	

#### Data collection

Bruker SMART CCD area-detector diffractometer	12466 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2007)	4365 independent reflections
$T_{\min} = 0.802$ , $T_{\max} = 0.874$	3391 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.027$
	Standard reflections: ?

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	298 parameters
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
4365 reflections	$\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots\text{O}1^{\text{i}}$	0.97	2.54	3.359 (3)	142
$\text{C}11-\text{H}11\cdots\text{O}1^{\text{ii}}$	0.93	2.57	3.199 (3)	126

Symmetry codes: (i)  $-x, -y, -z + 1$ ; (ii)  $-x, -y, -z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported financially by the Jiangxi Natural Science Foundation of China (grant No. 2009GZH0002).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5802).

### References

- Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hu, Y.-G., Li, G.-H. & Zhou, M.-H. (2007). *Acta Cryst. E63*, o1836–o1838.
- Kikuchi, H., Yamamoto, K., Horoiwa, S., Hirai, S., Kasahara, R., Hariguchi, N., Matsumoto, M. & Oshima, Y. (2006). *J. Med. Chem.* **49**, 4698–4706.
- Santagati, N. A., Prezzavento, O., Bousquet, E., Ronisvalle, G. & Spampinato, S. (2002). *J. Pharm. Pharmacol.* **54**, 717–728.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Xie, H., Meng, S.-M., Fan, Y.-Q. & Guo, Y. (2008). *Acta Cryst. E64*, o2434.

# supporting information

*Acta Cryst.* (2011). E67, o888 [doi:10.1107/S1600536811009226]

## **2-Dicyclohexylamino-3-phenyl-5,6-dihydro-8H-thiopyrano[4',3':4,5]thieno[2,3-d]pyrimidin-4(3H)-one**


**Xizhen Liang and Yueming Zhou**

### **S1. Comment**

The derivatives of heterocycles containing thienopyrimidine system are of great importance because of their remarkable biological activity for use as potential drugs. They proved to show significant antifungal, antibacterial, antimicrobial, anticonvulsant and angiotensin antagonistic activities. Also some of these compounds show good antimalarial or potent multitargeted receptor tyrosine kinase inhibitive activities (Santagati *et al.*, 2002; Kikuchi *et al.*, 2006). In addition, the title compound have exhibited a high potential as efficient luminophores for liquid and fluorescent properties and markers in biological or supramolecular systems. Some X-ray crystal structures of pyrimidinone derivatives have been reported (Hu *et al.*, 2007; Xie *et al.*, 2008). The bond lengths and angles are unexceptional. The thieno (A), the pyrimidinone (B) and the C9—C14 benzene (C), rings are, of course, planar and the dihedral angles between them are A/B = 3.73 (9) °, B/C = 71.60 (10) °. The thiopyrano ring in (I) (Fig 1) shows a distorted chair conformation [ $\varphi$  = 164.0 (3) ° and  $\theta$  = 128.83 (18) °, Puckering Amplitude = 0.6360 (18) %A]. The weak intermolecular C—H···O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimmers (Fig 2).


### **S2. Experimental**

To a solution of iminophosphorane (1.01 g, 2 mmol) in anhyd. CH<sub>2</sub>Cl<sub>2</sub> (15 ml) aromatic isocyanate (2.2 mmol) was added under nitrogen atmosphere at room temperature. After the reaction mixture was left unstirred for 6–12 h at 0–5°C, the iminophosphorane had disappeared (TLC monitored). The solvent was removed off under reduced pressure and Et<sub>2</sub>O / petroleum ether (1:2, 20 ml) was added to precipitate triphenylphosphine oxide. Removal of the solvent gave carbodi-imides, which were used directly without further purification. To the solution of in dichloromethane (15 ml) was added dialkylamine (2.2 mmol). After the reaction mixture was left unstirred for 4–6 h, the solvent was removed and anhyd. EtOH (15 mL) with several drops of EtONa in EtOH was added. The mixture was stirred for 6–12 h at room temperature. The solution was condensed and residue was recrystallized from EtOH to give yellow blocks of the title compound.



**Figure 1**

The molecular structure of (I) showing 50% displacement ellipsoids.

**Figure 2**

Packing diagram for (I) with C—H···O links shown as dashed lines.

### **2-Dicyclohexylamino-3-phenyl-5,6-dihydro-8*H*- thiopyrano[4',3':4,5]thieno[2,3-*d*]pyrimidin-4(3*H*)-one**

#### *Crystal data*

$C_{27}H_{33}N_3OS_2$

$M_r = 479.70$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.7118 (10)$  Å

$b = 22.601 (2)$  Å

$c = 10.924 (1)$  Å

$\beta = 109.939 (1)^\circ$

$V = 2486.1 (4)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 1024$

$D_x = 1.282 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4365 reflections

$\theta = 2.0\text{--}25.0^\circ$

$\mu = 0.24 \text{ mm}^{-1}$

$T = 293$  K

Block, yellow

$0.30 \times 0.25 \times 0.18$  mm

#### *Data collection*

Bruker DIFFRACTOMETER TYPE?????

    diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Absorption correction: multi-scan

    (SADABS; Bruker, 2007)

$T_{\min} = 0.802, T_{\max} = 0.874$

12466 measured reflections

4365 independent reflections

3391 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 25.0^\circ, \theta_{\min} = 2.0^\circ$

$h = -12 \rightarrow 11$   
 $k = -17 \rightarrow 26$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.097$

$S = 1.00$

4365 reflections

298 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

$l = -12 \rightarrow 12$

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0373P)^2 + 1.0362P]$   
where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.52363 (5)	0.09045 (3)	0.58192 (5)	0.05635 (17)
S2	0.36688 (6)	-0.00416 (3)	0.84325 (5)	0.06459 (19)
N2	0.39389 (15)	0.14368 (7)	0.35103 (15)	0.0446 (4)
N1	0.17515 (14)	0.11220 (7)	0.23197 (14)	0.0381 (4)
O1	0.06017 (13)	0.04696 (6)	0.31444 (12)	0.0520 (4)
C6	0.28078 (18)	0.07129 (8)	0.44376 (17)	0.0392 (4)
C7	0.38904 (18)	0.10337 (8)	0.44241 (18)	0.0414 (4)
C4	0.30430 (18)	0.03617 (8)	0.55936 (17)	0.0408 (4)
C8	0.28574 (17)	0.14842 (8)	0.24936 (17)	0.0382 (4)
C5	0.43038 (19)	0.04268 (9)	0.64211 (18)	0.0456 (5)
C3	0.2018 (2)	-0.00226 (10)	0.5853 (2)	0.0532 (5)
H3A	0.1593	-0.0265	0.5091	0.064*
H3B	0.1342	0.0228	0.5990	0.064*
C1	0.4921 (2)	0.01507 (10)	0.7740 (2)	0.0568 (6)
H1A	0.5547	0.0426	0.8312	0.068*
H1B	0.5405	-0.0202	0.7664	0.068*
C2	0.2587 (2)	-0.04225 (10)	0.7023 (2)	0.0587 (6)
H2A	0.3078	-0.0740	0.6798	0.070*
H2B	0.1862	-0.0599	0.7238	0.070*
N3	0.27679 (15)	0.18922 (7)	0.15363 (15)	0.0417 (4)
C9	0.08042 (18)	0.10101 (8)	0.10264 (17)	0.0392 (4)
C22	0.39403 (19)	0.20341 (9)	0.11691 (19)	0.0455 (5)
H22	0.3599	0.2280	0.0386	0.055*

C20	0.16227 (19)	0.23014 (9)	0.11292 (18)	0.0455 (5)
H20	0.0966	0.2139	0.1481	0.055*
C10	0.12331 (19)	0.07079 (9)	0.01470 (18)	0.0447 (5)
H10	0.2115	0.0589	0.0379	0.054*
C14	-0.05072 (19)	0.11717 (9)	0.07028 (19)	0.0485 (5)
H14	-0.0803	0.1362	0.1307	0.058*
C27	0.5036 (2)	0.23963 (10)	0.2115 (2)	0.0563 (5)
H27A	0.5426	0.2176	0.2918	0.068*
H27B	0.4669	0.2760	0.2319	0.068*
C23	0.4467 (2)	0.14866 (10)	0.0736 (2)	0.0576 (6)
H23A	0.3750	0.1290	0.0065	0.069*
H23B	0.4799	0.1217	0.1466	0.069*
C13	-0.1378 (2)	0.10476 (11)	-0.0530 (2)	0.0595 (6)
H13	-0.2263	0.1162	-0.0762	0.071*
C12	-0.0948 (2)	0.07562 (11)	-0.1418 (2)	0.0597 (6)
H12	-0.1538	0.0677	-0.2249	0.072*
C15	0.0939 (2)	0.23351 (11)	-0.0339 (2)	0.0597 (6)
H15A	0.0736	0.1939	-0.0692	0.072*
H15B	0.1529	0.2519	-0.0731	0.072*
C26	0.6108 (2)	0.25425 (11)	0.1533 (3)	0.0683 (7)
H26A	0.5735	0.2798	0.0783	0.082*
H26B	0.6822	0.2756	0.2172	0.082*
C19	0.1943 (2)	0.29232 (9)	0.1691 (2)	0.0593 (6)
H19A	0.2379	0.2901	0.2627	0.071*
H19B	0.2542	0.3116	0.1322	0.071*
C11	0.0352 (2)	0.05831 (10)	-0.1077 (2)	0.0533 (5)
H11	0.0640	0.0381	-0.1673	0.064*
C24	0.5580 (2)	0.16290 (11)	0.0206 (2)	0.0696 (7)
H24A	0.5961	0.1262	0.0035	0.083*
H24B	0.5211	0.1839	-0.0614	0.083*
C25	0.6657 (2)	0.19973 (11)	0.1127 (3)	0.0735 (7)
H25A	0.7126	0.1765	0.1892	0.088*
H25B	0.7289	0.2110	0.0709	0.088*
C16	-0.0336 (2)	0.26934 (13)	-0.0668 (2)	0.0789 (8)
H16A	-0.0734	0.2729	-0.1606	0.095*
H16B	-0.0959	0.2486	-0.0352	0.095*
C17	-0.0084 (3)	0.33031 (13)	-0.0069 (3)	0.0926 (10)
H17A	-0.0927	0.3501	-0.0222	0.111*
H17B	0.0419	0.3532	-0.0491	0.111*
C18	0.0670 (3)	0.32814 (12)	0.1378 (3)	0.0844 (8)
H18A	0.0110	0.3106	0.1816	0.101*
H18B	0.0886	0.3681	0.1704	0.101*
C28	0.16231 (18)	0.07445 (8)	0.33148 (17)	0.0387 (4)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0443 (3)	0.0596 (4)	0.0528 (3)	-0.0111 (3)	0.0006 (2)	0.0144 (3)

S2	0.0742 (4)	0.0727 (4)	0.0432 (3)	-0.0018 (3)	0.0152 (3)	0.0092 (3)
N2	0.0419 (9)	0.0445 (9)	0.0433 (9)	-0.0061 (7)	0.0092 (7)	0.0063 (8)
N1	0.0373 (8)	0.0404 (9)	0.0349 (8)	-0.0025 (7)	0.0104 (6)	0.0018 (7)
O1	0.0453 (8)	0.0625 (9)	0.0443 (8)	-0.0166 (7)	0.0101 (6)	0.0039 (7)
C6	0.0411 (10)	0.0366 (10)	0.0386 (10)	-0.0021 (8)	0.0117 (8)	-0.0009 (8)
C7	0.0404 (10)	0.0397 (11)	0.0413 (10)	-0.0016 (9)	0.0103 (8)	0.0024 (9)
C4	0.0457 (11)	0.0351 (10)	0.0383 (10)	-0.0020 (8)	0.0100 (8)	0.0004 (8)
C8	0.0394 (10)	0.0370 (10)	0.0388 (10)	0.0004 (8)	0.0141 (8)	-0.0005 (8)
C5	0.0487 (11)	0.0398 (11)	0.0426 (10)	-0.0043 (9)	0.0079 (9)	0.0034 (9)
C3	0.0517 (12)	0.0558 (13)	0.0463 (12)	-0.0094 (10)	0.0093 (10)	0.0103 (10)
C1	0.0571 (13)	0.0532 (13)	0.0480 (12)	-0.0049 (11)	0.0022 (10)	0.0092 (10)
C2	0.0591 (13)	0.0567 (14)	0.0568 (13)	-0.0099 (11)	0.0152 (11)	0.0150 (11)
N3	0.0416 (9)	0.0401 (9)	0.0445 (9)	0.0000 (7)	0.0162 (7)	0.0072 (7)
C9	0.0384 (10)	0.0393 (10)	0.0369 (10)	-0.0018 (8)	0.0091 (8)	0.0029 (8)
C22	0.0475 (11)	0.0442 (11)	0.0483 (11)	-0.0031 (9)	0.0208 (9)	0.0034 (9)
C20	0.0482 (11)	0.0447 (11)	0.0460 (11)	0.0075 (9)	0.0193 (9)	0.0099 (9)
C10	0.0422 (10)	0.0426 (11)	0.0487 (11)	-0.0026 (9)	0.0147 (9)	-0.0026 (9)
C14	0.0404 (11)	0.0585 (13)	0.0469 (11)	0.0016 (10)	0.0151 (9)	0.0003 (10)
C27	0.0519 (12)	0.0535 (13)	0.0639 (13)	-0.0087 (10)	0.0202 (11)	-0.0068 (11)
C23	0.0582 (13)	0.0525 (13)	0.0680 (14)	-0.0024 (11)	0.0292 (11)	-0.0060 (11)
C13	0.0364 (11)	0.0755 (16)	0.0592 (13)	0.0009 (11)	0.0066 (10)	0.0043 (12)
C12	0.0542 (13)	0.0705 (16)	0.0438 (12)	-0.0097 (12)	0.0032 (10)	-0.0024 (11)
C15	0.0596 (13)	0.0700 (15)	0.0468 (12)	0.0126 (12)	0.0146 (10)	0.0122 (11)
C26	0.0562 (14)	0.0533 (14)	0.0955 (18)	-0.0121 (11)	0.0261 (13)	0.0016 (13)
C19	0.0732 (15)	0.0461 (12)	0.0663 (14)	0.0049 (11)	0.0339 (12)	0.0014 (11)
C11	0.0595 (13)	0.0543 (13)	0.0468 (12)	-0.0083 (11)	0.0190 (10)	-0.0084 (10)
C24	0.0741 (16)	0.0686 (16)	0.0815 (17)	0.0040 (13)	0.0466 (14)	-0.0009 (14)
C25	0.0535 (14)	0.0666 (16)	0.111 (2)	0.0017 (13)	0.0422 (14)	0.0132 (15)
C16	0.0654 (16)	0.100 (2)	0.0686 (16)	0.0253 (15)	0.0187 (13)	0.0350 (16)
C17	0.094 (2)	0.078 (2)	0.119 (3)	0.0398 (17)	0.0537 (19)	0.0499 (19)
C18	0.104 (2)	0.0562 (16)	0.109 (2)	0.0245 (15)	0.0563 (19)	0.0098 (16)
C28	0.0406 (10)	0.0377 (10)	0.0375 (10)	-0.0027 (9)	0.0130 (8)	-0.0020 (8)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

S1—C7	1.7286 (19)	C14—C13	1.381 (3)
S1—C5	1.745 (2)	C14—H14	0.9300
S2—C2	1.799 (2)	C27—C26	1.526 (3)
S2—C1	1.803 (2)	C27—H27A	0.9700
N2—C8	1.308 (2)	C27—H27B	0.9700
N2—C7	1.365 (2)	C23—C24	1.527 (3)
N1—C8	1.399 (2)	C23—H23A	0.9700
N1—C28	1.425 (2)	C23—H23B	0.9700
N1—C9	1.453 (2)	C13—C12	1.375 (3)
O1—C28	1.216 (2)	C13—H13	0.9300
C6—C7	1.372 (3)	C12—C11	1.370 (3)
C6—C28	1.435 (2)	C12—H12	0.9300
C6—C4	1.439 (2)	C15—C16	1.522 (3)

C4—C5	1.353 (3)	C15—H15A	0.9700
C4—C3	1.500 (3)	C15—H15B	0.9700
C8—N3	1.373 (2)	C26—C25	1.497 (3)
C5—C1	1.501 (3)	C26—H26A	0.9700
C3—C2	1.514 (3)	C26—H26B	0.9700
C3—H3A	0.9700	C19—C18	1.521 (3)
C3—H3B	0.9700	C19—H19A	0.9700
C1—H1A	0.9700	C19—H19B	0.9700
C1—H1B	0.9700	C11—H11	0.9300
C2—H2A	0.9700	C24—C25	1.498 (3)
C2—H2B	0.9700	C24—H24A	0.9700
N3—C22	1.478 (2)	C24—H24B	0.9700
N3—C20	1.478 (2)	C25—H25A	0.9700
C9—C14	1.376 (3)	C25—H25B	0.9700
C9—C10	1.379 (3)	C16—C17	1.509 (4)
C22—C23	1.502 (3)	C16—H16A	0.9700
C22—C27	1.513 (3)	C16—H16B	0.9700
C22—H22	0.9800	C17—C18	1.511 (4)
C20—C15	1.521 (3)	C17—H17A	0.9700
C20—C19	1.525 (3)	C17—H17B	0.9700
C20—H20	0.9800	C18—H18A	0.9700
C10—C11	1.377 (3)	C18—H18B	0.9700
C10—H10	0.9300		
C7—S1—C5	91.39 (9)	H27A—C27—H27B	108.1
C2—S2—C1	96.33 (10)	C22—C23—C24	111.83 (19)
C8—N2—C7	115.60 (16)	C22—C23—H23A	109.2
C8—N1—C28	122.99 (14)	C24—C23—H23A	109.2
C8—N1—C9	120.81 (14)	C22—C23—H23B	109.2
C28—N1—C9	114.86 (14)	C24—C23—H23B	109.2
C7—C6—C28	117.97 (17)	H23A—C23—H23B	107.9
C7—C6—C4	113.91 (16)	C12—C13—C14	120.5 (2)
C28—C6—C4	128.07 (17)	C12—C13—H13	119.8
N2—C7—C6	127.12 (17)	C14—C13—H13	119.8
N2—C7—S1	121.94 (14)	C11—C12—C13	119.9 (2)
C6—C7—S1	110.80 (14)	C11—C12—H12	120.0
C5—C4—C6	111.25 (17)	C13—C12—H12	120.0
C5—C4—C3	124.39 (17)	C20—C15—C16	110.26 (18)
C6—C4—C3	124.36 (16)	C20—C15—H15A	109.6
N2—C8—N3	120.91 (16)	C16—C15—H15A	109.6
N2—C8—N1	122.10 (16)	C20—C15—H15B	109.6
N3—C8—N1	116.99 (15)	C16—C15—H15B	109.6
C4—C5—C1	127.50 (18)	H15A—C15—H15B	108.1
C4—C5—S1	112.64 (14)	C25—C26—C27	111.91 (19)
C1—C5—S1	119.85 (14)	C25—C26—H26A	109.2
C4—C3—C2	113.29 (17)	C27—C26—H26A	109.2
C4—C3—H3A	108.9	C25—C26—H26B	109.2
C2—C3—H3A	108.9	C27—C26—H26B	109.2

C4—C3—H3B	108.9	H26A—C26—H26B	107.9
C2—C3—H3B	108.9	C18—C19—C20	109.8 (2)
H3A—C3—H3B	107.7	C18—C19—H19A	109.7
C5—C1—S2	110.84 (15)	C20—C19—H19A	109.7
C5—C1—H1A	109.5	C18—C19—H19B	109.7
S2—C1—H1A	109.5	C20—C19—H19B	109.7
C5—C1—H1B	109.5	H19A—C19—H19B	108.2
S2—C1—H1B	109.5	C12—C11—C10	120.1 (2)
H1A—C1—H1B	108.1	C12—C11—H11	119.9
C3—C2—S2	113.16 (15)	C10—C11—H11	119.9
C3—C2—H2A	108.9	C25—C24—C23	112.6 (2)
S2—C2—H2A	108.9	C25—C24—H24A	109.1
C3—C2—H2B	108.9	C23—C24—H24A	109.1
S2—C2—H2B	108.9	C25—C24—H24B	109.1
H2A—C2—H2B	107.8	C23—C24—H24B	109.1
C8—N3—C22	120.49 (15)	H24A—C24—H24B	107.8
C8—N3—C20	118.98 (14)	C26—C25—C24	111.5 (2)
C22—N3—C20	118.51 (15)	C26—C25—H25A	109.3
C14—C9—C10	120.40 (18)	C24—C25—H25A	109.3
C14—C9—N1	121.13 (17)	C26—C25—H25B	109.3
C10—C9—N1	118.38 (16)	C24—C25—H25B	109.3
N3—C22—C23	110.62 (16)	H25A—C25—H25B	108.0
N3—C22—C27	117.73 (16)	C17—C16—C15	111.7 (2)
C23—C22—C27	111.94 (17)	C17—C16—H16A	109.3
N3—C22—H22	105.1	C15—C16—H16A	109.3
C23—C22—H22	105.1	C17—C16—H16B	109.3
C27—C22—H22	105.1	C15—C16—H16B	109.3
N3—C20—C15	113.60 (16)	H16A—C16—H16B	107.9
N3—C20—C19	113.94 (17)	C16—C17—C18	112.1 (2)
C15—C20—C19	109.83 (18)	C16—C17—H17A	109.2
N3—C20—H20	106.3	C18—C17—H17A	109.2
C15—C20—H20	106.3	C16—C17—H17B	109.2
C19—C20—H20	106.3	C18—C17—H17B	109.2
C11—C10—C9	119.78 (19)	H17A—C17—H17B	107.9
C11—C10—H10	120.1	C17—C18—C19	111.8 (2)
C9—C10—H10	120.1	C17—C18—H18A	109.3
C9—C14—C13	119.21 (19)	C19—C18—H18A	109.3
C9—C14—H14	120.4	C17—C18—H18B	109.3
C13—C14—H14	120.4	C19—C18—H18B	109.3
C22—C27—C26	110.57 (18)	H18A—C18—H18B	107.9
C22—C27—H27A	109.5	O1—C28—N1	120.33 (16)
C26—C27—H27A	109.5	O1—C28—C6	126.36 (17)
C22—C27—H27B	109.5	N1—C28—C6	113.24 (16)
C26—C27—H27B	109.5		
C8—N2—C7—C6	-5.7 (3)	C20—N3—C22—C23	-138.37 (18)
C8—N2—C7—S1	179.04 (14)	C8—N3—C22—C27	-72.5 (2)
C28—C6—C7—N2	7.6 (3)	C20—N3—C22—C27	91.2 (2)

C4—C6—C7—N2	-174.80 (18)	C8—N3—C20—C15	-131.05 (19)
C28—C6—C7—S1	-176.68 (14)	C22—N3—C20—C15	65.0 (2)
C4—C6—C7—S1	0.9 (2)	C8—N3—C20—C19	102.2 (2)
C5—S1—C7—N2	174.90 (17)	C22—N3—C20—C19	-61.8 (2)
C5—S1—C7—C6	-1.09 (16)	C14—C9—C10—C11	2.0 (3)
C7—C6—C4—C5	-0.2 (2)	N1—C9—C10—C11	178.53 (17)
C28—C6—C4—C5	177.14 (19)	C10—C9—C14—C13	-2.4 (3)
C7—C6—C4—C3	179.31 (19)	N1—C9—C14—C13	-178.87 (18)
C28—C6—C4—C3	-3.4 (3)	N3—C22—C27—C26	-175.41 (18)
C7—N2—C8—N3	177.25 (16)	C23—C22—C27—C26	54.7 (2)
C7—N2—C8—N1	-3.5 (3)	N3—C22—C23—C24	173.60 (18)
C28—N1—C8—N2	10.6 (3)	C27—C22—C23—C24	-53.0 (2)
C9—N1—C8—N2	-155.50 (17)	C9—C14—C13—C12	1.2 (3)
C28—N1—C8—N3	-170.12 (16)	C14—C13—C12—C11	0.5 (4)
C9—N1—C8—N3	23.8 (2)	N3—C20—C15—C16	171.91 (19)
C6—C4—C5—C1	178.0 (2)	C19—C20—C15—C16	-59.2 (2)
C3—C4—C5—C1	-1.4 (3)	C22—C27—C26—C25	-55.9 (3)
C6—C4—C5—S1	-0.7 (2)	N3—C20—C19—C18	-171.88 (18)
C3—C4—C5—S1	179.85 (16)	C15—C20—C19—C18	59.4 (2)
C7—S1—C5—C4	1.02 (17)	C13—C12—C11—C10	-1.0 (3)
C7—S1—C5—C1	-177.80 (18)	C9—C10—C11—C12	-0.3 (3)
C5—C4—C3—C2	-10.5 (3)	C22—C23—C24—C25	52.1 (3)
C6—C4—C3—C2	170.07 (19)	C27—C26—C25—C24	55.1 (3)
C4—C5—C1—S2	-23.8 (3)	C23—C24—C25—C26	-53.0 (3)
S1—C5—C1—S2	154.80 (12)	C20—C15—C16—C17	55.8 (3)
C2—S2—C1—C5	49.27 (17)	C15—C16—C17—C18	-52.8 (3)
C4—C3—C2—S2	47.5 (2)	C16—C17—C18—C19	53.3 (3)
C1—S2—C2—C3	-64.11 (18)	C20—C19—C18—C17	-56.4 (3)
N2—C8—N3—C22	35.6 (3)	C8—N1—C28—O1	174.84 (17)
N1—C8—N3—C22	-143.67 (16)	C9—N1—C28—O1	-18.3 (2)
N2—C8—N3—C20	-127.97 (19)	C8—N1—C28—C6	-8.0 (2)
N1—C8—N3—C20	52.7 (2)	C9—N1—C28—C6	158.83 (16)
C8—N1—C9—C14	-119.1 (2)	C7—C6—C28—O1	176.52 (19)
C28—N1—C9—C14	73.7 (2)	C4—C6—C28—O1	-0.7 (3)
C8—N1—C9—C10	64.4 (2)	C7—C6—C28—N1	-0.4 (2)
C28—N1—C9—C10	-102.80 (19)	C4—C6—C28—N1	-177.64 (17)
C8—N3—C22—C23	57.9 (2)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C2—H2B <sup>i</sup> —O1 <sup>i</sup>	0.97	2.54	3.359 (3)	142
C11—H11 <sup>ii</sup> —O1 <sup>ii</sup>	0.93	2.57	3.199 (3)	126

Symmetry codes: (i)  $-x, -y, -z+1$ ; (ii)  $-x, -y, -z$ .