

Bis{2-[4-(methylsulfanyl)phenyl]-1*H*-benzimidazol-3-ium} tetrabromidocuprate(II) dihydrate

M. N. Manjunatha,^a Mohamed Ziaulla,^a Ravish Sankolli,^b Noor Shahina Begum^{a*} and K. R. Nagasundara^a

^aDepartment of Chemistry, Bangalore University, Bangalore 560 001, India, and

^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, India

Correspondence e-mail: noorsb@rediffmail.com


Received 15 March 2011; accepted 6 April 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.059; wR factor = 0.127; data-to-parameter ratio = 15.1.

The asymmetric unit of the title compound, $(\text{C}_{14}\text{H}_{13}\text{N}_2\text{S})_2[\text{CuBr}_4] \cdot 2\text{H}_2\text{O}$, contains two cations, one anion and two solvent water molecules that are connected *via* $\text{O}-\text{H} \cdots \text{Br}$, $\text{N}-\text{H} \cdots \text{Br}$ and $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds into a two-dimensional polymeric structure. The cations are arranged in a head-to-tail fashion and form stacks along [100]. The central Cu^{II} atom of the anion is in a distorted tetrahedral environment.

Related literature

For general background to benzimidazoles and their derivatives, see: Huang & Scarborough *et al.* (1999); Preston (1974); Zhu *et al.* (2000). For related structures, see: Ziaulla *et al.* (2011).



Experimental

Crystal data

$(\text{C}_{14}\text{H}_{13}\text{N}_2\text{S})_2[\text{CuBr}_4] \cdot 2\text{H}_2\text{O}$

$M_r = 901.86$

Triclinic, $P\bar{1}$

$a = 7.6878$ (5) Å

$b = 11.8358$ (7) Å

$c = 18.5485$ (9) Å

$\alpha = 85.305$ (4)°

$\beta = 84.778$ (5)°

$\gamma = 80.692$ (5)°

$V = 1654.74$ (17) Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 5.65$ mm⁻¹

$T = 296$ K

$0.18 \times 0.16 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD

detector diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 1998)

$T_{\text{min}} = 0.430$, $T_{\text{max}} = 0.465$

27134 measured reflections

5805 independent reflections

3344 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.110$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$

$wR(F^2) = 0.127$

$S = 1.00$

5805 reflections

384 parameters

6 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.75$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.59$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3} \cdots \text{O1}^{\text{i}}$	0.86	1.86	2.703 (8)	165
$\text{N2}-\text{H2} \cdots \text{Br3}^{\text{ii}}$	0.86	2.44	3.275 (6)	162
$\text{O1}-\text{H1D} \cdots \text{Br3}^{\text{iii}}$	0.85 (6)	2.55 (7)	3.344 (6)	155
$\text{O2}-\text{H2A} \cdots \text{Br2}^{\text{iii}}$	0.83 (4)	2.96 (6)	3.735 (6)	155
$\text{O1}-\text{H1E} \cdots \text{Br1}$	0.84 (7)	2.53 (7)	3.359 (6)	170
$\text{O2}-\text{H2B} \cdots \text{Br4}$	0.85 (5)	2.77 (7)	3.597 (6)	166

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $x - 1, y, z$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *CAMERON* (Watkin *et al.*, 1996); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

NSB is thankful to the University Grants Commission (UGC), India, for financial assistance and the Department of Science and Technology, (DST), India, for the data collection facility under the IRHPA–DST program. MNM thanks the M. S. Ramaiah Institute of Technology, Bangalore, for their support and encouragement.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2356).

References

- Bruker. (1998). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Huang, W. & Scarborough, R. M. (1999). *Tetrahedron Lett.* **40**, 2665–2668.
- Preston, P. N. (1974). *Chem. Rev.* **74**, 279–314.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory, Oxford, England.
- Zhu, Z., Lipka, B., Drach, J. C. & Townsend, L. B. (2000). *J. Med. Chem.* **43**, 2430–2437.
- Ziaulla, M., Manjunatha, M. N., Sankolli, R., Nagasundara, K. R. & Begum, N. S. (2011). *Acta Cryst. E* **67**, o341–o342.

supporting information

Acta Cryst. (2011). E67, m578 [https://doi.org/10.1107/S1600536811012840]

**Bis{2-[4-(methylsulfanyl)phenyl]-1*H*-benzimidazol-3-ium}
tetrabromidocuprate(II) dihydrate**

M. N. Manjunatha, Mohamed Ziaulla, Ravish Sankolli, Noor Shahina Begum and K. R. Nagasundara

S1. Comment

The synthesis of benzimidazoles makes use of solid-phase synthesis *via o*-nitroanilines (Preston *et al.*, 1974; Huang & Scarborough, 1999). Benzimidazole derivatives are effective against the human cytomegalo virus (HCMV) (Zhu *et al.*, 2000). In addition benzimidazole derivatives exhibit a number of important pharmacological properties, such as antihistaminic, anti-ulcerative, antiallergic and antipyretic. In the title compound, as shown in Fig. 1, there are two cations, one tetrabromidocuprate(II) anion and two solvent water molecules in the asymmetric unit. The Cu^{II} atom shows strongly distorted tetrahedral geometry, coordinating with four terminal bromine atoms with the bond lengths in the range 2.3389 (1) Å to 2.4084 (1) Å. The Br—Cu—Br bond angles are between 96.18 (4)° and 139.53 (6)°. The benzimidazole and thiomethyl phenyl rings are virtually planar and inclined at a dihedral angle 2.67 (2)°. The bond lengths and angles for the benzimidazole cation of the molecule are in good agreement, within experimental errors, with those observed in other benzimidazole derivatives (Ziaulla *et al.*, 2011). The crystal structure is stabilized by N—H···O, O—H···Br and N—H···Br hydrogen bonds (Fig.2).

S2. Experimental

An ethanolic solution (15 ml) of the 2-(4-methylsulfanyl phenyl)-1*H*- benzimidazole) (0.960 g, 2 mmol) was added to a solution of copper(II) bromide (0.446 g, 1 mmol) in ethanol (25 ml). The mixture was then treated with 48% HBr (2–3 ml) followed by liquid Br₂ (2–3 ml). The mixture was refluxed for nearly six hours during which yellow crystals suitable for X-ray analysis were obtained. The crystals were washed with cold ethanol and dried in vacuum over P₂O₅ (yield 1.2 g, 85%).

S3. Refinement

The H atoms were placed at calculated positions and refined in the riding model approximation with C—H = 0.93–0.96 Å and N—H = 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. H atoms of water molecules were refined with restraints imposed on the O—H and H···H distances [O—H = 0.85 (2) Å, H···H = 1.39 (4) Å] and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.




Figure 1

ORTEP (Farrugia, 1997) view of the title compound, showing 50% probability ellipsoids.




Figure 2

Crystal packing of the title compound showing intermolecular interactions with dotted lines. H-atoms not involved in hydrogen bonding have been excluded.

Bis{2-[4-(methylsulfanyl)phenyl]-1*H*-benzimidazol-3-ium} tetrabromidocuprate(II) dihydrate

Crystal data

(C₁₄H₁₃N₂S)₂[CuBr₄]·2H₂O

*M*_r = 901.86

Triclinic, *P*1̄

Hall symbol: -P 1

a = 7.6878 (5) Å

b = 11.8358 (7) Å

c = 18.5485 (9) Å

α = 85.305 (4)°

β = 84.778 (5)°

γ = 80.692 (5)°

V = 1654.74 (17) Å³

Z = 2

F(000) = 886

*D*_x = 1.810 Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 5805 reflections
 $\theta = 2.7\text{--}25.0^\circ$
 $\mu = 5.65 \text{ mm}^{-1}$

$T = 296 \text{ K}$
 Block, yellow
 $0.18 \times 0.16 \times 0.16 \text{ mm}$

Data collection

Bruker SMART APEX CCD detector
 diffractometer
 Radiation source: Enhance (Mo) X-ray Source
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 1998)
 $T_{\min} = 0.430$, $T_{\max} = 0.465$

27134 measured reflections
 5805 independent reflections
 3344 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.110$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.7^\circ$
 $h = -9 \rightarrow 9$
 $k = -14 \rightarrow 14$
 $l = -22 \rightarrow 22$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.127$
 $S = 1.00$
 5805 reflections
 384 parameters
 6 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0351P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.75 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.59 \text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br3	0.84824 (12)	0.57905 (7)	0.30208 (4)	0.0669 (3)
Br2	0.87267 (11)	0.78902 (7)	0.15526 (5)	0.0683 (3)
Cu1	0.66741 (12)	0.74774 (8)	0.25235 (5)	0.0514 (3)
Br4	0.53451 (13)	0.82526 (8)	0.35966 (5)	0.0790 (3)
Br1	0.41165 (11)	0.76741 (8)	0.18456 (4)	0.0704 (3)
S1	0.6325 (3)	0.24760 (18)	0.32773 (11)	0.0633 (6)
S2	0.9931 (3)	0.15553 (19)	0.17091 (12)	0.0709 (7)
N1	0.0725 (8)	0.7087 (5)	0.4564 (3)	0.0510 (17)
H1	0.0834	0.7291	0.4108	0.061*
N2	0.1042 (7)	0.6081 (5)	0.5582 (3)	0.0455 (15)
H2	0.1382	0.5532	0.5896	0.055*
N3	0.6882 (8)	0.5966 (5)	-0.0737 (3)	0.0483 (16)

H3	0.7331	0.5473	−0.1045	0.058*
N4	0.6053 (7)	0.6754 (5)	0.0272 (3)	0.0491 (16)
H4	0.5875	0.6850	0.0729	0.059*
O1	0.2089 (9)	0.5361 (5)	0.1865 (3)	0.0744 (18)
O2	0.0818 (8)	0.8287 (6)	0.3236 (3)	0.0755 (17)
H1E	0.263 (9)	0.593 (5)	0.180 (5)	0.113*
H1D	0.103 (5)	0.557 (7)	0.204 (5)	0.113*
H2B	0.188 (5)	0.838 (8)	0.326 (4)	0.113*
H2A	0.070 (10)	0.811 (8)	0.282 (2)	0.113*
C1	0.6852 (11)	0.1254 (6)	0.3895 (4)	0.073 (3)
H1B	0.7592	0.0654	0.3642	0.109*
H1A	0.5782	0.0988	0.4094	0.109*
H1C	0.7466	0.1462	0.4280	0.109*
C2	0.4938 (9)	0.3474 (6)	0.3793 (4)	0.049 (2)
C3	0.4289 (10)	0.4503 (7)	0.3424 (4)	0.059 (2)
H3A	0.4626	0.4628	0.2933	0.071*
C4	0.3167 (9)	0.5329 (6)	0.3772 (4)	0.049 (2)
H4A	0.2709	0.5996	0.3508	0.059*
C5	0.2689 (9)	0.5208 (6)	0.4504 (4)	0.0406 (18)
C6	0.3317 (9)	0.4189 (6)	0.4874 (4)	0.0435 (18)
H6	0.2973	0.4073	0.5364	0.052*
C7	0.4443 (10)	0.3339 (6)	0.4531 (4)	0.050 (2)
H7	0.4878	0.2667	0.4795	0.060*
C8	0.1509 (9)	0.6090 (6)	0.4875 (4)	0.0418 (18)
C9	−0.0276 (9)	0.7731 (6)	0.5082 (4)	0.0480 (19)
C10	−0.0077 (9)	0.7088 (6)	0.5740 (4)	0.0460 (19)
C11	−0.0876 (10)	0.7504 (7)	0.6385 (4)	0.058 (2)
H11	−0.0746	0.7078	0.6826	0.070*
C12	−0.1869 (10)	0.8578 (7)	0.6341 (4)	0.063 (2)
H12	−0.2439	0.8880	0.6763	0.075*
C13	−0.2050 (10)	0.9230 (7)	0.5683 (4)	0.064 (2)
H13	−0.2710	0.9961	0.5678	0.077*
C14	−0.1274 (10)	0.8812 (6)	0.5045 (4)	0.063 (2)
H14	−0.1412	0.9239	0.4605	0.076*
C15	1.0901 (10)	0.0542 (6)	0.1072 (4)	0.077 (3)
H15C	1.1793	0.0857	0.0758	0.116*
H15A	1.0005	0.0372	0.0787	0.116*
H15B	1.1429	−0.0150	0.1326	0.116*
C16	0.9121 (9)	0.2772 (6)	0.1158 (4)	0.050 (2)
C17	0.9176 (10)	0.2808 (6)	0.0417 (4)	0.056 (2)
H17	0.9687	0.2168	0.0171	0.067*
C18	0.8455 (9)	0.3818 (6)	0.0031 (4)	0.050 (2)
H18	0.8514	0.3843	−0.0473	0.060*
C19	0.7664 (9)	0.4771 (6)	0.0380 (4)	0.0432 (18)
C20	0.7666 (10)	0.4719 (6)	0.1131 (4)	0.056 (2)
H20	0.7191	0.5363	0.1380	0.067*
C21	0.8361 (10)	0.3726 (6)	0.1511 (4)	0.061 (2)
H21	0.8315	0.3701	0.2015	0.074*

C22	0.6917 (9)	0.5791 (6)	−0.0009 (4)	0.0455 (19)
C23	0.6011 (9)	0.7062 (6)	−0.0912 (4)	0.0435 (18)
C24	0.5622 (10)	0.7637 (7)	−0.1566 (4)	0.061 (2)
H24	0.5965	0.7297	−0.2002	0.073*
C25	0.4703 (11)	0.8735 (7)	−0.1549 (5)	0.066 (2)
H25	0.4428	0.9147	−0.1984	0.080*
C26	0.4178 (10)	0.9240 (7)	−0.0901 (5)	0.064 (2)
H26	0.3564	0.9985	−0.0911	0.077*
C27	0.4541 (10)	0.8667 (6)	−0.0245 (4)	0.060 (2)
H27	0.4172	0.8999	0.0192	0.071*
C28	0.5485 (10)	0.7572 (6)	−0.0269 (4)	0.050 (2)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br3	0.0774 (6)	0.0605 (6)	0.0514 (5)	0.0114 (5)	0.0023 (4)	0.0136 (4)
Br2	0.0606 (6)	0.0670 (6)	0.0700 (6)	−0.0054 (4)	0.0039 (5)	0.0200 (5)
Cu1	0.0536 (6)	0.0493 (6)	0.0477 (6)	0.0012 (5)	−0.0026 (5)	−0.0023 (4)
Br4	0.0916 (7)	0.0806 (7)	0.0606 (6)	0.0093 (6)	−0.0038 (5)	−0.0273 (5)
Br1	0.0521 (5)	0.1001 (7)	0.0562 (6)	0.0001 (5)	−0.0059 (4)	−0.0116 (5)
S1	0.0654 (15)	0.0619 (14)	0.0568 (13)	0.0100 (12)	−0.0091 (11)	−0.0053 (11)
S2	0.0672 (15)	0.0632 (15)	0.0771 (16)	−0.0009 (12)	−0.0111 (13)	0.0147 (12)
N1	0.058 (4)	0.042 (4)	0.050 (4)	−0.001 (3)	−0.008 (3)	0.002 (3)
N2	0.049 (4)	0.050 (4)	0.037 (4)	−0.007 (3)	−0.009 (3)	0.001 (3)
N3	0.062 (4)	0.042 (4)	0.039 (4)	−0.001 (3)	−0.002 (3)	−0.008 (3)
N4	0.057 (4)	0.042 (4)	0.047 (4)	−0.008 (3)	−0.004 (3)	0.001 (3)
O1	0.078 (5)	0.078 (4)	0.060 (4)	0.008 (3)	0.003 (4)	−0.017 (3)
O2	0.084 (4)	0.089 (5)	0.054 (4)	−0.021 (4)	−0.011 (3)	0.011 (3)
C1	0.085 (7)	0.051 (5)	0.080 (6)	0.000 (5)	−0.018 (5)	0.000 (5)
C2	0.033 (4)	0.054 (5)	0.059 (5)	0.002 (4)	−0.013 (4)	−0.005 (4)
C3	0.053 (5)	0.072 (6)	0.044 (5)	0.005 (4)	−0.002 (4)	0.005 (4)
C4	0.048 (5)	0.052 (5)	0.041 (5)	0.005 (4)	−0.004 (4)	0.004 (4)
C5	0.045 (4)	0.038 (4)	0.039 (4)	−0.008 (3)	−0.006 (3)	0.004 (3)
C6	0.054 (5)	0.047 (4)	0.032 (4)	−0.017 (4)	−0.004 (4)	0.002 (3)
C7	0.059 (5)	0.047 (5)	0.045 (5)	−0.010 (4)	−0.010 (4)	0.008 (4)
C8	0.039 (4)	0.043 (4)	0.044 (5)	−0.008 (3)	−0.007 (4)	0.000 (3)
C9	0.049 (5)	0.051 (5)	0.042 (5)	−0.008 (4)	0.000 (4)	0.002 (4)
C10	0.045 (5)	0.050 (5)	0.045 (5)	−0.009 (4)	−0.004 (4)	−0.009 (4)
C11	0.052 (5)	0.066 (6)	0.054 (5)	−0.004 (4)	−0.004 (4)	0.001 (4)
C12	0.058 (6)	0.063 (6)	0.063 (6)	−0.002 (5)	0.000 (4)	−0.009 (5)
C13	0.062 (6)	0.055 (5)	0.068 (6)	0.010 (4)	0.001 (5)	−0.006 (5)
C14	0.070 (6)	0.048 (5)	0.067 (6)	0.006 (4)	−0.015 (5)	0.002 (4)
C15	0.063 (6)	0.051 (5)	0.112 (7)	−0.001 (4)	0.003 (5)	0.006 (5)
C16	0.047 (5)	0.047 (5)	0.056 (5)	−0.008 (4)	−0.005 (4)	0.003 (4)
C17	0.055 (5)	0.045 (5)	0.065 (6)	−0.002 (4)	0.000 (4)	−0.007 (4)
C18	0.060 (5)	0.052 (5)	0.040 (4)	−0.010 (4)	−0.007 (4)	−0.007 (4)
C19	0.048 (5)	0.033 (4)	0.047 (5)	−0.004 (3)	0.001 (4)	−0.004 (3)
C20	0.073 (6)	0.053 (5)	0.041 (5)	−0.005 (4)	−0.008 (4)	0.002 (4)

C21	0.078 (6)	0.053 (5)	0.048 (5)	−0.005 (5)	0.001 (4)	0.010 (4)
C22	0.049 (5)	0.041 (4)	0.048 (5)	−0.014 (4)	−0.003 (4)	−0.003 (4)
C23	0.039 (4)	0.042 (4)	0.051 (5)	−0.007 (4)	−0.006 (4)	0.000 (4)
C24	0.069 (6)	0.064 (6)	0.052 (5)	−0.021 (5)	−0.011 (4)	0.008 (4)
C25	0.077 (6)	0.061 (6)	0.067 (6)	−0.025 (5)	−0.034 (5)	0.022 (5)
C26	0.063 (6)	0.047 (5)	0.079 (7)	0.000 (4)	−0.020 (5)	0.004 (5)
C27	0.063 (6)	0.047 (5)	0.069 (6)	−0.005 (4)	−0.009 (5)	−0.008 (4)
C28	0.062 (5)	0.046 (5)	0.039 (5)	−0.003 (4)	−0.007 (4)	0.000 (4)

Geometric parameters (Å, °)

Br3—Cu1	2.4091 (11)	C6—H6	0.9300
Br2—Cu1	2.3579 (12)	C7—H7	0.9300
Cu1—Br4	2.3396 (12)	C9—C14	1.382 (9)
Cu1—Br1	2.3990 (12)	C9—C10	1.391 (9)
S1—C2	1.743 (7)	C10—C11	1.385 (9)
S1—C1	1.787 (7)	C11—C12	1.373 (9)
S2—C16	1.758 (7)	C11—H11	0.9300
S2—C15	1.782 (7)	C12—C13	1.395 (10)
N1—C8	1.346 (8)	C12—H12	0.9300
N1—C9	1.376 (8)	C13—C14	1.368 (9)
N1—H1	0.8600	C13—H13	0.9300
N2—C8	1.327 (8)	C14—H14	0.9300
N2—C10	1.388 (8)	C15—H15C	0.9600
N2—H2	0.8600	C15—H15A	0.9600
N3—C22	1.351 (8)	C15—H15B	0.9600
N3—C23	1.388 (8)	C16—C17	1.370 (9)
N3—H3	0.8600	C16—C21	1.373 (9)
N4—C22	1.342 (8)	C17—C18	1.402 (9)
N4—C28	1.386 (8)	C17—H17	0.9300
N4—H4	0.8600	C18—C19	1.373 (8)
O1—H1E	0.84 (7)	C18—H18	0.9300
O1—H1D	0.85 (6)	C19—C20	1.390 (9)
O2—H2B	0.85 (5)	C19—C22	1.421 (9)
O2—H2A	0.84 (2)	C20—C21	1.376 (9)
C1—H1B	0.9600	C20—H20	0.9300
C1—H1A	0.9600	C21—H21	0.9300
C1—H1C	0.9600	C23—C24	1.374 (9)
C2—C7	1.390 (9)	C23—C28	1.379 (9)
C2—C3	1.391 (9)	C24—C25	1.377 (10)
C3—C4	1.358 (9)	C24—H24	0.9300
C3—H3A	0.9300	C25—C26	1.385 (10)
C4—C5	1.376 (9)	C25—H25	0.9300
C4—H4A	0.9300	C26—C27	1.373 (10)
C5—C6	1.377 (8)	C26—H26	0.9300
C5—C8	1.443 (9)	C27—C28	1.381 (9)
C6—C7	1.374 (9)	C27—H27	0.9300

Br4—Cu1—Br2	139.54 (6)	C12—C11—H11	121.7
Br4—Cu1—Br1	99.30 (4)	C10—C11—H11	121.7
Br2—Cu1—Br1	97.76 (4)	C11—C12—C13	122.0 (7)
Br4—Cu1—Br3	99.91 (4)	C11—C12—H12	119.0
Br2—Cu1—Br3	96.13 (4)	C13—C12—H12	119.0
Br1—Cu1—Br3	130.74 (5)	C14—C13—C12	121.2 (7)
C2—S1—C1	104.8 (4)	C14—C13—H13	119.4
C16—S2—C15	103.5 (4)	C12—C13—H13	119.4
C8—N1—C9	109.9 (6)	C13—C14—C9	117.2 (7)
C8—N1—H1	125.0	C13—C14—H14	121.4
C9—N1—H1	125.0	C9—C14—H14	121.4
C8—N2—C10	109.9 (6)	S2—C15—H15C	109.5
C8—N2—H2	125.0	S2—C15—H15A	109.5
C10—N2—H2	125.0	H15C—C15—H15A	109.5
C22—N3—C23	109.6 (6)	S2—C15—H15B	109.5
C22—N3—H3	125.2	H15C—C15—H15B	109.5
C23—N3—H3	125.2	H15A—C15—H15B	109.5
C22—N4—C28	111.2 (6)	C17—C16—C21	119.4 (7)
C22—N4—H4	124.4	C17—C16—S2	124.3 (6)
C28—N4—H4	124.4	C21—C16—S2	116.3 (6)
H1E—O1—H1D	110 (5)	C16—C17—C18	119.5 (7)
H2B—O2—H2A	109 (5)	C16—C17—H17	120.2
S1—C1—H1B	109.5	C18—C17—H17	120.2
S1—C1—H1A	109.5	C19—C18—C17	121.6 (7)
H1B—C1—H1A	109.5	C19—C18—H18	119.2
S1—C1—H1C	109.5	C17—C18—H18	119.2
H1B—C1—H1C	109.5	C18—C19—C20	117.7 (6)
H1A—C1—H1C	109.5	C18—C19—C22	121.6 (7)
C7—C2—C3	117.7 (6)	C20—C19—C22	120.7 (6)
C7—C2—S1	126.4 (6)	C21—C20—C19	120.8 (7)
C3—C2—S1	116.0 (6)	C21—C20—H20	119.6
C4—C3—C2	120.7 (7)	C19—C20—H20	119.6
C4—C3—H3A	119.7	C16—C21—C20	121.0 (7)
C2—C3—H3A	119.7	C16—C21—H21	119.5
C3—C4—C5	121.9 (6)	C20—C21—H21	119.5
C3—C4—H4A	119.0	N4—C22—N3	106.5 (6)
C5—C4—H4A	119.0	N4—C22—C19	126.9 (7)
C4—C5—C6	117.7 (6)	N3—C22—C19	126.5 (6)
C4—C5—C8	121.9 (6)	C24—C23—C28	120.8 (7)
C6—C5—C8	120.3 (6)	C24—C23—N3	132.0 (7)
C7—C6—C5	121.2 (6)	C28—C23—N3	107.2 (6)
C7—C6—H6	119.4	C23—C24—C25	117.3 (7)
C5—C6—H6	119.4	C23—C24—H24	121.4
C6—C7—C2	120.7 (6)	C25—C24—H24	121.4
C6—C7—H7	119.7	C24—C25—C26	121.6 (7)
C2—C7—H7	119.7	C24—C25—H25	119.2
N2—C8—N1	107.8 (6)	C26—C25—H25	119.2
N2—C8—C5	126.5 (6)	C27—C26—C25	121.5 (7)

N1—C8—C5	125.7 (6)	C27—C26—H26	119.2
N1—C9—C14	132.3 (7)	C25—C26—H26	119.2
N1—C9—C10	106.2 (6)	C26—C27—C28	116.4 (7)
C14—C9—C10	121.5 (7)	C26—C27—H27	121.8
C11—C10—N2	132.5 (7)	C28—C27—H27	121.8
C11—C10—C9	121.3 (7)	C23—C28—C27	122.4 (7)
N2—C10—C9	106.1 (6)	C23—C28—N4	105.4 (6)
C12—C11—C10	116.7 (7)	C27—C28—N4	132.1 (7)
C1—S1—C2—C7	2.6 (8)	C15—S2—C16—C17	4.3 (8)
C1—S1—C2—C3	−178.0 (6)	C15—S2—C16—C21	−177.1 (6)
C7—C2—C3—C4	−2.0 (12)	C21—C16—C17—C18	0.0 (12)
S1—C2—C3—C4	178.6 (6)	S2—C16—C17—C18	178.6 (6)
C2—C3—C4—C5	2.9 (12)	C16—C17—C18—C19	−1.1 (12)
C3—C4—C5—C6	−3.2 (11)	C17—C18—C19—C20	2.6 (12)
C3—C4—C5—C8	179.4 (7)	C17—C18—C19—C22	−179.4 (7)
C4—C5—C6—C7	2.7 (11)	C18—C19—C20—C21	−3.1 (12)
C8—C5—C6—C7	−179.9 (7)	C22—C19—C20—C21	178.9 (8)
C5—C6—C7—C2	−1.9 (11)	C17—C16—C21—C20	−0.5 (13)
C3—C2—C7—C6	1.5 (11)	S2—C16—C21—C20	−179.2 (7)
S1—C2—C7—C6	−179.1 (6)	C19—C20—C21—C16	2.1 (13)
C10—N2—C8—N1	0.3 (8)	C28—N4—C22—N3	−1.4 (8)
C10—N2—C8—C5	178.6 (7)	C28—N4—C22—C19	−179.2 (7)
C9—N1—C8—N2	−0.3 (8)	C23—N3—C22—N4	1.1 (8)
C9—N1—C8—C5	−178.6 (7)	C23—N3—C22—C19	178.9 (7)
C4—C5—C8—N2	−176.3 (7)	C18—C19—C22—N4	177.1 (7)
C6—C5—C8—N2	6.4 (11)	C20—C19—C22—N4	−4.9 (12)
C4—C5—C8—N1	1.7 (12)	C18—C19—C22—N3	−0.2 (12)
C6—C5—C8—N1	−175.6 (7)	C20—C19—C22—N3	177.7 (7)
C8—N1—C9—C14	177.7 (8)	C22—N3—C23—C24	−179.4 (8)
C8—N1—C9—C10	0.2 (8)	C22—N3—C23—C28	−0.4 (8)
C8—N2—C10—C11	−177.9 (8)	C28—C23—C24—C25	0.4 (12)
C8—N2—C10—C9	−0.1 (8)	N3—C23—C24—C25	179.2 (8)
N1—C9—C10—C11	178.0 (7)	C23—C24—C25—C26	−0.5 (13)
C14—C9—C10—C11	0.2 (12)	C24—C25—C26—C27	−0.4 (14)
N1—C9—C10—N2	−0.1 (8)	C25—C26—C27—C28	1.2 (13)
C14—C9—C10—N2	−177.9 (7)	C24—C23—C28—C27	0.5 (12)
N2—C10—C11—C12	177.5 (8)	N3—C23—C28—C27	−178.6 (7)
C9—C10—C11—C12	0.1 (12)	C24—C23—C28—N4	178.7 (7)
C10—C11—C12—C13	−0.9 (12)	N3—C23—C28—N4	−0.4 (8)
C11—C12—C13—C14	1.6 (14)	C26—C27—C28—C23	−1.3 (12)
C12—C13—C14—C9	−1.3 (13)	C26—C27—C28—N4	−178.9 (8)
N1—C9—C14—C13	−176.7 (8)	C22—N4—C28—C23	1.1 (8)
C10—C9—C14—C13	0.4 (12)	C22—N4—C28—C27	179.1 (8)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C13—H13 $\cdots$ Br4 ⁱ	0.93	2.99	3.831 (8)	150
N3—H3 $\cdots$ O1 ⁱⁱ	0.86	1.86	2.703 (8)	165
C18—H18 $\cdots$ O1 ⁱⁱ	0.93	2.73	3.614 (9)	159
N2—H2 $\cdots$ Br3 ⁱⁱⁱ	0.86	2.44	3.275 (6)	162
O1—H1 <i>D</i> $\cdots$ Br3 ^{iv}	0.85 (6)	2.55 (7)	3.344 (6)	155
O2—H2 <i>A</i> $\cdots$ Br2 ^{iv}	0.83 (4)	2.96 (6)	3.735 (6)	155
O1—H1 <i>E</i> $\cdots$ Br1	0.84 (7)	2.53 (7)	3.359 (6)	170
O2—H2 <i>B</i> $\cdots$ Br4	0.85 (5)	2.77 (7)	3.597 (6)	166

Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $-x+1, -y+1, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x-1, y, z$.