

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dicyanido[tris(2-pyridylmethyl)amine]-cobalt(III) hexafluoridophosphate

Fan Yu^{a*} and Bao Li^b

^aSchool of Chemistry and Environmental Engineering, Jiangnan University, Wuhan, Hubei 430056, People's Republic of China, and ^bDepartment of Chemistry and Chemical Engineering, Huazhong University of Science and Technology, Wuhan 430074, People's Republic of China
Correspondence e-mail: yufan0714@163.com


Received 3 March 2011; accepted 31 March 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.025; wR factor = 0.070; data-to-parameter ratio = 12.3.

In the title complex, $[\text{Co}(\text{CN})_2(\text{C}_{18}\text{H}_{18}\text{N}_4)]\text{PF}_6$, the Co^{III} atom together with one of the pyridyl rings and two cyanide anions are located on a mirror plane, while the P atom is located on an inversion centre. The Co^{III} atom exhibits an octahedral geometry, coordinated by four N atoms from the tris(2-pyridylmethyl)amine ligand with an average $\text{Co}-\text{N}$ distance of 1.953 (2) Å, and two cyanide C atoms with an average $\text{Co}-\text{C}$ distance of 1.895 (2) Å. The crystal packing is stabilized by intermolecular $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{F}$ interactions.

Related literature

For related structures, see: Guo *et al.* (2007), Liu *et al.* (2010).



Experimental

Crystal data

 $[\text{Co}(\text{CN})_2(\text{C}_{18}\text{H}_{18}\text{N}_4)]\text{PF}_6$ $M_r = 546.30$ Orthorhombic, $Pbcm$

$a = 10.703$ (2) Å
 $b = 13.472$ (3) Å
 $c = 15.151$ (3) Å
 $V = 2184.7$ (8) Å³

 $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.93$ mm⁻¹ $T = 293$ K $0.40 \times 0.30 \times 0.25$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.722$, $T_{\max} = 0.792$

26914 measured reflections
 2173 independent reflections
 2053 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.070$
 $S = 1.08$
 2173 reflections

176 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.27$ e Å⁻³
 $\Delta\rho_{\min} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3A}\cdots\text{N4}^{\text{i}}$	0.93	2.60	3.339 (3)	137
$\text{C6}-\text{H6A}\cdots\text{F3}^{\text{i}}$	0.96	2.29	3.234 (2)	169
$\text{C7}-\text{H7A}\cdots\text{F2}^{\text{ii}}$	0.93	2.44	3.128 (3)	131
$\text{C9}-\text{H9A}\cdots\text{N5}^{\text{iii}}$	0.93	2.57	3.410 (2)	151

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y, -z + \frac{1}{2}$; (iii) $-x + 2, -y, z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Jiangnan University and Huazhong University of Science and Technology are thanked for their generous financial support and for a start-up grant.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2083).

References

- Bruker (1997). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (1999). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Guo, Y., Feng, Y. H., Liu, Z. Q. & Liao, D. Z. (2007). *J. Coord. Chem.* **60**, 2713–2720.
 Liu, T., Zhang, Y. J., Kanegawa, S. & Sato, O. (2010). *J. Am. Chem. Soc.* **132**, 8250–8251.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2011). E67, m566 [https://doi.org/10.1107/S1600536811012001]

Dicyanido[tris(2-pyridylmethyl)amine]cobalt(III) hexafluoridophosphate**Fan Yu and Bao Li****S1. Comment**

Transitional metal-cyanide systems have been extensively investigated due to their versatile structure and physical properties, especially in the field of molecular-based magnets. A large number of cyanide-bridged heterobimetallic or homometallic coordination complexes that exhibit excellent magnetic properties has been constructed by using the hexacyanoferrate(III) and hexacyanocobaltate(III) anions as templates (Liu *et al.*, 2010). In contrast, complexes constructed by using the octacyanotungsten(IV) anion are rare. In an attempt to synthesize new complexes, we decided to use octacyanotungsten(IV) anions as template and new compounds containing cyanides have been obtained. The octacyanotungsten(IV) anion was not coordinated to Co atom *via* cyanide bridges, but acts as source of *in situ* cyanide generation.

In the title compound, the cobalt(III) centers are coordinated by two cyanides and tris(2-pyridylmethyl)amine (Fig. 1). Each cobalt(III) ion is coordinated by four N atoms with average Co—N distance of 1.957 (2) Å and two C atoms with average Co—C distance of 1.895 (2) Å in a rigid octahedral geometry, in accordance with those observed in other [Co(N)₄(CN)₂][−] units (Guo *et al.*, 2007). The dihedral angle of two types of pyridyl rings is about 80.17°, indicating the nearly perpendicular occupation of these pyridyl rings. The crystal packing is stabilized by C—H⋯N and C—H⋯F hydrogen bonding interactions (Table 1, Fig. 2).

S2. Experimental

0.01 mmol K₄W(CN)₈ in 5 ml H₂O was added in a tube, and 3 ml H₂O was layered on as a buffer. A solution containing 0.1 mmol CoCl₂·6H₂O, 0.11 mmol tris(2-pyridylmethyl)amine and 0.11 mmol KPF₆ in 5 mL acetone and 1 ml H₂O was stirred for 30 min and layered on top of the previous solution. After half a year, yellow crystals were obtained.

S3. Refinement

All H atoms were placed geometrically with C—H = 0.93 (aromatic) or 0.96–0.97 Å (CH₂), and refined using a riding atom model with their isotropic displacement factors, *U*_{iso} fixed at 1.2 times the *U*_{eq} of the parent C atom.




Figure 1

Molecular structure of the title compound showing atomic numbering and 30% probability displacement ellipsoids. Symmetry codes: (i) $x, y, -z + 1/2$; (ii) $x, -y + 1/2, -z$.




Figure 2

The packing of title compound, viewed down the b axis. Symmetry codes: (i) $x, y, -z + 1/2$; (ii) $x, -y + 1/2, -z$.

Dicyanido[tris(2-pyridylmethyl)amine]cobalt(III) hexafluoridophosphate

Crystal data

[Co(CN)₂(C₁₈H₁₈N₄)]PF₆ $M_r = 546.30$ Orthorhombic, *Pbcm*

Hall symbol: -P 2c 2b

 $a = 10.703 (2) \text{ \AA}$ $b = 13.472 (3) \text{ \AA}$ $c = 15.151 (3) \text{ \AA}$ $V = 2184.7 (8) \text{ \AA}^3$ $Z = 4$ $F(000) = 1104$ $D_x = 1.658 \text{ Mg m}^{-3}$ $D_m = 1.658 \text{ Mg m}^{-3}$ D_m measured by not measuredMo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 27644 reflections

 $\theta = 6.1\text{--}54.9^\circ$ $\mu = 0.93 \text{ mm}^{-1}$ $T = 293 \text{ K}$

Block, yellow

 $0.40 \times 0.30 \times 0.25 \text{ mm}$

Data collection

Bruker SMART APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 0 pixels mm^{-1} ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.722$, $T_{\max} = 0.792$

26914 measured reflections

2173 independent reflections

2053 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.028$ $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 3.3^\circ$ $h = -13 \rightarrow 13$ $k = -16 \rightarrow 14$ $l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.025$ $wR(F^2) = 0.070$ $S = 1.08$

2173 reflections

176 parameters

0 restraints

1.234 constraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.043P)^2 + 0.5717P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.27 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.82856 (2)	0.057628 (17)	0.2500	0.02656 (10)
N1	0.64937 (15)	0.02901 (13)	0.2500	0.0308 (3)
N2	0.82831 (10)	0.04135 (9)	0.37718 (8)	0.0326 (3)

N3	0.85307 (16)	−0.08775 (13)	0.2500	0.0328 (4)
N4	0.8016 (2)	0.28241 (14)	0.2500	0.0499 (5)
N5	1.10402 (19)	0.11315 (17)	0.2500	0.0508 (5)
C1	0.5562 (2)	0.09626 (16)	0.2500	0.0412 (5)
H1A	0.5761	0.1634	0.2500	0.049*
C2	0.4321 (2)	0.0686 (2)	0.2500	0.0501 (6)
H2A	0.3694	0.1163	0.2500	0.060*
C3	0.4029 (2)	−0.0304 (2)	0.2500	0.0503 (6)
H3A	0.3199	−0.0507	0.2500	0.060*
C4	0.4976 (2)	−0.09968 (17)	0.2500	0.0470 (5)
H4A	0.4792	−0.1671	0.2500	0.056*
P1	0.43435 (6)	0.2500	0.0000	0.04841 (17)
C6	0.7279 (2)	−0.13885 (15)	0.2500	0.0408 (5)
H6A	0.7219	−0.1806	0.3012	0.049*
F2	0.54077 (17)	0.23947 (14)	0.07289 (13)	0.1109 (6)
F1	0.43287 (15)	0.13526 (10)	−0.01878 (13)	0.0969 (5)
C13	0.80941 (18)	0.19751 (15)	0.2500	0.0333 (4)
C8	0.76782 (18)	0.07851 (15)	0.52469 (11)	0.0524 (4)
H8A	0.7322	0.1225	0.5647	0.063*
F3	0.32765 (16)	0.23608 (13)	0.07209 (10)	0.0978 (5)
C10	0.86385 (17)	−0.07628 (14)	0.49189 (11)	0.0447 (4)
H10A	0.8939	−0.1378	0.5098	0.054*
C7	0.77754 (15)	0.10373 (12)	0.43633 (10)	0.0417 (3)
H7A	0.7483	0.1652	0.4175	0.050*
C12	0.92406 (15)	−0.11166 (11)	0.33220 (10)	0.0400 (3)
H12A	1.0122	−0.0977	0.3241	0.048*
H12B	0.9146	−0.1813	0.3469	0.048*
C9	0.81121 (17)	−0.01197 (16)	0.55256 (11)	0.0510 (4)
H9A	0.8053	−0.0299	0.6117	0.061*
C14	1.0012 (2)	0.08893 (15)	0.2500	0.0340 (4)
C11	0.87139 (14)	−0.04806 (10)	0.40424 (10)	0.0357 (3)
C5	0.6211 (2)	−0.06763 (14)	0.2500	0.0336 (4)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02638 (16)	0.02409 (15)	0.02921 (15)	0.00029 (9)	0.000	0.000
N1	0.0296 (8)	0.0301 (8)	0.0326 (8)	−0.0016 (6)	0.000	0.000
N2	0.0312 (6)	0.0339 (6)	0.0328 (6)	−0.0008 (4)	−0.0014 (4)	−0.0005 (5)
N3	0.0355 (9)	0.0281 (8)	0.0348 (8)	0.0041 (7)	0.000	0.000
N4	0.0456 (11)	0.0296 (9)	0.0746 (14)	−0.0006 (8)	0.000	0.000
N5	0.0318 (10)	0.0664 (13)	0.0541 (11)	−0.0040 (9)	0.000	0.000
C1	0.0311 (10)	0.0343 (10)	0.0583 (13)	0.0002 (8)	0.000	0.000
C2	0.0318 (12)	0.0533 (14)	0.0654 (16)	0.0017 (9)	0.000	0.000
C3	0.0301 (11)	0.0612 (14)	0.0595 (14)	−0.0113 (10)	0.000	0.000
C4	0.0449 (13)	0.0394 (11)	0.0566 (13)	−0.0155 (10)	0.000	0.000
P1	0.0461 (3)	0.0473 (3)	0.0519 (3)	0.000	0.000	−0.0053 (3)
C6	0.0452 (12)	0.0269 (9)	0.0505 (12)	−0.0053 (8)	0.000	0.000

F2	0.0979 (12)	0.1154 (13)	0.1194 (13)	0.0305 (10)	−0.0483 (10)	0.0004 (10)
F1	0.0994 (11)	0.0502 (7)	0.1411 (14)	−0.0045 (7)	0.0406 (10)	−0.0180 (8)
C13	0.0278 (9)	0.0317 (10)	0.0403 (10)	−0.0021 (7)	0.000	0.000
C8	0.0529 (10)	0.0681 (11)	0.0362 (8)	0.0019 (9)	0.0042 (7)	−0.0091 (8)
F3	0.0994 (12)	0.1130 (13)	0.0809 (10)	−0.0082 (9)	0.0396 (8)	−0.0240 (9)
C10	0.0443 (8)	0.0509 (9)	0.0389 (8)	−0.0040 (7)	−0.0058 (7)	0.0106 (7)
C7	0.0447 (9)	0.0440 (8)	0.0363 (7)	0.0040 (6)	0.0014 (6)	−0.0046 (6)
C12	0.0448 (8)	0.0345 (7)	0.0409 (8)	0.0085 (6)	−0.0051 (6)	0.0058 (6)
C9	0.0497 (9)	0.0721 (12)	0.0314 (7)	−0.0062 (8)	0.0000 (7)	0.0057 (8)
C14	0.0330 (11)	0.0348 (9)	0.0341 (9)	0.0025 (8)	0.000	0.000
C11	0.0330 (7)	0.0379 (7)	0.0361 (7)	−0.0021 (6)	−0.0041 (6)	0.0044 (6)
C5	0.0383 (11)	0.0328 (10)	0.0296 (9)	−0.0057 (8)	0.000	0.000

Geometric parameters (Å, °)

Co1—C14	1.895 (2)	C4—C5	1.390 (3)
Co1—C13	1.896 (2)	C4—H4A	0.9300
Co1—N2 ⁱ	1.9393 (13)	P1—F1 ⁱⁱ	1.5719 (14)
Co1—N2	1.9393 (13)	P1—F1	1.5719 (14)
Co1—N1	1.9563 (17)	P1—F3 ⁱⁱ	1.5913 (15)
Co1—N3	1.9760 (18)	P1—F3	1.5913 (15)
N1—C5	1.337 (3)	P1—F2	1.5928 (16)
N1—C1	1.347 (3)	P1—F2 ⁱⁱ	1.5928 (16)
N2—C7	1.343 (2)	C6—C5	1.492 (3)
N2—C11	1.3534 (19)	C6—H6A	0.9600
N3—C12	1.4940 (17)	C8—C9	1.371 (3)
N3—C12 ⁱ	1.4940 (17)	C8—C7	1.385 (2)
N3—C6	1.507 (3)	C8—H8A	0.9300
N4—C13	1.147 (3)	C10—C9	1.383 (3)
N5—C14	1.148 (3)	C10—C11	1.384 (2)
C1—C2	1.379 (3)	C10—H10A	0.9300
C1—H1A	0.9300	C7—H7A	0.9300
C2—C3	1.370 (4)	C12—C11	1.498 (2)
C2—H2A	0.9300	C12—H12A	0.9700
C3—C4	1.378 (4)	C12—H12B	0.9700
C3—H3A	0.9300	C9—H9A	0.9300
C14—Co1—C13	83.35 (8)	F1 ⁱⁱ —P1—F3	89.11 (10)
C14—Co1—N2 ⁱ	91.52 (3)	F1—P1—F3	90.06 (9)
C13—Co1—N2 ⁱ	96.45 (4)	F3 ⁱⁱ —P1—F3	88.28 (14)
C14—Co1—N2	91.52 (3)	F1 ⁱⁱ —P1—F2	88.24 (9)
C13—Co1—N2	96.45 (4)	F1—P1—F2	92.58 (10)
N2 ⁱ —Co1—N2	167.01 (7)	F3 ⁱⁱ —P1—F2	178.30 (9)
C14—Co1—N1	178.51 (8)	F3—P1—F2	91.54 (11)
C13—Co1—N1	95.16 (8)	F1 ⁱⁱ —P1—F2 ⁱⁱ	92.58 (10)
N2 ⁱ —Co1—N1	88.64 (3)	F1—P1—F2 ⁱⁱ	88.24 (9)
N2—Co1—N1	88.64 (3)	F3 ⁱⁱ —P1—F2 ⁱⁱ	91.54 (11)
C14—Co1—N3	95.23 (8)	F3—P1—F2 ⁱⁱ	178.30 (9)

C13—Co1—N3	178.58 (8)	F2—P1—F2 ⁱⁱ	88.69 (16)
N2 ⁱ —Co1—N3	83.58 (4)	C5—C6—N3	112.79 (17)
N2—Co1—N3	83.58 (4)	C5—C6—H6A	109.0
N1—Co1—N3	86.26 (7)	N3—C6—H6A	109.0
C5—N1—C1	119.17 (18)	N4—C13—Co1	177.95 (19)
C5—N1—Co1	114.45 (14)	C9—C8—C7	119.36 (16)
C1—N1—Co1	126.38 (15)	C9—C8—H8A	120.3
C7—N2—C11	119.50 (13)	C7—C8—H8A	120.3
C7—N2—Co1	126.35 (11)	C9—C10—C11	119.30 (16)
C11—N2—Co1	113.65 (10)	C9—C10—H10A	120.3
C12—N3—C12 ⁱ	112.94 (16)	C11—C10—H10A	120.3
C12—N3—C6	110.72 (10)	N2—C7—C8	121.45 (16)
C12 ⁱ —N3—C6	110.72 (10)	N2—C7—H7A	119.3
C12—N3—Co1	106.33 (9)	C8—C7—H7A	119.3
C12 ⁱ —N3—Co1	106.33 (9)	N3—C12—C11	107.02 (12)
C6—N3—Co1	109.56 (12)	N3—C12—H12A	110.3
N1—C1—C2	122.0 (2)	C11—C12—H12A	110.3
N1—C1—H1A	119.0	N3—C12—H12B	110.3
C2—C1—H1A	119.0	C11—C12—H12B	110.3
C3—C2—C1	118.9 (2)	H12A—C12—H12B	108.6
C3—C2—H2A	120.6	C8—C9—C10	119.36 (15)
C1—C2—H2A	120.6	C8—C9—H9A	120.3
C2—C3—C4	119.4 (2)	C10—C9—H9A	120.3
C2—C3—H3A	120.3	N5—C14—Co1	176.3 (2)
C4—C3—H3A	120.3	N2—C11—C10	121.03 (15)
C3—C4—C5	119.3 (2)	N2—C11—C12	114.62 (13)
C3—C4—H4A	120.4	C10—C11—C12	124.34 (14)
C5—C4—H4A	120.4	N1—C5—C4	121.2 (2)
F1 ⁱⁱ —P1—F1	178.85 (13)	N1—C5—C6	116.95 (18)
F1 ⁱⁱ —P1—F3 ⁱⁱ	90.06 (9)	C4—C5—C6	121.88 (19)
F1—P1—F3 ⁱⁱ	89.11 (10)		
C14—Co1—N1—C5	180.0	C2—C3—C4—C5	0.0
C13—Co1—N1—C5	180.0	C12—N3—C6—C5	116.97 (11)
N2 ⁱ —Co1—N1—C5	83.65 (4)	C12 ⁱ —N3—C6—C5	−116.97 (11)
N2—Co1—N1—C5	−83.65 (4)	Co1—N3—C6—C5	0.0
N3—Co1—N1—C5	0.0	C14—Co1—C13—N4	0.0
C14—Co1—N1—C1	0.0	N2 ⁱ —Co1—C13—N4	−90.79 (3)
C13—Co1—N1—C1	0.0	N2—Co1—C13—N4	90.79 (3)
N2 ⁱ —Co1—N1—C1	−96.35 (4)	N1—Co1—C13—N4	180.0
N2—Co1—N1—C1	96.35 (4)	N3—Co1—C13—N4	0.0
N3—Co1—N1—C1	180.0	C11—N2—C7—C8	−0.2 (2)
C14—Co1—N2—C7	107.30 (14)	Co1—N2—C7—C8	171.16 (13)
C13—Co1—N2—C7	23.82 (14)	C9—C8—C7—N2	0.2 (3)
N2 ⁱ —Co1—N2—C7	−149.2 (2)	C12 ⁱ —N3—C12—C11	156.60 (11)
N1—Co1—N2—C7	−71.22 (14)	C6—N3—C12—C11	−78.58 (16)
N3—Co1—N2—C7	−157.61 (14)	Co1—N3—C12—C11	40.35 (14)
C14—Co1—N2—C11	−80.90 (11)	C7—C8—C9—C10	0.0 (3)

C13—Co1—N2—C11	−164.38 (11)	C11—C10—C9—C8	−0.1 (3)
N2 ⁱ —Co1—N2—C11	22.6 (4)	C13—Co1—C14—N5	0.0
N1—Co1—N2—C11	100.58 (11)	N2 ⁱ —Co1—C14—N5	96.31 (4)
N3—Co1—N2—C11	14.19 (11)	N2—Co1—C14—N5	−96.31 (4)
C14—Co1—N3—C12	60.30 (10)	N1—Co1—C14—N5	0.0
C13—Co1—N3—C12	60.30 (10)	N3—Co1—C14—N5	180.0
N2 ⁱ —Co1—N3—C12	151.25 (11)	C7—N2—C11—C10	0.1 (2)
N2—Co1—N3—C12	−30.65 (10)	Co1—N2—C11—C10	−172.35 (12)
N1—Co1—N3—C12	−119.70 (10)	C7—N2—C11—C12	179.11 (14)
C14—Co1—N3—C12 ⁱ	−60.30 (10)	Co1—N2—C11—C12	6.70 (16)
C13—Co1—N3—C12 ⁱ	−60.30 (10)	C9—C10—C11—N2	0.1 (2)
N2 ⁱ —Co1—N3—C12 ⁱ	30.65 (10)	C9—C10—C11—C12	−178.84 (16)
N2—Co1—N3—C12 ⁱ	−151.25 (11)	N3—C12—C11—N2	−31.69 (18)
N1—Co1—N3—C12 ⁱ	119.70 (10)	N3—C12—C11—C10	147.32 (16)
C14—Co1—N3—C6	180.0	C1—N1—C5—C4	0.0
C13—Co1—N3—C6	180.0	Co1—N1—C5—C4	180.0
N2 ⁱ —Co1—N3—C6	−89.05 (3)	C1—N1—C5—C6	180.0
N2—Co1—N3—C6	89.05 (3)	Co1—N1—C5—C6	0.0
N1—Co1—N3—C6	0.0	C3—C4—C5—N1	0.0
C5—N1—C1—C2	0.0	C3—C4—C5—C6	180.0
Co1—N1—C1—C2	180.0	N3—C6—C5—N1	0.0
N1—C1—C2—C3	0.0	N3—C6—C5—C4	180.0
C1—C2—C3—C4	0.0		

Symmetry codes: (i) $x, y, -z+1/2$; (ii) $x, -y+1/2, -z$.

Hydrogen-bond geometry ($\text{\AA}$, $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3A $\cdots$ N4 ⁱⁱⁱ	0.93	2.60	3.339 (3)	137
C6—H6A $\cdots$ F3 ⁱⁱⁱ	0.96	2.29	3.234 (2)	169
C7—H7A $\cdots$ F2 ⁱ	0.93	2.44	3.128 (3)	131
C9—H9A $\cdots$ N5 ^{iv}	0.93	2.57	3.410 (2)	151

Symmetry codes: (i) $x, y, -z+1/2$; (iii) $-x+1, y-1/2, -z+1/2$; (iv) $-x+2, -y, z+1/2$.