

Bis[*N*-(2-furylmethyl)ethane-1,2-diamine]bis(perchlorato)copper(II)

Wei Xiao,^a Shi-Rong Li,^b Hong Zhou,^a Zhi-Quan Pan^{a*} and Qimao Huang^a

^aKey Laboratory for Green Chemical Processes of the Ministry of Education, Wuhan Institute of Technology, Wuhan, 430073, People's Republic of China, and ^bHubei Key Laboratory of Biologic Resources Protection and Utilization, Hubei Institute for Nationalities, Enshi, 44500, People's Republic of China
Correspondence e-mail: zhiqpan@163.com


Received 23 March 2011; accepted 1 April 2011

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.049; wR factor = 0.112; data-to-parameter ratio = 13.5.

In the title complex, $[\text{Cu}(\text{ClO}_4)_2(\text{C}_7\text{H}_{12}\text{N}_2\text{O})_2]$, the Cu(II) ion lies on a crystallographic inversion centre. The coordination sphere around Cu(II) ion can be described as tetragonally distorted octahedral with two perchlorate O atoms occupying the apical positions and four N atoms from two *N*¹-(2-furylmethyl)ethane-1,2-diamine ligands in the basal plane.

Related literature

For copper complexes with polyamine ligands, see: Souza *et al.* (2009); Patra *et al.* (2007); Zhou *et al.* (2009). For the synthesis, see: Wang *et al.* (2009).



Experimental

Crystal data

$[\text{Cu}(\text{ClO}_4)_2(\text{C}_7\text{H}_{12}\text{N}_2\text{O})_2]$
 $M_r = 542.81$
Monoclinic, $P2_1/c$
 $a = 9.736$ (8) Å
 $b = 11.899$ (9) Å
 $c = 9.466$ (7) Å
 $\beta = 94.227$ (12)°

$V = 1093.6$ (14) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.30$ mm⁻¹
 $T = 291$ K
 $0.28 \times 0.24 \times 0.22$ mm

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.712$, $T_{\max} = 0.763$

5510 measured reflections
1914 independent reflections
1589 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.112$
 $S = 1.07$
1914 reflections

142 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.26$ e Å⁻³
 $\Delta\rho_{\min} = -0.70$ e Å⁻³

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors would like to thank the National Natural Science Foundation of China (20871097 and 20971102) and the Foundation of the Excellent Middle-Young Innovation Group of the Education Department of Hubei Province, China (T200802).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2094).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Patra, A. K., Nethaji, M. & Chakravarty, A. R. (2007). *J. Inorg. Biochem.* **101**, 233–244.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Souza, B., Bortoluzzi, A. J., Bortolotto, T., Fischer, F. L., Terenzi, H., Ferreira, D. E. C., Rocha, W. R. & Neves, A. (2009). *Dalton Trans.* pp. 2027–2035.
- Wang, K. W., Qian, X. H. & Cui, J. N. (2009). *Tetrahedron*, **50**, 10377–10382.
- Zhou, H., Chen, L., Chen, R., Peng, Z. H., Song, Y., Pan, Z. Q., Huang, Q. M., Hu, X. L. & Bai, Z. W. (2009). *CrystEngComm*, **11**, 671–679.

supporting information

Acta Cryst. (2011). E67, m701 [https://doi.org/10.1107/S1600536811012232]

Bis[N-(2-furylmethyl)ethane-1,2-diamine]bis(perchlorato)copper(II)

Wei Xiao, Shi-Rong Li, Hong Zhou, Zhi-Quan Pan and Qimao Huang

S1. Comment

Recently, study of copper complex with polyamine has been given considerable attention (Souza *et al.*, 2009; Patra *et al.*, 2007; Zhou *et al.*, 2009). In this paper, we report on the synthesis and the crystal structure determination of the title complex obtained by the reaction of $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ with the polyamine ligand N^1 -(furan-2-ylmethyl)ethane-1,2-diamine.

In the title complex, $[\text{Cu}(\text{C}_{14}\text{H}_{24}\text{N}_4\text{O}_2)_2](\text{ClO}_4)_2$, the Cu(II) ion lies on a crystallographic inversion centre. The coordination sphere around Cu(II) ion can be best described as slightly distorted octahedral. The basal plane is composed of four nitrogen atoms from the two polyamine ligands with the Cu-N distances of 2.001 (4) and 2.049 (4) Å. The apical positions are occupied by two oxygen atoms from two perchlorate anions with a Cu-O distance of 2.492 (4) Å.

S2. Experimental

N^1 -(furan-2-ylmethyl)ethane-1,2-diamine (L) was prepared according to the literature method (Wang *et al.*, 2009). $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (0.25 mmol, 0.093 g) dissolved in 10 ml H_2O was added dropwise to a solution of L (0.5 mmol, 0.071 g) in 10 ml H_2O . The mixture was stirred at ambient temperature for about 12 h and filtrated. The light blue crystals suitable for X-ray diffraction were obtained by the slow evaporation of the mother solution at ambient temperature for 3 weeks.

S3. Refinement

All H atoms for C-H distances were placed in calculated positions and included in the refinement in the riding-model approximation, with $U(\text{H})$ set to $-1.2U_{\text{eq}}$ of the parent atom.




Figure 1

A view of the title complex cation, with displacement ellipsoids at the 30% probability level. H atoms are excluded for clarity. Unlabelled atoms are related to labelled atoms by inversion symmetry.

Bis[*N*-(2-furylmethyl)ethane-1,2-diamine]bis(perchlorato)copper (II)

Crystal data

[Cu(ClO₄)₂(C₇H₁₂N₂O)₂]

M_r = 542.81

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 9.736 (8) Å

b = 11.899 (9) Å

c = 9.466 (7) Å

β = 94.227 (12)°

V = 1093.6 (14) Å³

Z = 2

F(000) = 558

D_x = 1.648 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 1812 reflections

θ = 2.7–22.6°

μ = 1.30 mm⁻¹

T = 291 K

Block, blue

0.28 × 0.24 × 0.22 mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

T_{min} = 0.712, *T_{max}* = 0.763

5510 measured reflections

1914 independent reflections

1589 reflections with *I* > 2σ(*I*)

R_{int} = 0.039

θ_{max} = 25.1°, θ_{min} = 2.7°

h = -9→11

k = -14→12

l = -11→11

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$ $wR(F^2) = 0.112$ $S = 1.07$

1914 reflections

142 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.05P)^2 + 1.33P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.70 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4721 (5)	0.3096 (4)	0.7859 (4)	0.0440 (10)
H1	0.3854	0.2781	0.7661	0.053*
C2	0.5127 (4)	0.3583 (4)	0.9007 (5)	0.0421 (10)
H2	0.4651	0.3609	0.9823	0.051*
C3	0.6429 (5)	0.4080 (4)	0.8837 (5)	0.0476 (11)
H3	0.6939	0.4531	0.9484	0.057*
C4	0.6766 (4)	0.3770 (4)	0.7570 (4)	0.0404 (10)
C5	0.8085 (4)	0.3823 (4)	0.6885 (4)	0.0417 (10)
H5A	0.8831	0.3906	0.7616	0.050*
H5B	0.8220	0.3116	0.6405	0.050*
C6	0.7106 (5)	0.4843 (4)	0.4717 (5)	0.0468 (11)
H6A	0.6259	0.4492	0.4970	0.056*
H6B	0.6924	0.5634	0.4536	0.056*
C7	0.7570 (5)	0.4316 (4)	0.3458 (5)	0.0466 (11)
H7A	0.6909	0.4462	0.2662	0.056*
H7B	0.7621	0.3509	0.3599	0.056*
Cl1	0.90322 (11)	0.80490 (9)	0.47487 (11)	0.0422 (3)
Cu1	1.0000	0.5000	0.5000	0.0320 (2)
N1	0.8171 (4)	0.4721 (3)	0.5882 (4)	0.0465 (9)
H1B	0.8199	0.5368	0.6377	0.056*
N2	0.8951 (4)	0.4749 (3)	0.3128 (4)	0.0451 (9)
H2A	0.8864	0.5396	0.2638	0.054*
H2B	0.9387	0.4245	0.2611	0.054*
O1	0.5768 (3)	0.3107 (3)	0.6947 (3)	0.0493 (8)
O2	0.9102 (3)	0.6962 (3)	0.4956 (3)	0.0493 (8)

O3	0.8334 (3)	0.8393 (2)	0.3418 (3)	0.0476 (8)
O4	1.0257 (3)	0.8481 (2)	0.4616 (3)	0.0462 (7)
O5	0.8431 (3)	0.8640 (2)	0.5750 (3)	0.0446 (7)

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.044 (3)	0.046 (2)	0.042 (2)	−0.018 (2)	0.0009 (19)	0.004 (2)
C2	0.041 (2)	0.045 (2)	0.042 (2)	0.0089 (19)	0.0103 (19)	0.0034 (19)
C3	0.048 (3)	0.049 (3)	0.048 (2)	−0.015 (2)	0.015 (2)	−0.009 (2)
C4	0.035 (2)	0.057 (3)	0.0306 (19)	−0.0051 (19)	0.0070 (17)	0.0011 (18)
C5	0.042 (2)	0.041 (2)	0.043 (2)	0.0078 (19)	0.0053 (19)	0.0011 (18)
C6	0.045 (3)	0.051 (3)	0.046 (2)	−0.006 (2)	0.012 (2)	0.019 (2)
C7	0.045 (3)	0.048 (3)	0.044 (2)	−0.012 (2)	−0.0140 (19)	0.012 (2)
Cl1	0.0397 (6)	0.0441 (6)	0.0442 (6)	0.0033 (4)	0.0113 (4)	−0.0060 (4)
Cu1	0.0324 (4)	0.0315 (4)	0.0333 (4)	−0.0005 (3)	0.0110 (3)	0.0024 (3)
N1	0.043 (2)	0.050 (2)	0.048 (2)	−0.0093 (17)	0.0127 (17)	0.0170 (17)
N2	0.043 (2)	0.043 (2)	0.048 (2)	−0.0126 (16)	−0.0073 (17)	−0.0004 (16)
O1	0.053 (2)	0.0482 (17)	0.0475 (17)	−0.0162 (15)	0.0109 (15)	−0.0149 (14)
O2	0.0527 (19)	0.0448 (18)	0.0530 (18)	0.0136 (14)	0.0218 (15)	0.0132 (14)
O3	0.0467 (17)	0.0419 (17)	0.0529 (18)	−0.0135 (14)	−0.0053 (14)	0.0056 (14)
O4	0.0469 (18)	0.0466 (17)	0.0474 (16)	−0.0021 (14)	0.0193 (14)	0.0084 (14)
O5	0.0430 (17)	0.0442 (16)	0.0467 (16)	0.0178 (14)	0.0040 (13)	−0.0015 (13)

Geometric parameters ($\text{\AA}$, $^\circ$)

C1—C2	1.268 (6)	C6—H6B	0.9700
C1—O1	1.384 (5)	C7—N2	1.494 (6)
C1—H1	0.9300	C7—H7A	0.9700
C2—C3	1.419 (6)	C7—H7B	0.9700
C2—H2	0.9300	Cl1—O2	1.309 (3)
C3—C4	1.319 (6)	Cl1—O4	1.313 (3)
C3—H3	0.9300	Cl1—O5	1.348 (3)
C4—O1	1.352 (5)	Cl1—O3	1.446 (3)
C4—C5	1.482 (6)	Cu1—N2	2.001 (4)
C5—N1	1.436 (5)	Cu1—N2 ⁱ	2.001 (4)
C5—H5A	0.9700	Cu1—N1	2.049 (4)
C5—H5B	0.9700	Cu1—N1 ⁱ	2.049 (4)
C6—C7	1.448 (6)	N1—H1B	0.9000
C6—N1	1.465 (6)	N2—H2A	0.9000
C6—H6A	0.9700	N2—H2B	0.9000
C2—C1—O1	109.5 (4)	N2—C7—H7B	109.4
C2—C1—H1	125.3	H7A—C7—H7B	108.0
O1—C1—H1	125.3	O2—Cl1—O4	111.3 (2)
C1—C2—C3	108.5 (4)	O2—Cl1—O5	115.5 (2)
C1—C2—H2	125.8	O4—Cl1—O5	107.9 (2)
C3—C2—H2	125.8	O2—Cl1—O3	115.2 (2)

C4—C3—C2	105.7 (4)	O4—Cl1—O3	100.24 (19)
C4—C3—H3	127.1	O5—Cl1—O3	105.3 (2)
C2—C3—H3	127.1	N2—Cu1—N2 ⁱ	180.000 (1)
C3—C4—O1	109.9 (4)	N2—Cu1—N1	86.25 (16)
C3—C4—C5	131.9 (4)	N2 ⁱ —Cu1—N1	93.75 (16)
O1—C4—C5	116.9 (3)	N2—Cu1—N1 ⁱ	93.75 (16)
N1—C5—C4	114.6 (4)	N2 ⁱ —Cu1—N1 ⁱ	86.25 (16)
N1—C5—H5A	108.6	N1—Cu1—N1 ⁱ	180.0
C4—C5—H5A	108.6	C5—N1—C6	120.0 (4)
N1—C5—H5B	108.6	C5—N1—Cu1	119.0 (3)
C4—C5—H5B	108.6	C6—N1—Cu1	105.4 (3)
H5A—C5—H5B	107.6	C5—N1—H1B	107.1
C7—C6—N1	109.3 (4)	C6—N1—H1B	107.4
C7—C6—H6A	109.8	Cu1—N1—H1B	94.6
N1—C6—H6A	109.8	C7—N2—Cu1	106.0 (3)
C7—C6—H6B	109.8	C7—N2—H2A	110.5
N1—C6—H6B	109.8	Cu1—N2—H2A	110.5
H6A—C6—H6B	108.3	C7—N2—H2B	110.5
C6—C7—N2	111.2 (4)	Cu1—N2—H2B	110.5
C6—C7—H7A	109.4	H2A—N2—H2B	108.7
N2—C7—H7A	109.4	C4—O1—C1	105.8 (3)
C6—C7—H7B	109.4		
O1—C1—C2—C3	7.7 (5)	N2—Cu1—N1—C5	121.4 (3)
C1—C2—C3—C4	−4.6 (5)	N2 ⁱ —Cu1—N1—C5	−58.6 (3)
C2—C3—C4—O1	−0.5 (5)	N2—Cu1—N1—C6	−16.7 (3)
C2—C3—C4—C5	−166.5 (5)	N2 ⁱ —Cu1—N1—C6	163.3 (3)
C3—C4—C5—N1	−102.5 (6)	C6—C7—N2—Cu1	36.2 (4)
O1—C4—C5—N1	92.3 (5)	N1—Cu1—N2—C7	−9.8 (3)
N1—C6—C7—N2	−53.2 (5)	N1 ⁱ —Cu1—N2—C7	170.2 (3)
C4—C5—N1—C6	−54.1 (5)	C3—C4—O1—C1	4.9 (5)
C4—C5—N1—Cu1	173.9 (3)	C5—C4—O1—C1	173.3 (4)
C7—C6—N1—C5	−96.6 (5)	C2—C1—O1—C4	−7.9 (5)
C7—C6—N1—Cu1	41.1 (4)		

Symmetry code: (i) $-x+2, -y+1, -z+1$.