

3,4-Dinitro-1*H*-pyrazole benzene 0.25-solvate

Yong-Xiang Li, Shan Du and Jian-Long Wang*

School of Chemical Engineering and Environment, North University of China, Taiyuan, People's Republic of China
Correspondence e-mail: wangjianlong@nuc.edu.cn


Received 11 April 2011; accepted 27 April 2011

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.060; wR factor = 0.105; data-to-parameter ratio = 13.9.

The asymmetric unit of the title compound, $4\text{C}_3\text{H}_2\text{N}_2\text{O}_4 \cdot \text{C}_6\text{H}_6$, contains two independent dinitropyrazole molecules and half a benzene solvent molecule, which lies on a crystallographic inversion centre. Each pyrazole ring is essentially planar (mean deviations of 0.009 and 0.002 Å), with the two nitro groups rotated out of the plane [dihedral angles = 11.7 (2)/31.1 (1) and 21.8 (2)/25.0 (1)° for the two molecules].

Related literature

For the biological properties of polynitropyrazoles, see: Alejandre-Durán *et al.* (1986); Grigor'ev *et al.* (1998); Xuan *et al.* (1999). For their detonation properties, see: Keshavarz *et al.* (2007); Zaitsev *et al.* (2009). For the synthesis, see: Katritzky *et al.* (2005).



Experimental

Crystal data

$2\text{C}_3\text{H}_2\text{N}_2\text{O}_4 \cdot 0.5\text{C}_6\text{H}_6$
 $M_r = 355.23$
Monoclinic, $P2_1/n$
 $a = 7.4579$ (15) Å
 $b = 9.787$ (2) Å
 $c = 19.534$ (4) Å
 $\beta = 94.87$ (3)°

$V = 1420.7$ (5) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.15$ mm⁻¹
 $T = 123$ K
0.30 × 0.20 × 0.20 mm

Data collection

Rigaku R-Axis RAPID IP diffractometer
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.957$, $T_{\max} = 0.971$

3454 measured reflections
3256 independent reflections
1267 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.084$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.105$
 $S = 0.92$
3256 reflections
235 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.27$ e Å⁻³
 $\Delta\rho_{\min} = -0.33$ e Å⁻³

Data collection: *RAPID-AUTO* (Rigaku, 2000); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku, 2000); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the China North Industries Group Corporation for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2110).

References

- Alejandre-Durán, E., Ruiz-Rubio, M., Claramunt, R. M., López, C. & Pueyo, C. (1986). *Environ. Mutagen.* **8**, 611–619.
- Grigor'ev, N. B., Kalinkina, M. A., Chechekin, G. V., Nikitin, V. B. & Engalycheva, G. N. (1998). *Pharm. Chem. J.* **32**, 127–131.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Katritzky, A. R., Scriven, E. F. V., Majumder, S., Akhmedova, R. G., Akhmedov, N. G. & Vakulenko, A. V. (2005). *ARKIVOC*, **iii**, 179–191.
- Keshavarz, M. H., Pouretdal, H. R. & Semnani, A. (2007). *J. Hazard. Mater.* **141**, 803–807.
- Rigaku (2000). *RAPID-AUTO* and *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xuan, B., Zhou, Y. H. & Verma, R. S. (1999). *J. Ocul. Pharmacol. Ther.* **15**, 135–142.
- Zaitsev, A. A., Dalinger, I. L. & Shaevelev, S. A. (2009). *Russ. Chem. Rev.* **78**, 589–627.

supporting information

Acta Cryst. (2011). E67, o1369 [https://doi.org/10.1107/S1600536811015996]

3,4-Dinitro-1*H*-pyrazole benzene 0.25-solvate

Yong-Xiang Li, Shan Du and Jian-Long Wang

S1. Comment

Polynitropyrazole systems have been investigated extensively because of their biological activity (Alejandre-Durán *et al.*, 1986; Grigor'ev *et al.*, 1998; Xuan *et al.*, 1999). Recently, these so called "high energy density materials" have attracted renewed attention because of their favorable detonation performance (Keshavarz *et al.*, 2007; Zaitsev *et al.*, 2009). As a potential candidate, 3,4-dinitropyrazole was synthesized by the nitration of pyrazole (Katritzky *et al.*, 2005). Here we report the crystal structure of the title compound, the benzene solvate 4(C₃H₂N₂O₄). C₆H₆ (I).

In the crystal structure of (I) (Fig. 1), the nitro groups are twisted with respect to the pyrazole plane, making dihedral angles of 11.7° (N1/O1, O2), 31.1° (N2/O3, O4) (molecule A) and 21.8° (N5/O5, O6), 25.0° (N6/O7, O8) (molecule B).

S2. Experimental

The title compound was prepared by the nitric acid of pyrazole according to the literature method (Katritzky *et al.*, 2005). Single crystals suitable for X-ray diffraction were obtained by evaporation of a solution of the compound in benzene at room temperature.

S3. Refinement

All H atoms were positioned geometrically and treated as riding, with C—H bond lengths constrained to 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, and the N—H bond = 0.87 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N})$.




Figure 1

The molecular structure of the asymmetric unit of the title compound with the atom numbering scheme. The benzene molecule has inversion symmetry [symmetry code (i): $-x + 2, -y, -z$]. Hydrogen atoms are omitted and displacement ellipsoids are drawn at the 30% probability level.

3,4-Dinitro-1*H*-pyrazole benzene 0.25-solvate*Crystal data*2C₃H₂N₄O₄·0.5C₆H₆ $M_r = 355.23$ Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

 $a = 7.4579 (15) \text{ \AA}$ $b = 9.787 (2) \text{ \AA}$ $c = 19.534 (4) \text{ \AA}$ $\beta = 94.87 (3)^\circ$ $V = 1420.7 (5) \text{ \AA}^3$ $Z = 4$ $F(000) = 724$ $D_x = 1.661 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3256 reflections

 $\theta = 2.1\text{--}27.5^\circ$ $\mu = 0.15 \text{ mm}^{-1}$ $T = 123 \text{ K}$

Block, colorless

 $0.30 \times 0.20 \times 0.20 \text{ mm}$ *Data collection*

Rigaku R-Axis RAPID IP

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $10.00 \text{ pixels mm}^{-1}$ ω scans

Absorption correction: multi-scan

(ABSCOR; Higashi, 1995)

 $T_{\min} = 0.957, T_{\max} = 0.971$

3256 measured reflections

3256 independent reflections

1267 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.084$ $\theta_{\max} = 27.5^\circ, \theta_{\min} = 2.1^\circ$ $h = -9 \rightarrow 9$ $k = -12 \rightarrow 12$ $l = -25 \rightarrow 25$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.060$ $wR(F^2) = 0.105$ $S = 0.92$

3256 reflections

235 parameters

2 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from

neighbouring sites

H atoms treated by a mixture of independent

and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.27 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.33 \text{ e \AA}^{-3}$

Extinction correction: SHELXL97 (Sheldrick,

2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0081 (6)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.9887 (4)	0.1907 (3)	0.19586 (16)	0.0327 (8)
C2	1.1357 (4)	0.1088 (4)	0.19047 (17)	0.0420 (9)

H2	1.2585	0.1359	0.1953	0.050*
C3	0.8410 (4)	0.1014 (3)	0.18468 (16)	0.0289 (8)
C4	0.5086 (4)	0.5510 (3)	0.09984 (17)	0.0387 (9)
C5	0.3685 (4)	0.6383 (3)	0.11466 (16)	0.0344 (8)
C6	0.6606 (4)	0.6268 (4)	0.10959 (18)	0.0538 (11)
H6	0.7801	0.5975	0.1045	0.065*
C7	1.1648 (8)	0.0615 (7)	0.0108 (2)	0.0902 (16)
H7	1.2790	0.1045	0.0180	0.108*
C8	1.0067 (11)	0.1372 (5)	0.0150 (2)	0.0921 (19)
H8	1.0132	0.2318	0.0258	0.111*
C9	0.8469 (8)	0.0760 (7)	0.0040 (2)	0.0880 (17)
H9	0.7399	0.1279	0.0063	0.106*
N1	0.9970 (4)	0.3361 (3)	0.20565 (14)	0.0409 (7)
N2	0.6477 (3)	0.1259 (3)	0.18791 (15)	0.0431 (8)
N3	1.0721 (3)	−0.0172 (3)	0.17704 (15)	0.0415 (8)
N4	0.8896 (3)	−0.0246 (3)	0.17325 (13)	0.0364 (7)
N5	0.5049 (4)	0.4072 (3)	0.08235 (17)	0.0540 (9)
N6	0.1748 (3)	0.6197 (3)	0.10954 (15)	0.0422 (7)
N7	0.6079 (4)	0.7520 (4)	0.12796 (17)	0.0533 (9)
N8	0.4295 (3)	0.7621 (3)	0.13217 (14)	0.0455 (8)
O1	1.1467 (3)	0.3853 (3)	0.22096 (13)	0.0593 (8)
O2	0.8580 (3)	0.4027 (3)	0.19682 (14)	0.0638 (8)
O3	0.6020 (3)	0.2140 (3)	0.22658 (13)	0.0616 (8)
O4	0.5471 (3)	0.0518 (3)	0.15166 (14)	0.0585 (8)
O5	0.6340 (4)	0.3642 (3)	0.05424 (15)	0.0840 (10)
O6	0.3762 (4)	0.3390 (3)	0.09674 (17)	0.0864 (10)
O7	0.1130 (3)	0.5307 (3)	0.07052 (13)	0.0636 (8)
O8	0.0861 (3)	0.6953 (3)	0.14284 (14)	0.0631 (8)
H7A	0.684 (5)	0.820 (4)	0.127 (3)	0.17 (2)*
H3	1.138 (3)	−0.091 (2)	0.1735 (17)	0.052 (11)*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0275 (18)	0.0321 (19)	0.038 (2)	−0.0016 (15)	0.0019 (15)	0.0026 (17)
C2	0.0259 (18)	0.042 (2)	0.057 (2)	0.0014 (17)	−0.0032 (16)	−0.004 (2)
C3	0.0226 (16)	0.0290 (18)	0.0343 (18)	0.0030 (14)	−0.0015 (14)	0.0029 (16)
C4	0.0326 (19)	0.039 (2)	0.044 (2)	0.0008 (17)	−0.0042 (15)	−0.001 (2)
C5	0.0306 (18)	0.034 (2)	0.038 (2)	−0.0019 (16)	−0.0017 (14)	−0.0027 (18)
C6	0.0279 (19)	0.070 (3)	0.063 (3)	−0.001 (2)	−0.0027 (18)	0.013 (3)
C7	0.112 (5)	0.101 (5)	0.056 (3)	−0.015 (4)	−0.007 (3)	0.007 (3)
C8	0.177 (6)	0.047 (3)	0.047 (3)	−0.006 (4)	−0.019 (4)	−0.001 (3)
C9	0.127 (5)	0.089 (5)	0.046 (3)	0.043 (4)	−0.004 (3)	−0.002 (3)
N1	0.0403 (18)	0.0392 (19)	0.0422 (18)	−0.0050 (15)	−0.0027 (14)	−0.0002 (16)
N2	0.0299 (16)	0.0406 (19)	0.058 (2)	−0.0009 (15)	−0.0004 (15)	0.0044 (18)
N3	0.0257 (15)	0.042 (2)	0.056 (2)	0.0115 (15)	−0.0010 (13)	−0.0006 (17)
N4	0.0232 (14)	0.0343 (17)	0.0511 (18)	0.0016 (12)	0.0007 (12)	0.0042 (15)
N5	0.051 (2)	0.048 (2)	0.061 (2)	0.0108 (18)	−0.0078 (17)	−0.0066 (19)

N6	0.0311 (16)	0.0382 (18)	0.056 (2)	−0.0012 (15)	−0.0003 (14)	0.0049 (18)
N7	0.045 (2)	0.049 (2)	0.063 (2)	−0.0178 (18)	−0.0099 (16)	0.0041 (19)
N8	0.0431 (18)	0.0332 (18)	0.059 (2)	−0.0049 (15)	−0.0009 (15)	−0.0002 (17)
O1	0.0446 (15)	0.0533 (17)	0.0775 (19)	−0.0211 (14)	−0.0100 (13)	−0.0038 (16)
O2	0.0497 (15)	0.0403 (16)	0.099 (2)	0.0119 (13)	−0.0072 (15)	0.0013 (16)
O3	0.0338 (14)	0.0637 (19)	0.089 (2)	0.0096 (14)	0.0136 (13)	−0.0253 (18)
O4	0.0272 (13)	0.0502 (17)	0.096 (2)	−0.0024 (12)	−0.0094 (12)	−0.0134 (17)
O5	0.071 (2)	0.078 (2)	0.105 (2)	0.0246 (17)	0.0199 (17)	−0.030 (2)
O6	0.082 (2)	0.0369 (18)	0.141 (3)	−0.0096 (16)	0.015 (2)	−0.0024 (19)
O7	0.0371 (14)	0.066 (2)	0.085 (2)	−0.0130 (14)	−0.0100 (13)	−0.0203 (18)
O8	0.0405 (15)	0.0539 (18)	0.098 (2)	0.0059 (14)	0.0231 (14)	−0.0095 (18)

Geometric parameters (Å, °)

C1—C2	1.370 (4)	C8—C9	1.335 (7)
C1—C3	1.409 (4)	C8—H8	0.9500
C1—N1	1.436 (4)	C9—C7 ⁱ	1.377 (7)
C2—N3	1.339 (4)	C9—H9	0.9500
C2—H2	0.9500	N1—O2	1.224 (3)
C3—N4	1.310 (4)	N1—O1	1.229 (3)
C3—N2	1.468 (3)	N2—O3	1.214 (3)
C4—C6	1.355 (4)	N2—O4	1.225 (3)
C4—C5	1.398 (4)	N3—N4	1.359 (3)
C4—N5	1.448 (4)	N3—H3	0.876 (10)
C5—N8	1.328 (4)	N5—O6	1.221 (4)
C5—N6	1.451 (4)	N5—O5	1.223 (4)
C6—N7	1.344 (4)	N6—O8	1.217 (3)
C6—H6	0.9500	N6—O7	1.222 (3)
C7—C9 ⁱ	1.377 (7)	N7—N8	1.344 (4)
C7—C8	1.401 (7)	N7—H7A	0.878 (10)
C7—H7	0.9500		
C2—C1—C3	104.2 (3)	C8—C9—C7 ⁱ	120.8 (5)
C2—C1—N1	124.3 (3)	C8—C9—H9	119.6
C3—C1—N1	131.3 (3)	C7 ⁱ —C9—H9	119.6
N3—C2—C1	106.3 (3)	O2—N1—O1	124.5 (3)
N3—C2—H2	126.8	O2—N1—C1	118.8 (3)
C1—C2—H2	126.8	O1—N1—C1	116.6 (3)
N4—C3—C1	112.7 (3)	O3—N2—O4	126.1 (3)
N4—C3—N2	116.7 (3)	O3—N2—C3	118.1 (3)
C1—C3—N2	130.5 (3)	O4—N2—C3	115.7 (3)
C6—C4—C5	105.5 (3)	C2—N3—N4	113.3 (3)
C6—C4—N5	124.4 (3)	C2—N3—H3	125 (2)
C5—C4—N5	130.0 (3)	N4—N3—H3	121 (2)
N8—C5—C4	111.4 (3)	C3—N4—N3	103.4 (3)
N8—C5—N6	116.7 (3)	O6—N5—O5	125.4 (3)
C4—C5—N6	131.7 (3)	O6—N5—C4	118.5 (3)
N7—C6—C4	106.1 (3)	O5—N5—C4	116.1 (3)

N7—C6—H6	127.0	O8—N6—O7	125.0 (3)
C4—C6—H6	127.0	O8—N6—C5	118.0 (3)
C9 ⁱ —C7—C8	119.4 (5)	O7—N6—C5	117.0 (3)
C9 ⁱ —C7—H7	120.3	N8—N7—C6	113.4 (3)
C8—C7—H7	120.3	N8—N7—H7A	126 (3)
C9—C8—C7	119.8 (5)	C6—N7—H7A	119 (3)
C9—C8—H8	120.1	C5—N8—N7	103.7 (3)
C7—C8—H8	120.1		
C3—C1—C2—N3	0.1 (4)	C1—C3—N2—O3	−29.0 (5)
N1—C1—C2—N3	−175.8 (3)	N4—C3—N2—O4	−31.6 (4)
C2—C1—C3—N4	−0.1 (4)	C1—C3—N2—O4	152.5 (3)
N1—C1—C3—N4	175.4 (3)	C1—C2—N3—N4	−0.1 (4)
C2—C1—C3—N2	175.9 (3)	C1—C3—N4—N3	0.0 (4)
N1—C1—C3—N2	−8.6 (6)	N2—C3—N4—N3	−176.6 (3)
C6—C4—C5—N8	0.5 (4)	C2—N3—N4—C3	0.1 (4)
N5—C4—C5—N8	176.4 (4)	C6—C4—N5—O6	155.9 (4)
C6—C4—C5—N6	175.6 (3)	C5—C4—N5—O6	−19.3 (6)
N5—C4—C5—N6	−8.5 (6)	C6—C4—N5—O5	−23.7 (5)
C5—C4—C6—N7	−1.0 (4)	C5—C4—N5—O5	161.0 (3)
N5—C4—C6—N7	−177.2 (3)	N8—C5—N6—O8	−25.2 (4)
C9 ⁱ —C7—C8—C9	−0.8 (8)	C4—C5—N6—O8	159.9 (3)
C7—C8—C9—C7 ⁱ	0.8 (9)	N8—C5—N6—O7	153.1 (3)
C2—C1—N1—O2	166.3 (3)	C4—C5—N6—O7	−21.8 (5)
C3—C1—N1—O2	−8.5 (6)	C4—C6—N7—N8	1.2 (4)
C2—C1—N1—O1	−12.0 (5)	C4—C5—N8—N7	0.3 (4)
C3—C1—N1—O1	173.2 (3)	N6—C5—N8—N7	−175.7 (3)
N4—C3—N2—O3	146.8 (3)	C6—N7—N8—C5	−0.9 (4)

Symmetry code: (i) $-x+2, -y, -z$.