

Diaquabis(propane-1,3-diamine)-copper(II) bis[diamminetetakis(thiocyanato- κ N)chromate(III)] dimethyl sulfoxide octasolvate

Vitalina M. Nikitina,^a Oksana V. Nesterova,^a Roman I. Zubatyuk,^b Oleg V. Shishkin^b and Julia A. Rusanova^{a*}

^aNational Taras Shevchenko University, Department of Chemistry, Volodymyrska Str. 64, 01033 Kyiv, Ukraine, and ^bInstitute for Scintillation Materials, "Institute for Single Crystals", National Academy of Sciences of Ukraine, Lenina ave. 60, Kharkov 61001, Ukraine

Correspondence e-mail: rusanova_j@yahoo.com


Received 11 May 2011; accepted 26 May 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.037; wR factor = 0.097; data-to-parameter ratio = 24.0.

The ionic title complex, $[\text{Cu}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2][\text{Cr}(\text{NCS})_4(\text{NH}_3)_2] \cdot 8\text{C}_2\text{H}_6\text{OS}$, consists of complex $[\text{Cu}(\text{dipr})_2(\text{H}_2\text{O})_2]^{2+}$ copper cations (dipr is propane-1,3-diamine), complex $[\text{Cr}(\text{NCS})_4(\text{NH}_3)_2]^-$ anions and uncoordinated solvent dimethyl sulfoxide (DMSO) molecules. All the metal atoms lie on crystallographic centers of symmetry. The cations are connected to the anions through $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds between the NH_3 molecules of the anion and the water molecules of the cation. The DMSO molecules are involved in hydrogen-bonded linkage of the $[\text{Cr}(\text{NCS})_4(\text{NH}_3)_2]^-$ anions into supramolecular chains through bridging O atoms. A network of hydrogen bonds as well as short $\text{S} \cdots \text{S}$ contacts [3.5159 (12) and 3.4880 (12) Å] between the NCS groups of the complex anions link the molecules into a three-dimensional supramolecular network.

Related literature

For background to direct synthesis see: Nesterov *et al.* (2004, 2006); Kovbasyuk *et al.* (1997, 1998); Vassilyeva *et al.* (1997). For the structures of related complexes, see: Zhang *et al.* (2001); Cucos *et al.* (2006); Cherkasova & Gorunova (2003); Kolotilov *et al.* (2010).



Experimental

Crystal data

$[\text{Cu}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2] \cdot$ $[\text{Cr}(\text{NCS})_4(\text{NH}_3)_2] \cdot 8(\text{C}_2\text{H}_6\text{OS})$	$\beta = 89.664$ (7) $^\circ$
$M_r = 1509.63$	$\gamma = 61.535$ (10) $^\circ$
Triclinic, $P\bar{1}$	$V = 1724.9$ (3) Å ³
$a = 12.2609$ (11) Å	$Z = 1$
$b = 12.2772$ (12) Å	Mo $K\alpha$ radiation
$c = 13.8578$ (12) Å	$\mu = 1.15$ mm ⁻¹
$\alpha = 72.466$ (8) $^\circ$	$T = 100$ K
	$0.6 \times 0.4 \times 0.3$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer	15229 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	8482 independent reflections
$T_{\min} = 0.58$, $T_{\max} = 0.71$	6966 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$	353 parameters
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.76$ e Å ⁻³
8482 reflections	$\Delta\rho_{\text{min}} = -0.56$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1}-\text{H1E} \cdots \text{O2}^i$	0.87	1.88	2.732 (2)	165
$\text{O1}-\text{H1E} \cdots \text{S5}^i$	0.87	2.87	3.5905 (18)	142
$\text{O1}-\text{H1F} \cdots \text{O4}$	0.87	1.98	2.786 (2)	153
$\text{N1}-\text{H1A} \cdots \text{O2}^{ii}$	0.92	2.10	2.995 (3)	163
$\text{N1}-\text{H1B} \cdots \text{S4}^{iii}$	0.92	2.70	3.490 (2)	145
$\text{N2}-\text{H2A} \cdots \text{O4}$	0.92	2.33	3.052 (2)	135
$\text{N2}-\text{H2B} \cdots \text{O3}$	0.92	2.24	3.091 (3)	153
$\text{N5}-\text{H5A} \cdots \text{O2}$	0.91	2.11	3.003 (2)	167
$\text{N5}-\text{H5B} \cdots \text{O3}^{iv}$	0.91	2.21	3.079 (2)	161
$\text{N5}-\text{H5C} \cdots \text{O5}^v$	0.91	2.09	2.966 (2)	160
$\text{N8}-\text{H8A} \cdots \text{O1}^{iii}$	0.91	2.08	2.956 (2)	162
$\text{N8}-\text{H8B} \cdots \text{O5}^{vi}$	0.91	2.19	3.054 (2)	159
$\text{N8}-\text{H8C} \cdots \text{O3}^{vii}$	0.91	2.10	2.981 (2)	162

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $x - 1, y + 1, z$; (iii) $-x, -y + 1, -z$; (iv) $x + 1, y - 1, z$; (v) $-x + 1, -y + 1, -z + 1$; (vi) $-x, -y + 1, -z + 1$; (vii) $x, y - 1, z$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors gratefully acknowledge the Ukrainian State Fund for Fundamental Researchers (SFFR) for financial support of the Research Program (Chemistry).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2167).

References

- Cherkasova, T. G. & Gorunova, I. P. (2003). *Zh. Neorg. Khim.* **48**, 611–615.
 Cucos, A., Avarvari, N., Andruh, M., Journaux, Y., Muller, A. & Schmidtman, M. (2006). *Eur. J. Inorg. Chem.* pp. 903–907.
 Kolotilov, S. V., Cador, O., Gavrilenko, K. S., Golhen, S., Ouahab, L. & Pavlishchuk, V. V. (2010). *Eur. J. Inorg. Chem.* **8**, 1255–1266.

- Kovbasyuk, L. A., Babich, O. A. & Kokozay, V. N. (1997). *Polyhedron*, pp. 161–163.
- Kovbasyuk, L. A., Vassilyeva, O. Yu., Kokozay, V. N., Linert, W., Reedijk, J., Skelton, B. W. & Oliver, A. G. (1998). *J. Chem. Soc. Dalton Trans.* pp. 2735–2738.
- Nesterov, D. S., Kokozay, V. N., Dyakonenko, V. V., Shishkin, O. V., Jezierska, J., Ozarowski, F., Kirillov, A. M., Kopylovich, M. N. & Pombeiro, A. J. L. (2006). *Chem. Commun.* pp. 4605–4607.
- Nesterov, D. S., Makhankova, V. G., Vassilyeva, O. Yu., Kokozay, V. N., Kovbasyuk, L. A., Skelton, B. W. & Jezierska, J. (2004). *Inorg. Chem.* pp. 7868–7876.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vassilyeva, O. Yu., Kokozay, V. N., Zhukova, N. I. & Kovbasyuk, L. A. (1997). *Polyhedron*, pp. 263–266.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhang, K.-L., Chen, W., Xu, Y., Wang, Z., Zhong, Z. J. & You, X.-Z. (2001). *Polyhedron*, pp. 2033–2036.

supporting information

Acta Cryst. (2011). E67, m849–m850 [doi:10.1107/S160053681102023X]

Diaquabis(propane-1,3-diamine)copper(II) bis[diamminetetakis(thiocyanato- κ N)chromate(III)] dimethyl sulfoxide octasolvate

Vitalina M. Nikitina, Oksana V. Nesterova, Roman I. Zubatyuk, Oleg V. Shishkin and Julia A. Rusanova

S1. Comment


It has been shown that direct synthesis is an efficient method to obtain novel homo- and heterometallic complexes (Nesterov *et al.* (2004, 2006); Kovbasyuk *et al.* (1997, 1998); Vassilyeva *et al.* (1997)). Continuing our investigations in this paper we present a novel Cu/Cr heterometallic ionic complex which has been synthesized using zerovalent copper, Reinecke's salt and 1,3-propylenediamine as starting materials. As it shown on Fig. 1 Cu atoms in complex cations are in square bypyramidal coordination environment with four N atoms in equatorial position and two O atoms of the H₂O molecules in axial position. The Cr centers are coordinated to six N atoms - four NCS-groups in equatorial position and two NH₃ molecules in axial position. The bond distances and angles in the title molecule agree well with the corresponding bond distances and angles reported in closely related compounds (Zhang *et al.*, 2001, Cucos *et al.*, 2006; Cherkasova *et al.*, 2003; Kolotilov *et al.*, 2010). There are short interanionic S...S contacts between NCS-groups of the complex anions with the distances 3.5159 (12) and 3.4880 (12) Å whereas sum of standard Van-der-Vaals radius of the sulfur atom is 3.68 Å. The crystal packing of the title compound is presented on Fig 2.

S2. Experimental

For the preparation of the title compound, copper powder 0.04 g (0.63 mmol), NH₄[Cr(NCS)₄(NH₃)₂].H₂O 0.45 g (1.26 mmol), 0.11 ml (1.26 mmol) 1,3-propylenediamine (dipr), 20 ml of DMSO, were heated to 323–333 K and stirred magnetically for 15 min, until total dissolution of the copper powder was observed. Addition of a few ml of the PrⁱOH to the cooled solution leads to precipitation within few days of the dark violet crystals suitable for X-ray analysis. They were collected by filter-suction, washed with dry PrⁱOH and finally dried in *vacuo* at room temperature (yield: 0.59 g, 69%).

S3. Refinement

The structure was solved by direct methods and refined by the full-matrix least-squares technique in the anisotropic approximation for non-hydrogen atoms using the *BRUKER SHELXTL* program package. All hydrogen atoms were placed at calculated positions which were refined as 'riding' model.

**Figure 1**

Molecular view of the title compound. DMSO molecules are omitted for clarity. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

Crystal packing of the title compound along *b* axis.

Diaquabis(propane-1,3-diamine)copper(II) bis[diamminetetrakis(thiocyanato- κ N)chromate(III)] dimethyl sulfoxide octasolvate

Crystal data

$[\text{Cu}(\text{C}_3\text{H}_{10}\text{N}_2)_2(\text{H}_2\text{O})_2] \cdot [\text{Cr}(\text{NCS})_4(\text{NH}_3)_2] \cdot 8(\text{C}_2\text{H}_6\text{OS})$

$M_r = 1509.63$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 12.2609$ (11) Å

$b = 12.2772$ (12) Å

$c = 13.8578$ (12) Å

$\alpha = 72.466$ (8)°

$\beta = 89.664$ (7)°

$\gamma = 61.535$ (10)°

$V = 1724.9$ (3) Å³

$Z = 1$

$F(000) = 789$

$D_x = 1.453$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4303 reflections

$\theta = 2.9\text{--}25.0^\circ$

$\mu = 1.15$ mm⁻¹

$T = 100$ K

Block, violet

$0.6 \times 0.4 \times 0.3$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3
diffractometer

Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator

Detector resolution: 16.1827 pixels mm⁻¹

ω and ϕ scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Oxford Diffraction, 2010)

$T_{\min} = 0.58$, $T_{\max} = 0.71$
 15229 measured reflections
 8482 independent reflections
 6966 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

$\theta_{\max} = 29.9^\circ$, $\theta_{\min} = 2.9^\circ$
 $h = -16 \rightarrow 16$
 $k = -16 \rightarrow 15$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.097$
 $S = 1.08$
 8482 reflections
 353 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0448P)^2 + 1.0385P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.76 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.56 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Xcalibur; Oxford Diffraction, (2010) CrysAlisPro, Oxford Diffraction (2010). Oxford Diffraction Ltd., Version 1.171.34.44 (release 25-10-2010 CrysAlis171 .NET) (compiled Oct 25 2010,18:11:34) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.13010 (15)	0.77778 (15)	-0.02683 (12)	0.0259 (3)
H1E	0.1484	0.8037	-0.0871	0.039*
H1F	0.1991	0.7425	0.0160	0.039*
Cu1	0.0000	1.0000	0.0000	0.01782 (9)
Cr1	1.0000	0.0000	0.5000	0.01666 (10)
Cr2	0.0000	0.5000	0.0000	0.01895 (11)
S1	1.22765 (6)	0.19198 (6)	0.58717 (5)	0.03186 (14)
S2	0.60022 (5)	0.32965 (5)	0.52979 (4)	0.02642 (13)
S3	0.28491 (6)	0.43926 (7)	0.26932 (6)	0.03845 (16)
S4	0.34351 (5)	0.12439 (6)	-0.04325 (5)	0.02791 (13)
S5	0.64253 (5)	0.22031 (5)	0.19080 (4)	0.02553 (12)
S6	0.30881 (5)	1.08392 (5)	0.23806 (4)	0.02493 (12)
S7	0.47022 (5)	0.58296 (6)	0.11082 (4)	0.02518 (12)
S8	0.17949 (5)	0.48082 (5)	0.67464 (4)	0.02321 (12)
O2	0.78274 (14)	0.13337 (16)	0.19770 (12)	0.0264 (3)
O3	0.18069 (15)	1.09692 (16)	0.22392 (13)	0.0276 (3)
O4	0.36425 (15)	0.72167 (16)	0.05789 (12)	0.0275 (3)
O5	0.16592 (15)	0.58928 (15)	0.71241 (12)	0.0263 (3)

N1	-0.10137 (17)	0.93537 (17)	0.09092 (14)	0.0220 (4)
H1A	-0.1522	1.0008	0.1162	0.026*
H1B	-0.1535	0.9272	0.0496	0.026*
N2	0.13971 (17)	0.92577 (18)	0.12042 (13)	0.0212 (4)
H2A	0.2133	0.9049	0.0940	0.025*
H2B	0.1236	0.9936	0.1440	0.025*
N3	1.08964 (17)	0.08516 (17)	0.53719 (13)	0.0209 (4)
N4	0.83883 (17)	0.13557 (17)	0.52307 (14)	0.0220 (4)
N5	0.96720 (16)	0.11943 (17)	0.34908 (13)	0.0186 (3)
H5A	0.9124	0.1123	0.3108	0.022*
H5B	1.0410	0.0939	0.3241	0.022*
H5C	0.9338	0.2046	0.3460	0.022*
N6	0.12608 (17)	0.47359 (18)	0.10717 (15)	0.0242 (4)
N7	0.13479 (17)	0.35414 (18)	-0.03865 (15)	0.0239 (4)
N8	-0.03243 (17)	0.36323 (17)	0.10530 (14)	0.0213 (4)
H8A	-0.0753	0.3381	0.0719	0.026*
H8B	-0.0786	0.3996	0.1501	0.026*
H8C	0.0423	0.2917	0.1403	0.026*
C1	-0.0366 (2)	0.8106 (2)	0.17926 (17)	0.0254 (5)
H1C	0.0134	0.7365	0.1541	0.031*
H1D	-0.0998	0.7942	0.2157	0.031*
C2	0.0489 (2)	0.8171 (2)	0.25237 (17)	0.0266 (5)
H2C	0.0741	0.7434	0.3179	0.032*
H2D	0.0018	0.9003	0.2669	0.032*
C3	0.1658 (2)	0.8099 (2)	0.21183 (16)	0.0256 (5)
H3A	0.2224	0.8027	0.2669	0.031*
H3B	0.2103	0.7291	0.1935	0.031*
C4	1.1462 (2)	0.1305 (2)	0.55823 (16)	0.0215 (4)
C5	0.7391 (2)	0.2164 (2)	0.52692 (16)	0.0207 (4)
C6	0.1937 (2)	0.4591 (2)	0.17413 (18)	0.0240 (4)
C7	0.2218 (2)	0.2583 (2)	-0.04054 (16)	0.0219 (4)
C8	0.5722 (2)	0.1351 (3)	0.1611 (2)	0.0333 (5)
H8D	0.5821	0.1333	0.0913	0.050*
H8E	0.4826	0.1801	0.1659	0.050*
H8F	0.6131	0.0447	0.2098	0.050*
C9	0.6182 (3)	0.1991 (3)	0.3196 (2)	0.0557 (10)
H9A	0.6514	0.1053	0.3578	0.084*
H9B	0.5280	0.2480	0.3214	0.084*
H9C	0.6619	0.2323	0.3511	0.084*
C10	0.4189 (2)	0.9225 (2)	0.2392 (2)	0.0311 (5)
H10A	0.4043	0.8578	0.2903	0.047*
H10B	0.5046	0.9045	0.2568	0.047*
H10C	0.4083	0.9167	0.1712	0.047*
C11	0.3450 (3)	1.0572 (3)	0.3702 (2)	0.0402 (6)
H11A	0.2926	1.1390	0.3842	0.060*
H11B	0.4340	1.0294	0.3870	0.060*
H11C	0.3285	0.9883	0.4122	0.060*
C12	0.6073 (2)	0.5762 (3)	0.0626 (2)	0.0329 (5)

H12A	0.6020	0.5784	-0.0086	0.049*
H12B	0.6818	0.4946	0.1046	0.049*
H12C	0.6135	0.6520	0.0652	0.049*
C13	0.5117 (2)	0.5740 (3)	0.23673 (19)	0.0354 (6)
H13A	0.5284	0.6457	0.2328	0.053*
H13B	0.5873	0.4892	0.2718	0.053*
H13C	0.4423	0.5824	0.2750	0.053*
C14	0.0292 (2)	0.4947 (3)	0.6669 (2)	0.0331 (5)
H14A	0.0074	0.4758	0.7358	0.050*
H14B	0.0302	0.4317	0.6367	0.050*
H14C	-0.0334	0.5844	0.6239	0.050*
C15	0.1823 (3)	0.5356 (3)	0.54062 (19)	0.0348 (6)
H15A	0.1095	0.6237	0.5080	0.052*
H15B	0.1791	0.4744	0.5104	0.052*
H15C	0.2599	0.5389	0.5298	0.052*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0239 (8)	0.0248 (8)	0.0277 (8)	-0.0124 (7)	0.0004 (6)	-0.0066 (6)
Cu1	0.01624 (17)	0.01650 (17)	0.01893 (17)	-0.00831 (14)	0.00019 (13)	-0.00339 (13)
Cr1	0.0145 (2)	0.0143 (2)	0.0189 (2)	-0.00612 (17)	0.00164 (17)	-0.00447 (17)
Cr2	0.0162 (2)	0.0144 (2)	0.0248 (2)	-0.00683 (18)	0.00464 (18)	-0.00617 (18)
S1	0.0354 (3)	0.0315 (3)	0.0385 (3)	-0.0230 (3)	0.0043 (3)	-0.0136 (3)
S2	0.0196 (3)	0.0210 (3)	0.0285 (3)	-0.0031 (2)	0.0046 (2)	-0.0073 (2)
S3	0.0249 (3)	0.0358 (3)	0.0474 (4)	-0.0128 (3)	-0.0097 (3)	-0.0085 (3)
S4	0.0189 (3)	0.0210 (3)	0.0415 (3)	-0.0054 (2)	0.0044 (2)	-0.0154 (2)
S5	0.0205 (3)	0.0208 (3)	0.0282 (3)	-0.0047 (2)	-0.0019 (2)	-0.0087 (2)
S6	0.0225 (3)	0.0225 (3)	0.0303 (3)	-0.0126 (2)	0.0066 (2)	-0.0071 (2)
S7	0.0186 (3)	0.0235 (3)	0.0323 (3)	-0.0096 (2)	0.0007 (2)	-0.0095 (2)
S8	0.0230 (3)	0.0177 (2)	0.0258 (3)	-0.0086 (2)	0.0018 (2)	-0.0060 (2)
O2	0.0186 (8)	0.0266 (8)	0.0288 (8)	-0.0080 (6)	-0.0002 (6)	-0.0083 (7)
O3	0.0203 (8)	0.0248 (8)	0.0363 (9)	-0.0109 (7)	0.0059 (7)	-0.0093 (7)
O4	0.0193 (8)	0.0240 (8)	0.0304 (8)	-0.0060 (6)	0.0000 (6)	-0.0061 (6)
O5	0.0292 (8)	0.0210 (8)	0.0282 (8)	-0.0115 (7)	0.0020 (7)	-0.0093 (6)
N1	0.0193 (9)	0.0189 (9)	0.0257 (9)	-0.0096 (7)	0.0015 (7)	-0.0046 (7)
N2	0.0185 (8)	0.0228 (9)	0.0208 (8)	-0.0095 (7)	0.0018 (7)	-0.0065 (7)
N3	0.0192 (9)	0.0166 (8)	0.0228 (8)	-0.0069 (7)	0.0025 (7)	-0.0048 (7)
N4	0.0202 (9)	0.0183 (9)	0.0233 (9)	-0.0078 (7)	0.0020 (7)	-0.0046 (7)
N5	0.0183 (8)	0.0173 (8)	0.0191 (8)	-0.0081 (7)	0.0033 (7)	-0.0059 (6)
N6	0.0192 (9)	0.0208 (9)	0.0308 (10)	-0.0094 (7)	0.0033 (8)	-0.0076 (8)
N7	0.0201 (9)	0.0198 (9)	0.0306 (9)	-0.0089 (7)	0.0064 (7)	-0.0085 (7)
N8	0.0205 (9)	0.0168 (8)	0.0269 (9)	-0.0100 (7)	0.0041 (7)	-0.0067 (7)
C1	0.0270 (11)	0.0196 (10)	0.0251 (10)	-0.0115 (9)	0.0022 (9)	-0.0018 (8)
C2	0.0279 (12)	0.0238 (11)	0.0197 (10)	-0.0085 (9)	0.0026 (9)	-0.0041 (8)
C3	0.0235 (11)	0.0232 (11)	0.0197 (10)	-0.0073 (9)	-0.0039 (8)	-0.0012 (8)
C4	0.0215 (10)	0.0176 (10)	0.0212 (10)	-0.0078 (8)	0.0026 (8)	-0.0045 (8)
C5	0.0222 (10)	0.0176 (10)	0.0204 (9)	-0.0096 (8)	0.0020 (8)	-0.0046 (8)

C6	0.0176 (10)	0.0173 (10)	0.0324 (11)	-0.0073 (8)	0.0041 (9)	-0.0046 (9)
C7	0.0194 (10)	0.0235 (11)	0.0267 (10)	-0.0122 (9)	0.0067 (8)	-0.0110 (9)
C8	0.0254 (12)	0.0302 (12)	0.0427 (14)	-0.0155 (10)	0.0030 (10)	-0.0075 (11)
C9	0.0343 (15)	0.064 (2)	0.0323 (14)	0.0075 (14)	0.0003 (12)	-0.0237 (14)
C10	0.0228 (11)	0.0241 (11)	0.0393 (13)	-0.0085 (9)	0.0057 (10)	-0.0070 (10)
C11	0.0467 (16)	0.0494 (17)	0.0368 (14)	-0.0317 (14)	0.0075 (12)	-0.0169 (12)
C12	0.0197 (11)	0.0330 (13)	0.0436 (14)	-0.0112 (10)	0.0056 (10)	-0.0133 (11)
C13	0.0267 (12)	0.0367 (14)	0.0314 (12)	-0.0080 (11)	-0.0039 (10)	-0.0101 (11)
C14	0.0312 (13)	0.0326 (13)	0.0399 (13)	-0.0195 (11)	0.0069 (11)	-0.0119 (11)
C15	0.0510 (16)	0.0367 (14)	0.0314 (12)	-0.0301 (13)	0.0165 (11)	-0.0166 (11)

Geometric parameters (Å, °)

O1—H1E	0.8692	N4—C5	1.165 (3)
O1—H1F	0.8704	N5—H5A	0.9100
Cu1—O1	2.5655 (16)	N5—H5B	0.9100
Cu1—N1	2.0202 (18)	N5—H5C	0.9100
Cu1—N1 ⁱ	2.0202 (18)	N6—C6	1.162 (3)
Cu1—N2 ⁱ	2.0442 (17)	N7—C7	1.159 (3)
Cu1—N2	2.0442 (17)	N8—H8A	0.9100
Cr1—N4	1.9888 (18)	N8—H8B	0.9100
Cr1—N4 ⁱⁱ	1.9888 (18)	N8—H8C	0.9100
Cr1—N3 ⁱⁱ	1.9983 (19)	C1—C2	1.508 (3)
Cr1—N3	1.9983 (19)	C1—H1C	0.9900
Cr1—N5	2.0702 (17)	C1—H1D	0.9900
Cr1—N5 ⁱⁱ	2.0702 (17)	C2—C3	1.509 (3)
Cr2—N6	1.9898 (19)	C2—H2C	0.9900
Cr2—N6 ⁱⁱⁱ	1.9898 (19)	C2—H2D	0.9900
Cr2—N7 ⁱⁱⁱ	1.9928 (18)	C3—H3A	0.9900
Cr2—N7	1.9928 (19)	C3—H3B	0.9900
Cr2—N8	2.0640 (17)	C8—H8D	0.9800
Cr2—N8 ⁱⁱⁱ	2.0641 (17)	C8—H8E	0.9800
S1—C4	1.623 (2)	C8—H8F	0.9800
S2—C5	1.615 (2)	C9—H9A	0.9800
S3—C6	1.617 (2)	C9—H9B	0.9800
S4—C7	1.622 (2)	C9—H9C	0.9800
S5—O2	1.5149 (16)	C10—H10A	0.9800
S5—C8	1.769 (3)	C10—H10B	0.9800
S5—C9	1.770 (3)	C10—H10C	0.9800
S6—O3	1.5090 (17)	C11—H11A	0.9800
S6—C11	1.780 (3)	C11—H11B	0.9800
S6—C10	1.783 (2)	C11—H11C	0.9800
S7—O4	1.5086 (16)	C12—H12A	0.9800
S7—C13	1.777 (3)	C12—H12B	0.9800
S7—C12	1.779 (2)	C12—H12C	0.9800
S8—O5	1.5108 (17)	C13—H13A	0.9800
S8—C14	1.769 (3)	C13—H13B	0.9800
S8—C15	1.781 (2)	C13—H13C	0.9800

N1—C1	1.483 (3)	C14—H14A	0.9800
N1—H1A	0.9200	C14—H14B	0.9800
N1—H1B	0.9200	C14—H14C	0.9800
N2—C3	1.489 (3)	C15—H15A	0.9800
N2—H2A	0.9200	C15—H15B	0.9800
N2—H2B	0.9200	C15—H15C	0.9800
N3—C4	1.159 (3)		
H1E—O1—H1F	105.3	Cr2—N8—H8B	109.5
N1—Cu1—N1 ⁱ	180.00 (11)	H8A—N8—H8B	109.5
N1—Cu1—N2 ⁱ	87.84 (7)	Cr2—N8—H8C	109.5
N1 ⁱ —Cu1—N2 ⁱ	92.16 (7)	H8A—N8—H8C	109.5
N1—Cu1—N2	92.16 (7)	H8B—N8—H8C	109.5
N1 ⁱ —Cu1—N2	87.84 (7)	N1—C1—C2	110.69 (19)
N2 ⁱ —Cu1—N2	180.0	N1—C1—H1C	109.5
N1—Cu1—O1	92.27 (6)	C2—C1—H1C	109.5
N1 ⁱ —Cu1—O1	87.73 (6)	N1—C1—H1D	109.5
N2 ⁱ —Cu1—O1	94.59 (6)	C2—C1—H1D	109.5
N2—Cu1—O1	85.41 (6)	H1C—C1—H1D	108.1
N4—Cr1—N4 ⁱⁱ	180.00 (10)	C1—C2—C3	113.34 (19)
N4—Cr1—N3 ⁱⁱ	89.65 (8)	C1—C2—H2C	108.9
N4 ⁱⁱ —Cr1—N3 ⁱⁱ	90.35 (8)	C3—C2—H2C	108.9
N4—Cr1—N3	90.35 (8)	C1—C2—H2D	108.9
N4 ⁱⁱ —Cr1—N3	89.65 (8)	C3—C2—H2D	108.9
N3 ⁱⁱ —Cr1—N3	180.00 (6)	H2C—C2—H2D	107.7
N4—Cr1—N5	90.24 (7)	N2—C3—C2	113.53 (18)
N4 ⁱⁱ —Cr1—N5	89.76 (7)	N2—C3—H3A	108.9
N3 ⁱⁱ —Cr1—N5	91.87 (7)	C2—C3—H3A	108.9
N3—Cr1—N5	88.13 (7)	N2—C3—H3B	108.9
N4—Cr1—N5 ⁱⁱ	89.76 (7)	C2—C3—H3B	108.9
N4 ⁱⁱ —Cr1—N5 ⁱⁱ	90.24 (7)	H3A—C3—H3B	107.7
N3 ⁱⁱ —Cr1—N5 ⁱⁱ	88.13 (7)	N3—C4—S1	178.9 (2)
N3—Cr1—N5 ⁱⁱ	91.87 (7)	N4—C5—S2	178.86 (19)
N5—Cr1—N5 ⁱⁱ	180.000 (1)	N6—C6—S3	178.4 (2)
N6—Cr2—N6 ⁱⁱⁱ	180.0	N7—C7—S4	179.9 (3)
N6—Cr2—N7 ⁱⁱⁱ	90.37 (8)	S5—C8—H8D	109.5
N6 ⁱⁱⁱ —Cr2—N7 ⁱⁱⁱ	89.63 (8)	S5—C8—H8E	109.5
N6—Cr2—N7	89.63 (8)	H8D—C8—H8E	109.5
N6 ⁱⁱⁱ —Cr2—N7	90.37 (8)	S5—C8—H8F	109.5
N7 ⁱⁱⁱ —Cr2—N7	180.00 (10)	H8D—C8—H8F	109.5
N6—Cr2—N8	89.48 (8)	H8E—C8—H8F	109.5
N6 ⁱⁱⁱ —Cr2—N8	90.52 (8)	S5—C9—H9A	109.5
N7 ⁱⁱⁱ —Cr2—N8	91.17 (7)	S5—C9—H9B	109.5
N7—Cr2—N8	88.83 (7)	H9A—C9—H9B	109.5
N6—Cr2—N8 ⁱⁱⁱ	90.52 (8)	S5—C9—H9C	109.5
N6 ⁱⁱⁱ —Cr2—N8 ⁱⁱⁱ	89.48 (8)	H9A—C9—H9C	109.5
N7 ⁱⁱⁱ —Cr2—N8 ⁱⁱⁱ	88.83 (7)	H9B—C9—H9C	109.5
N7—Cr2—N8 ⁱⁱⁱ	91.17 (7)	S6—C10—H10A	109.5

N8—Cr2—N8 ⁱⁱⁱ	180.00 (14)	S6—C10—H10B	109.5
O2—S5—C8	105.70 (11)	H10A—C10—H10B	109.5
O2—S5—C9	104.89 (11)	S6—C10—H10C	109.5
C8—S5—C9	98.98 (17)	H10A—C10—H10C	109.5
O3—S6—C11	105.98 (12)	H10B—C10—H10C	109.5
O3—S6—C10	105.91 (11)	S6—C11—H11A	109.5
C11—S6—C10	97.80 (13)	S6—C11—H11B	109.5
O4—S7—C13	106.08 (11)	H11A—C11—H11B	109.5
O4—S7—C12	106.13 (11)	S6—C11—H11C	109.5
C13—S7—C12	97.55 (13)	H11A—C11—H11C	109.5
O5—S8—C14	105.80 (11)	H11B—C11—H11C	109.5
O5—S8—C15	106.19 (11)	S7—C12—H12A	109.5
C14—S8—C15	97.93 (13)	S7—C12—H12B	109.5
C1—N1—Cu1	120.07 (14)	H12A—C12—H12B	109.5
C1—N1—H1A	107.3	S7—C12—H12C	109.5
Cu1—N1—H1A	107.3	H12A—C12—H12C	109.5
C1—N1—H1B	107.3	H12B—C12—H12C	109.5
Cu1—N1—H1B	107.3	S7—C13—H13A	109.5
H1A—N1—H1B	106.9	S7—C13—H13B	109.5
C3—N2—Cu1	121.96 (14)	H13A—C13—H13B	109.5
C3—N2—H2A	106.8	S7—C13—H13C	109.5
Cu1—N2—H2A	106.8	H13A—C13—H13C	109.5
C3—N2—H2B	106.8	H13B—C13—H13C	109.5
Cu1—N2—H2B	106.8	S8—C14—H14A	109.5
H2A—N2—H2B	106.7	S8—C14—H14B	109.5
C4—N3—Cr1	177.22 (18)	H14A—C14—H14B	109.5
C5—N4—Cr1	173.03 (18)	S8—C14—H14C	109.5
Cr1—N5—H5A	109.5	H14A—C14—H14C	109.5
Cr1—N5—H5B	109.5	H14B—C14—H14C	109.5
H5A—N5—H5B	109.5	S8—C15—H15A	109.5
Cr1—N5—H5C	109.5	S8—C15—H15B	109.5
H5A—N5—H5C	109.5	H15A—C15—H15B	109.5
H5B—N5—H5C	109.5	S8—C15—H15C	109.5
C6—N6—Cr2	175.38 (19)	H15A—C15—H15C	109.5
C7—N7—Cr2	166.49 (18)	H15B—C15—H15C	109.5
Cr2—N8—H8A	109.5		

Symmetry codes: (i) $-x, -y+2, -z$; (ii) $-x+2, -y, -z+1$; (iii) $-x, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1E \cdots O2 ^{iv}	0.87	1.88	2.732 (2)	165
O1—H1E \cdots S5 ^{iv}	0.87	2.87	3.5905 (18)	142
O1—H1F \cdots O4	0.87	1.98	2.786 (2)	153
N1—H1A \cdots O2 ^v	0.92	2.10	2.995 (3)	163
N1—H1B \cdots S4 ⁱⁱⁱ	0.92	2.70	3.490 (2)	145
N2—H2A \cdots O4	0.92	2.33	3.052 (2)	135

N2—H2B···O3	0.92	2.24	3.091 (3)	153
N5—H5A···O2	0.91	2.11	3.003 (2)	167
N5—H5B···O3 ^{vi}	0.91	2.21	3.079 (2)	161
N5—H5C···O5 ^{vii}	0.91	2.09	2.966 (2)	160
N8—H8A···O1 ⁱⁱⁱ	0.91	2.08	2.956 (2)	162
N8—H8B···O5 ^{viii}	0.91	2.19	3.054 (2)	159
N8—H8C···O3 ^{ix}	0.91	2.10	2.981 (2)	162

Symmetry codes: (iii) $-x, -y+1, -z$; (iv) $-x+1, -y+1, -z$; (v) $x-1, y+1, z$; (vi) $x+1, y-1, z$; (vii) $-x+1, -y+1, -z+1$; (viii) $-x, -y+1, -z+1$; (ix) $x, y-1, z$.