

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

A new Mo^{VI} Schiff base complex: methanol[*N'*-(3-methoxy-2-oxidobenzylidene)benzohydrazidato]dioxido-molybdenum(VI)

 Iran Sheikhshoae,^{a*} Vratislav Langer^b and Seyed Ali Yasrebi^a
^aDepartment of Chemistry, Shahid Bahonar University, Kerman, Iran, and

^bEnvironmental Inorganic Chemistry, Department of Chemical and Biological Engineering, Chalmers University of Technology, SE-41296 Göteborg, Sweden

Correspondence e-mail: i_shoae@yahoo.com


Received 20 May 2011; accepted 26 May 2011

 Key indicators: single-crystal X-ray study; *T* = 173 K; mean $\sigma(\text{C}-\text{C}) = 0.004 \text{ \AA}$; *R* factor = 0.038; *wR* factor = 0.104; data-to-parameter ratio = 25.6.

In the title benzilidene Schiff base molybdenum(VI) complex, $[\text{Mo}(\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_3)\text{O}_2(\text{CH}_3\text{OH})]$, the Mo^{VI} ion is coordinated by two oxide O atoms and by two O atoms and one N atom of the tridentate *N'*-(3-methoxy-2-oxidobenzylidene)benzohydrazidate (*L*) Schiff base ligand. The methanol O atom completes the distorted octahedral configuration of the Mo^{VI} atom. Strong O—H···N hydrogen bonds form a *C*(5) chain around a *Z*₁ screw axis. Weak C—H—O hydrogen bonds are also present.

Related literature

For general background, see: Alizadeh *et al.* (1999); Ambroziak *et al.* (2004); Archer & Wang (1990); Bagherzadeh & Amini (2009); Bagherzadeh *et al.* (2008); Bhatia *et al.* (1981); Bindlish *et al.* (1978); Blake *et al.* (1995); Chang *et al.* (1998); Costamagna *et al.* (1992); Dhar & Taploo (1982); Hatefi *et al.* (2009); Holm (1990); Jalali-Heravi *et al.* (1999); Johnson *et al.* (1996); Maurya *et al.* (1997); Sheikhshoae & Fabian (2009); Yamada (1999). For details of the synthesis, see: Perrin *et al.* (1990). For related structures, see: Dinda *et al.* (2006); Glowiak *et al.* (2003); Liimatainen *et al.* (2000); Monadi *et al.* (2009); Niaz *et al.* (2010); Pramaniky *et al.* (2007); Rao *et al.* (1999); Rezaeifard *et al.* (2010); Saeednia *et al.* (2009); Sheikhshoae *et al.* (2009); Vrdoljak *et al.* (2010).



Experimental

Crystal data

 $[\text{Mo}(\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_3)\text{O}_2(\text{CH}_3\text{O})]$
 $M_r = 428.25$

 Monoclinic, *C*2/*c*
 $a = 29.400 (13) \text{ \AA}$
 $b = 8.553 (4) \text{ \AA}$
 $c = 14.391 (6) \text{ \AA}$
 $\beta = 112.993 (8)^\circ$
 $V = 3331 (2) \text{ \AA}^3$
 $Z = 8$

 Mo *K* α radiation

 $\mu = 0.82 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
 $0.58 \times 0.54 \times 0.46 \text{ mm}$

Data collection

 Bruker SMART CCD
 diffractometer

 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 2003)

 $T_{\min} = 0.368$, $T_{\max} = 0.703$

27788 measured reflections

5867 independent reflections

 4401 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.104$
 $S = 1.01$

5867 reflections

229 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 1.38 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.80 \text{ e \AA}^{-3}$
Table 1

 Selected geometric parameters (\AA , $^\circ$).

Mo1—O1	2.0281 (19)	Mo1—O5	1.7093 (19)
Mo1—O2	1.9391 (17)	Mo1—N1	2.248 (2)
Mo1—O4	1.7096 (18)	Mo1—O1M	2.3374 (18)

Table 2

 Hydrogen-bond geometry (\AA , $^\circ$).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1M—H1M···N2 ⁱ	0.84	1.87	2.700 (3)	173
C1M—H1BM···O1 ⁱⁱ	0.98	2.58	3.402 (3)	141
C6—H6···O4 ⁱⁱⁱ	0.95	2.53	3.450 (3)	162
C11—H11···O1	0.95	2.44	2.767 (3)	100
C15—H15···O1M ⁱⁱ	0.95	2.54	3.427 (3)	156

 Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT* and *SADABS* (Sheldrick, 2003); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

This work was supported by the Shahid Bahonar University of Kerman.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2692).

References

- Alizadeh, N., Ershad, S., Naeimi, H., Sharghi, H. & Shamsipur, M. (1999). *Pol. J. Chem.* **73**, 915–925.
 Ambroziak, K., Pelech, R., Milchert, E., Dziembowska, T. & Rozwadowski, Z. (2004). *J. Mol. Catal. A*, **211**, 9–16.
 Archer, R. D. & Wang, B. (1990). *Inorg. Chem.* **29**, 39–43.

- Bagherzadeh, M. & Amini, M. (2009). *Inorg. Chem. Commun.* **12**, 21–25.
- Bagherzadeh, M., Tahsini, L., Latifi, R., Ellern, A. & Woo, L. K. (2008). *Inorg. Chim. Acta*, **361**, 2019–2024.
- Bhatia, S. C., Bindlish, J. M., Saini, A. R. & Jain, P. C. (1981). *J. Chem. Soc. Dalton Trans.* pp. 1773–1779.
- Bindlish, J. M., Bhatia, S. C., Gautam, P. & Jain, P. C. (1978). *Indian J. Chem. Sect. A*, **16**, 279–282.
- Blake, A. B., Chipperfield, J. R., Hussain, W., Paschke, R. & Sinn, E. (1995). *Inorg. Chem.* **34**, 1125–1129.
- Brandenburg, K. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2003). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chang, S., Jones, L., Wang, C., Henling, L. M. & Grubbs, R. H. (1998). *Organometallics*, **17**, 3460–3465.
- Costamagna, J., Vargas, J., Latorre, R., Alvarado, A. & Mena, G. (1992). *Coord. Chem. Rev.* **119**, 67–88.
- Dhar, D. N. & Taploo, C. L. (1982). *J. Sci. Ind. Res.* **41**, 501–506.
- Dinda, R., Ghosh, S., Falvello, L. R., Tomas, M. & Mak, T. C. W. (2006). *Polyhedron*, **25**, 2375–2382.
- Glowiak, T., Jerzykiewicz, L., Sobczak, J. M. & Ziolkowski, J. J. (2003). *Inorg. Chim. Acta*, **356**, 387–392.
- Hatefi, M., Moghadam, M., Sheikhshoaei, I., Mirkhani, V., Tangestaninejad, S., Mohammadpoor-Baltork, I. & Kargar, H. (2009). *Appl. Catal. A*, **370**, 66–71.
- Holm, R. H. (1990). *Coord. Chem. Rev.* **100**, 183–221.
- Jalali-Heravi, M., Khandar, A. A. & Sheikhshoaei, I. (1999). *Spectrochim. Acta Part A*, **55**, 2537–2544.
- Johnson, C. P., Atwood, J. L., Steed, J. W., Bauer, C. B. & Rogers, R. D. (1996). *Inorg. Chem.* **35**, 2602–2610.
- Liimatainen, J., Lehtonen, A. & Sillanpaa, R. (2000). *Polyhedron*, **19**, 1133–1138.
- Maurya, R. C., Mishra, D. D., Rao, N. S., Verma, R. & Rao, N. N. (1997). *Indian J. Chem. Sect A*, **36**, 599–601.
- Monadi, N., Sheikhshoaei, I., Rezaeifard, A. & Stoeckli-Evans, H. (2009). *Acta Cryst. E* **65**, m1124–m1125.
- Niaz, M., Iran, S. & Abdolreza, R. (2010). *Acta Cryst. E* **66**, m202.
- Perrin, D. D., Armarego, W. L. & Perrin, D. R. (1990). *Purification of Laboratory Chemicals*, 2nd ed. New York: Pergamon.
- Pramaniky, N. R., Ghosh, S., Raychaudhuri, T. K., Chaudhuri, S., Drew, M. G. B. & Mandal, S. S. S. (2007). *J. Coord. Chem.* **60**, 2177–2190.
- Rao, S. N., Munshi, K. N., Rao, N. N., Bhadbhade, M. M. & Suresh, E. (1999). *Polyhedron*, **18**, 2491–2497.
- Rezaeifard, A., Sheikhshoaei, I., Monadi, N. & Stoeckli-Evans, H. (2010). *Eur. J. Inorg. Chem.* pp. 799–806.
- Saeednia, S., Sheikhshoaei, I. & Stoeckli-Evans, H. (2009). *Acta Cryst. E* **65**, m1591.
- Sheikhshoaei, I. & Fabian, W. M. F. (2009). *Curr. Org. Chem.* **13**, 149–171.
- Sheikhshoaei, I., Rezaeifard, A., Monadi, N. & Kaafi, S. (2009). *Polyhedron*, **28**, 733–738.
- Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Vrdoljak, V., Dilovic, L., Rubcic, M., Pavelic, S. K., Kralj, M., Matkovic-Catogovic, D., Piantanida, I., Novak, P., Rozman, A. & Cindric, M. (2010). *Eur. J. Med. Chem.* **45**, 38–48.
- Yamada, S. (1999). *Coord. Chem. Rev.* **190–192**, 537–555.

supporting information

Acta Cryst. (2011). E67, m839–m840 [doi:10.1107/S1600536811020101]

A new Mo^{VI} Schiff base complex: methanol[*N'*-(3-methoxy-2-oxidobenzylidene)benzohydrazidato]dioxidomolybdenum(VI)

Iran Sheikhshoae, Vratislav Langer and Seyed Ali Yasrebi

S1. Comment

The condensation products of primary amines and aldehydes or ketones ($RCH=NR'$), where R and R' represent alkyl and/or aryl constituents), are called Schiff bases (Dhar & Taploo, 1982); they play an important role in inorganic chemistry (Blake *et al.*, 1995; Johnson *et al.*, 1996; Alizadeh *et al.*, 1999) as they easily form stable complexes with most transition metal ions. Transition metal compounds containing the Schiff base ligands have been of interest for many years (Yamada, 1999; Chang *et al.*, 1998; Archer & Wang, 1990; Sheikhshoae & Fabian, 2009; Jalali-Heravi *et al.*, 1999). Many complexes play an important role in the developing of coordination chemistry related to catalytic (Holm, 1990; Rao *et al.*, 1999; Bagherzadeh *et al.*, 2008; Ambroziak *et al.*, 2004; Bagherzadeh & Amini, 2009; Hatefi *et al.*, 2009) and enzymatic reactions (Bindlish *et al.*, 1978; Bhatia *et al.*, 1981; Costamagna *et al.*, 1992). High valent metal oxo species have demonstrated the ability to catalyze the oxidation of a variety of organic substrates, *via* homogeneous as well as heterogeneous routes (Holm, 1990; Maurya *et al.*, 1997). In particular, the Mo-catalyzed alkene to epoxide conversion has received most attention. In the course of our studies on transition metal Schiff base complexes (Sheikhshoae *et al.*, 2009; Monadi *et al.*, 2009), we have synthesized a dioxo molybdenum complex with a tridentate ligand (L) and the crystal structure of the title complex (I) is presented here.

The molecular structure of (I) is shown in Fig. 1. and the selected bond distances and angles are given in Table 1. The calculated bond valence by *PLATON* (version 91110: Spek, 2009) for Mo atom is 5.89, *i.e.* it exhibits the oxidation state +VI. Mo atom is surrounded by two O atoms and one N atom of the tridentate Schiff base ligand 3-methoxy-salicylidenbenzoyl hydrazine in a distorted octahedral configuration. The Mo—O distances of the oxo ligands (O4 and O5) are significantly shorter [average 1.7094 (18) Å] than the corresponding distances to the O atoms of the tridentate ligands (O1 and O2) [average 1.98 (6) Å]. The Mo—N distance is longer [2.248 (2) Å] and the distance to methanol O atom is longest.

The aromatic ring C1/C2/C3/C4/C5/C6 is slightly distorted ($\chi^2 = 56.6$), while the phenyl ring C10/C11/C12/C13/C14/C15 is planar ($\chi^2 = 4.7$); the dihedral angle between them is 23.09 (12)°.

There are strong hydrogen bonds present between the hydroxy group of methanol and N2ⁱ atom in an adjacent complex [symmetry code: (i): $-x + 1/2, y + 1/2, -z + 1/2$] forming thus a C(5) chain around 2₁ screw axis, see Fig.2. There are also weak hydrogen bonds of the type in the C—H—O hydrogen bonds present, see Table 2. Fig. 3 shows the packing of the complexes in the unit cell, but there are no π - π interactions present.

Comparison with other similar structures is presented in Table 3. In all complexes molybdenum atom is coordinated by two oxo oxygen atoms, one nitrogen atom of C=N group and one oxygen atom from solvent (methanol or ethanol). In all these complexes, the longest bond from the solvent coordination indicates the most labile site available for substitution. In addition, in all the investigated complexes the bonds between molybdenum atom and nitrogen atom of imine group

(C=N) are long (average: 2.26 (2) Å). Two oxo oxygen atoms are almost in *cis* position towards each other in all the cases [the average of the O=Mo=O angles: 106.1 (5)°]. The angles of N—Mo=O in the complexes indicate that one of the oxo oxygen atoms is in *cis* and the other one is in *trans* position toward nitrogen atom of imine group. The coordination geometry around Mo atoms is highly distorted octahedral in [MoO₂(*L*) (solvent)] complexes, where (*L*) is a tridentate ligand containing an imine group and MeOH or EtOH as a solvent. Table 4 provides a comparison of important frequencies in the IR spectra of (I) and some dioxo-molybdenum(VI) complexes.

S2. Experimental

All reagents were used as supplied by Fluka and Merck without further purification. Solvents used for the reaction were purified and dried by conventional methods (Perrin *et al.*, 1990).

NMR spectra were obtained on a BRUKER AVANCE DRX500 (500 MHz) spectrometer. Proton chemical shifts δ are reported in p.p.m. relative to an internal standard of Me₄Si. Elemental analyses were performed by using Heraeus CHN-O-RAPID elemental analyzer. IR spectra were recorded in KBr pellets using Shimadzu 435 spectrophotometer.

Synthesis of ligand (*L*): To a solution of 0.136 g (0.001 mol) benzohydrazide in 20 ml ethanol was added a solution of 0.152 g (0.001 mol) 3-methoxysalicylaldehyde in 10 ml ethanol, and refluxed for 5 h. The solvent was evaporated on a rotatory evaporator and the solid dissolved in 10 ml ethanol and filtered off. The filtrate was left overnight to give yellow crystals. Yield: 202 mg (75%). *Anal. Calc.* for C₁₅H₁₄N₂O₃: C, 66.66; H, 5.18; N, 10.37. Found: C, 66.79; H, 5.20; N, 10.39%. IR (KBr, cm⁻¹) ν_{\max} 1652.9 (s, C=O), 1614 (m, C=N). ¹H NMR: (d₆-DMSO): δ 3.81 (s, 3H, methoxy group), δ 6.84–7.96 (m, 8H, ArH), δ 8.671 (s, 1H, —CH=N— group), δ 11.02 (s, 1H, OH group), δ 12.09 (s, 1H, NH group).

Synthesis of the title metal complex (I) [Mo O₂(*L*)(CH₃OH)]: The title dioxomolybdenum(VI) complex was prepared by mixing MoO₂(acac)₂ (acac=acetylacetonate) with the ligand in a 1:1 molar ratio using 30 ml dry methanol as a solvent, followed by refluxing the solution for 4 h. Deeply orange crystals were collected by filtration and dried in the room temperature, yield: 83%. ¹H NMR: δ 3.80 (s, 3H, methoxy group), δ 7.006–8.01 (m, 8 H, Ar H), δ 8.92 (s, 1H, —CH=N— group), δ 4.08–4.11 (s, 1H, OH methanol group); δ 3.16 (d, 3H CH₃ in methanol). IR spectra are presented in Table 4 [frequencies ν (N—H) and ν (C=O) were not observed].

S3. Refinement

Aromatic hydrogen atoms were refined isotropically with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, and their positions were constrained to an ideal geometry using an appropriate riding model, (C—H = 0.95 Å). For methyl groups, O—C—H angles (109.5°) were kept fixed, while the torsion angle was allowed to refine with the starting positions based on the circular Fourier synthesis averaged using the local 3-fold axis with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ and a constrained C—H distance of 0.98 Å was applied. For hydroxy group, the O—H distance (0.84 Å) and C—O—H angle (109.5°) were kept fixed, while the torsion angle was allowed to refine with the starting position based on the maximum on the circular Fourier synthesis, $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.





Figure 1

The atom numbering scheme for (I), with atomic displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

The hydrogen-pattern (dashed lines) in (I). C(5) chains are formed in the b-direction. Symmetry code: (i): $-x + 1/2, y + 1/2, -z + 1/2$.




Figure 3

Content of the unit cell for (I) in projection along the *b* axis.

methanol[*N'*-(3-methoxy-2-oxidobenzylidene)benzohydrazidato]dioxidomolybdenum(VI)

Crystal data

[Mo(C₁₅H₁₂N₂O₃)O₂(CH₄O)]

M_r = 428.25

Monoclinic, *C2/c*

Hall symbol: -*C* 2yc

a = 29.400 (13) Å

b = 8.553 (4) Å

c = 14.391 (6) Å

β = 112.993 (8)°

V = 3331 (2) Å³

Z = 8

F(000) = 1728

D_x = 1.708 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 5066 reflections

θ = 2.5–32.3°

μ = 0.82 mm⁻¹

T = 173 K

Block, orange

0.58 × 0.54 × 0.46 mm

Data collection

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2003)

T_{min} = 0.368, *T_{max}* = 0.703

27788 measured reflections

5867 independent reflections

4401 reflections with *I* > 2σ(*I*)

R_{int} = 0.065

θ_{max} = 32.8°, θ_{min} = 2.5°

h = -43→43

k = -12→12

l = -21→21

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.038

wR(*F*²) = 0.104

S = 1.01

5867 reflections

229 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0566*P*)² + 2.343*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 1.38 e Å⁻³

Δρ_{min} = -1.80 e Å⁻³

Special details

Experimental. Data were collected at 173 K using a Siemens *SMART* CCD diffractometer equipped with LT-2 A cooling device. A full sphere of reciprocal space was scanned by 0.3° steps in ω with a crystal-to-detector distance of 3.97 cm, 1 second per frame. Preliminary orientation matrix was obtained from the first 100 frames using *SMART* (Bruker, 2003). The collected frames were integrated using the preliminary orientation matrix which was updated every 100 frames. Final cell parameters were obtained by refinement on the position of 5066 reflections with $I > 10\sigma(I)$ after integration of all the frames data using *SAINTE* (Bruker, 2003).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mo1	0.169567 (7)	0.55438 (2)	0.093970 (13)	0.01640 (7)
O1	0.23804 (6)	0.54087 (18)	0.09191 (12)	0.0191 (3)
O2	0.11949 (6)	0.47242 (19)	0.13716 (13)	0.0209 (3)
O3	0.02944 (7)	0.4772 (2)	0.13247 (15)	0.0299 (4)
O4	0.17354 (6)	0.74754 (19)	0.12674 (13)	0.0244 (4)
O5	0.13736 (7)	0.5485 (2)	-0.03385 (13)	0.0254 (4)
N1	0.19324 (7)	0.3025 (2)	0.11307 (13)	0.0163 (3)
N2	0.24201 (7)	0.2753 (2)	0.12187 (14)	0.0176 (4)
C1	0.11885 (8)	0.1902 (3)	0.12040 (16)	0.0185 (4)
C2	0.09645 (8)	0.3331 (3)	0.12693 (16)	0.0187 (4)
C3	0.04804 (9)	0.3332 (3)	0.12641 (17)	0.0226 (4)
C4	0.02337 (9)	0.1919 (3)	0.11911 (19)	0.0285 (5)
H4	-0.0093	0.1918	0.1170	0.034*
C5	0.04618 (10)	0.0501 (3)	0.1148 (2)	0.0282 (5)
H5	0.0291	-0.0456	0.1111	0.034*
C6	0.09347 (9)	0.0485 (3)	0.11585 (19)	0.0236 (5)
H6	0.1089	-0.0482	0.1135	0.028*
C7	0.16837 (8)	0.1801 (3)	0.12059 (16)	0.0189 (4)
H7	0.1832	0.0799	0.1264	0.023*
C8	-0.02257 (10)	0.4859 (4)	0.1098 (2)	0.0357 (6)
H8A	-0.0409	0.4339	0.0452	0.054*
H8B	-0.0327	0.5958	0.1053	0.054*
H8C	-0.0297	0.4340	0.1633	0.054*
C9	0.26226 (8)	0.4068 (3)	0.11017 (16)	0.0171 (4)
C10	0.31418 (8)	0.4117 (3)	0.11898 (17)	0.0186 (4)
C11	0.33256 (9)	0.5503 (3)	0.09429 (18)	0.0212 (4)
H11	0.3115	0.6383	0.0706	0.025*
C12	0.38163 (10)	0.5592 (3)	0.1044 (2)	0.0265 (5)
H12	0.3938	0.6532	0.0874	0.032*
C13	0.41308 (10)	0.4306 (3)	0.1396 (2)	0.0279 (5)

H13	0.4466	0.4370	0.1470	0.033*
C14	0.39471 (9)	0.2927 (3)	0.1637 (2)	0.0277 (5)
H14	0.4158	0.2048	0.1873	0.033*
C15	0.34575 (9)	0.2829 (3)	0.15337 (18)	0.0228 (5)
H15	0.3336	0.1883	0.1697	0.027*
O1M	0.21855 (6)	0.51974 (19)	0.26524 (12)	0.0200 (3)
H1M	0.2288	0.6039	0.2964	0.030*
C1M	0.21418 (10)	0.4038 (3)	0.33394 (17)	0.0240 (5)
H1AM	0.2338	0.4359	0.4035	0.036*
H1BM	0.2263	0.3030	0.3206	0.036*
H1CM	0.1794	0.3935	0.3244	0.036*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mo1	0.01570 (10)	0.01803 (10)	0.01522 (9)	0.00265 (7)	0.00577 (7)	0.00205 (7)
O1	0.0194 (8)	0.0192 (8)	0.0205 (7)	0.0034 (6)	0.0098 (6)	0.0041 (6)
O2	0.0171 (8)	0.0222 (8)	0.0249 (8)	0.0010 (6)	0.0099 (7)	-0.0004 (6)
O3	0.0191 (9)	0.0358 (10)	0.0363 (10)	0.0057 (7)	0.0126 (8)	0.0020 (8)
O4	0.0271 (9)	0.0205 (8)	0.0273 (9)	0.0058 (7)	0.0126 (7)	0.0045 (7)
O5	0.0224 (9)	0.0332 (10)	0.0186 (8)	0.0054 (7)	0.0058 (7)	0.0033 (7)
N1	0.0156 (9)	0.0190 (8)	0.0143 (8)	0.0003 (7)	0.0058 (7)	-0.0004 (6)
N2	0.0152 (8)	0.0213 (9)	0.0167 (8)	0.0019 (7)	0.0066 (7)	-0.0006 (7)
C1	0.0159 (10)	0.0238 (11)	0.0145 (9)	-0.0014 (8)	0.0046 (8)	-0.0003 (8)
C2	0.0159 (10)	0.0254 (11)	0.0138 (9)	0.0005 (8)	0.0047 (8)	0.0019 (8)
C3	0.0163 (10)	0.0308 (12)	0.0197 (10)	0.0008 (9)	0.0060 (8)	0.0007 (9)
C4	0.0172 (11)	0.0400 (14)	0.0291 (12)	-0.0050 (10)	0.0099 (10)	0.0022 (11)
C5	0.0220 (12)	0.0321 (13)	0.0308 (13)	-0.0080 (10)	0.0107 (10)	-0.0001 (10)
C6	0.0212 (11)	0.0248 (11)	0.0234 (11)	-0.0041 (9)	0.0074 (9)	-0.0007 (9)
C7	0.0177 (10)	0.0217 (10)	0.0163 (9)	0.0008 (8)	0.0055 (8)	0.0000 (8)
C8	0.0194 (12)	0.0552 (18)	0.0344 (14)	0.0113 (12)	0.0125 (11)	0.0076 (13)
C9	0.0184 (10)	0.0207 (10)	0.0115 (8)	0.0011 (8)	0.0050 (8)	-0.0002 (7)
C10	0.0181 (10)	0.0236 (11)	0.0154 (9)	-0.0003 (8)	0.0079 (8)	-0.0022 (8)
C11	0.0208 (11)	0.0233 (11)	0.0209 (10)	0.0015 (9)	0.0095 (9)	0.0031 (9)
C12	0.0248 (12)	0.0288 (12)	0.0285 (12)	-0.0037 (10)	0.0133 (10)	0.0024 (10)
C13	0.0198 (11)	0.0356 (14)	0.0304 (13)	-0.0017 (10)	0.0120 (10)	-0.0005 (10)
C14	0.0214 (12)	0.0304 (13)	0.0335 (13)	0.0034 (10)	0.0130 (10)	0.0012 (10)
C15	0.0205 (11)	0.0228 (11)	0.0280 (12)	0.0021 (9)	0.0128 (9)	0.0023 (9)
O1M	0.0241 (8)	0.0205 (8)	0.0152 (7)	-0.0036 (6)	0.0076 (6)	-0.0017 (6)
C1M	0.0307 (13)	0.0247 (11)	0.0149 (10)	0.0001 (9)	0.0071 (9)	0.0027 (8)

Geometric parameters (Å, °)

Mo1—O1	2.0281 (19)	C6—H6	0.9500
Mo1—O2	1.9391 (17)	C7—H7	0.9500
Mo1—O4	1.7096 (18)	C8—H8A	0.9800
Mo1—O5	1.7093 (19)	C8—H8B	0.9800
Mo1—N1	2.248 (2)	C8—H8C	0.9800

Mo1—O1M	2.3374 (18)	C9—C10	1.482 (3)
O1—C9	1.321 (3)	C10—C15	1.400 (3)
O2—C2	1.350 (3)	C10—C11	1.404 (3)
O3—C3	1.364 (3)	C11—C12	1.394 (4)
O3—C8	1.435 (3)	C11—H11	0.9500
N1—C7	1.306 (3)	C12—C13	1.399 (4)
N1—N2	1.409 (3)	C12—H12	0.9500
N2—C9	1.313 (3)	C13—C14	1.396 (4)
C1—C2	1.409 (3)	C13—H13	0.9500
C1—C6	1.411 (3)	C14—C15	1.391 (3)
C1—C7	1.457 (3)	C14—H14	0.9500
C2—C3	1.420 (3)	C15—H15	0.9500
C3—C4	1.392 (4)	O1M—C1M	1.441 (3)
C4—C5	1.399 (4)	O1M—H1M	0.8400
C4—H4	0.9500	C1M—H1AM	0.9800
C5—C6	1.384 (4)	C1M—H1BM	0.9800
C5—H5	0.9500	C1M—H1CM	0.9800
O4—Mo1—O5	105.93 (8)	N1—C7—H7	118.5
O4—Mo1—O2	103.86 (8)	C1—C7—H7	118.5
O5—Mo1—O2	99.42 (9)	O3—C8—H8A	109.5
O4—Mo1—O1	95.61 (7)	O3—C8—H8B	109.5
O5—Mo1—O1	96.80 (8)	H8A—C8—H8B	109.5
O2—Mo1—O1	150.03 (7)	O3—C8—H8C	109.5
O4—Mo1—N1	155.24 (8)	H8A—C8—H8C	109.5
O5—Mo1—N1	96.81 (7)	H8B—C8—H8C	109.5
O2—Mo1—N1	81.49 (7)	N2—C9—O1	122.2 (2)
O1—Mo1—N1	71.67 (6)	N2—C9—C10	121.1 (2)
C9—O1—Mo1	120.49 (14)	O1—C9—C10	116.7 (2)
C2—O2—Mo1	134.07 (14)	C15—C10—C11	119.1 (2)
C3—O3—C8	116.8 (2)	C15—C10—C9	121.7 (2)
C7—N1—N2	116.36 (18)	C11—C10—C9	119.2 (2)
C7—N1—Mo1	128.46 (15)	C12—C11—C10	120.3 (2)
N2—N1—Mo1	115.13 (13)	C12—C11—H11	119.9
C9—N2—N1	110.11 (18)	C10—C11—H11	119.9
C2—C1—C6	119.7 (2)	C11—C12—C13	120.4 (2)
C2—C1—C7	122.9 (2)	C11—C12—H12	119.8
C6—C1—C7	117.3 (2)	C13—C12—H12	119.8
O2—C2—C1	123.0 (2)	C14—C13—C12	119.3 (2)
O2—C2—C3	117.3 (2)	C14—C13—H13	120.4
C1—C2—C3	119.6 (2)	C12—C13—H13	120.4
O3—C3—C4	125.4 (2)	C15—C14—C13	120.6 (2)
O3—C3—C2	115.2 (2)	C15—C14—H14	119.7
C4—C3—C2	119.4 (2)	C13—C14—H14	119.7
C3—C4—C5	120.8 (2)	C14—C15—C10	120.4 (2)
C3—C4—H4	119.6	C14—C15—H15	119.8
C5—C4—H4	119.6	C10—C15—H15	119.8
C6—C5—C4	120.3 (2)	C1M—O1M—H1M	109.5

C6—C5—H5	119.9	O1M—C1M—H1AM	109.5
C4—C5—H5	119.9	O1M—C1M—H1BM	109.5
C5—C6—C1	120.2 (2)	H1AM—C1M—H1BM	109.5
C5—C6—H6	119.9	O1M—C1M—H1CM	109.5
C1—C6—H6	119.9	H1AM—C1M—H1CM	109.5
N1—C7—C1	123.0 (2)	H1BM—C1M—H1CM	109.5
O4—Mo1—O1—C9	152.56 (16)	O2—C2—C3—C4	-178.1 (2)
O5—Mo1—O1—C9	-100.61 (16)	C1—C2—C3—C4	-0.2 (3)
O2—Mo1—O1—C9	21.8 (2)	O3—C3—C4—C5	-179.1 (2)
N1—Mo1—O1—C9	-5.67 (15)	C2—C3—C4—C5	1.6 (4)
O4—Mo1—O2—C2	174.04 (19)	C3—C4—C5—C6	-1.2 (4)
O5—Mo1—O2—C2	64.9 (2)	C4—C5—C6—C1	-0.6 (4)
O1—Mo1—O2—C2	-56.9 (3)	C2—C1—C6—C5	1.9 (3)
N1—Mo1—O2—C2	-30.6 (2)	C7—C1—C6—C5	-179.6 (2)
O4—Mo1—N1—C7	120.7 (2)	N2—N1—C7—C1	175.46 (18)
O5—Mo1—N1—C7	-82.56 (19)	Mo1—N1—C7—C1	-1.8 (3)
O2—Mo1—N1—C7	15.99 (19)	C2—C1—C7—N1	-9.5 (3)
O1—Mo1—N1—C7	-177.5 (2)	C6—C1—C7—N1	172.0 (2)
O4—Mo1—N1—N2	-56.6 (2)	N1—N2—C9—O1	-0.6 (3)
O5—Mo1—N1—N2	100.14 (14)	N1—N2—C9—C10	178.14 (18)
O2—Mo1—N1—N2	-161.31 (14)	Mo1—O1—C9—N2	5.8 (3)
O1—Mo1—N1—N2	5.20 (13)	Mo1—O1—C9—C10	-173.01 (14)
C7—N1—N2—C9	178.27 (19)	N2—C9—C10—C15	-8.8 (3)
Mo1—N1—N2—C9	-4.1 (2)	O1—C9—C10—C15	170.0 (2)
Mo1—O2—C2—C1	30.2 (3)	N2—C9—C10—C11	172.5 (2)
Mo1—O2—C2—C3	-151.97 (17)	O1—C9—C10—C11	-8.6 (3)
C6—C1—C2—O2	176.2 (2)	C15—C10—C11—C12	-0.3 (3)
C7—C1—C2—O2	-2.2 (3)	C9—C10—C11—C12	178.3 (2)
C6—C1—C2—C3	-1.5 (3)	C10—C11—C12—C13	-0.2 (4)
C7—C1—C2—C3	-179.9 (2)	C11—C12—C13—C14	0.5 (4)
C8—O3—C3—C4	-12.9 (4)	C12—C13—C14—C15	-0.3 (4)
C8—O3—C3—C2	166.5 (2)	C13—C14—C15—C10	-0.3 (4)
O2—C2—C3—O3	2.5 (3)	C11—C10—C15—C14	0.5 (3)
C1—C2—C3—O3	-179.6 (2)	C9—C10—C15—C14	-178.1 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1M—H1M \cdots N2 ⁱ	0.84	1.87	2.700 (3)	173
C1M—H1BM \cdots O1 ⁱⁱ	0.98	2.58	3.402 (3)	141
C6—H6 \cdots O4 ⁱⁱⁱ	0.95	2.53	3.450 (3)	162
C11—H11 \cdots O1	0.95	2.44	2.767 (3)	100
C15—H15 \cdots O1M ⁱⁱ	0.95	2.54	3.427 (3)	156

Symmetry codes: (i) $-x+1/2, y+1/2, -z+1/2$; (ii) $-x+1/2, y-1/2, -z+1/2$; (iii) $x, y-1, z$.