

**{N-Methyl-N'-[1-(pyridin-2-yl)ethylidene]ethane-1,2-diamine- $\kappa^3 N,N',N''$ }-bis(thiocyanato- $\kappa N$ )zinc(II)**

Xian-Wen Li

Department of Chemistry and Chemical Engineering, Minjiang University, Fuzhou 350108, People's Republic of China  
Correspondence e-mail: xianwen\_li@126.com


Received 24 May 2011; accepted 25 May 2011

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C-C}) = 0.007\text{ \AA}$ ; disorder in main residue;  $R$  factor = 0.040;  $wR$  factor = 0.105; data-to-parameter ratio = 16.7.

In the title compound,  $[\text{Zn}(\text{NCS})_2(\text{C}_{10}\text{H}_{15}\text{N}_3)]$ , the Zn atom is five-coordinated by the three N-donor atoms of the Schiff base ligand and by two N atoms from two thiocyanate anions, forming a distorted  $\text{ZnN}_5$  trigonal-bipyramidal coordination geometry for the metal ion. The side chain of the ligand is disordered over two sets of sites in a 0.655 (12):0.345 (12) ratio. In the crystal, molecules are linked by  $\text{N-H}\cdots\text{S}$  hydrogen bonds, generating [100] chains.

## Related literature

For the biological activity of Schiff base compounds, see: Panneerselvam *et al.* (2005); Shi *et al.* (2007); Singh *et al.* (2006, 2007); Zhong *et al.* (2006). For the Schiff base complexes we reported previously, see: Li & Qiu (2008a,b).



## Experimental

### Crystal data

$[\text{Zn}(\text{NCS})_2(\text{C}_{10}\text{H}_{15}\text{N}_3)]$	$c = 14.3766 (6)\text{ \AA}$
$M_r = 358.78$	$\beta = 101.853 (2)^\circ$
Monoclinic, $P2_{1}/n$	$V = 1597.30 (11)\text{ \AA}^3$
$a = 7.6674 (3)\text{ \AA}$	$Z = 4$
$b = 14.8062 (5)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 1.80\text{ mm}^{-1}$   
 $T = 298\text{ K}$

$0.20 \times 0.20 \times 0.18\text{ mm}$

### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2004)  
 $T_{\min} = 0.715$ ,  $T_{\max} = 0.738$

9004 measured reflections  
3352 independent reflections  
2095 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.029$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.105$   
 $S = 1.02$   
3352 reflections  
201 parameters

12 restraints  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.46\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.46\text{ e \AA}^{-3}$

**Table 1**  
Selected bond lengths ( $\text{\AA}$ ).

Zn1—N5	1.974 (4)	Zn1—N3	2.163 (4)
Zn1—N4	1.986 (4)	Zn1—N1	2.195 (3)
Zn1—N2	2.088 (3)		

**Table 2**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3A $\cdots$ S1 <sup>i</sup>	0.91	2.66	3.551 (5)	165

Symmetry code: (i)  $x - 1, y, z$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We thank Minjiang University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5892).

## References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Li, X.-W. & Qiu, Y. (2008a). *Acta Cryst. E* **64**, m113.
- Li, X.-W. & Qiu, Y. (2008b). *Acta Cryst. E* **64**, m218.
- Panneerselvam, P., Nair, R. R., Vijayalakshmi, G., Subramanian, E. H. & Krishnan, S. (2005). *Eur. J. Med. Chem.* **40**, 225–229.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shi, L., Ge, H.-M., Tan, S.-H., Li, H.-Q., Song, Y.-C., Zhu, H.-L. & Tan, R.-X. (2007). *Eur. J. Med. Chem.* **42**, 558–564.
- Singh, K., Barwa, M. S. & Tyagi, P. (2006). *Eur. J. Med. Chem.* **41**, 147–153.
- Singh, K., Barwa, M. S. & Tyagi, P. (2007). *Eur. J. Med. Chem.* **42**, 394–402.
- Zhong, X., Yi, J., Sun, J., Wei, H.-L., Liu, W.-S. & Yu, K.-B. (2006). *Eur. J. Med. Chem.* **41**, 1090–1092.

# supporting information

*Acta Cryst.* (2011). E67, m829 [doi:10.1107/S1600536811019945]

## {N-Methyl-N'-[1-(pyridin-2-yl)ethylidene]ethane-1,2-diamine- $\kappa^3$ N,N',N''}bis-(thiocyanato- $\kappa$ N)zinc(II)

Xian-Wen Li

### S1. Comment

Schiff base compounds have been reported to have excellent biological activity (Shi *et al.*, 2007; Panneerselvam *et al.*, 2005). The metal complexes derived from the Schiff bases also have excellent biological activity (Singh *et al.*, 2006, 2007; Zhong *et al.*, 2006). As a continuation of our work on Schiff base complexes (Li & Qiu, 2008*a,b*), we report herein the crystal structure of the title zinc complex, (I).

In the title mononuclear zinc(II) complex, the Zn atom is five-coordinated by the three donor atoms (N1, N2, and N3) of the Schiff baes ligand, and two N atoms from two thiocyanate ligands, forming a slightly distorted trigonal-bipyramidal geometry (Fig. 1). The coordinate bond values (Table 1) are within normal ranges.


### S2. Experimental

The title compound was obtained by the reaction of equimolar quantities (0.1 mmol each) of 2-acetylpyridine, *N*-methyl-ethane-1,2-diamine, sodium thiocyanate, and zinc acetate in ethanol. Colorless blocks of (I) were obtained by the slow evaporation of the filtrate in air.

### S3. Refinement

H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93–0.97 Å and with  $U_{\text{iso}}(\text{H}) = 1.2$  (1.5 for methyl groups) times  $U_{\text{eq}}(\text{C})$ .

The amino H atoms were located in a difference map and refined with N—H distance restrained to 0.90 (1) Å. The remaining H atoms were positioned geometrically (C—H = 0.93–0.97 Å, N—H = 0.91 Å) and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and 1.5 $U_{\text{eq}}$ (methyl C). The C9—N3—C10 moiety is disordered over two sites, with occupancies of 0.655 (3) and 0.345 (3).

**Figure 1**

The molecular structure of the title compound, showing 30% probability displacement ellipsoids. Only the major component of the disordered group is shown.

### {N-Methyl-N'-[1-(pyridin-2-yl)ethylidene]ethane-1,2-diamine- $\kappa^3N,N',N''$ }bis(thiocyanato- $\kappa N$ )zinc(II)

#### Crystal data

$[Zn(NCS)_2(C_{10}H_{15}N_3)]$

$M_r = 358.78$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 7.6674 (3) \text{ \AA}$

$b = 14.8062 (5) \text{ \AA}$

$c = 14.3766 (6) \text{ \AA}$

$\beta = 101.853 (2)^\circ$

$V = 1597.30 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 736$

$D_x = 1.492 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2372 reflections

$\theta = 2.6\text{--}25.1^\circ$

$\mu = 1.80 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Block, colorless

$0.20 \times 0.20 \times 0.18 \text{ mm}$

#### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 2004)

$T_{\min} = 0.715$ ,  $T_{\max} = 0.738$

9004 measured reflections

3352 independent reflections

2095 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$   
 $\theta_{\text{max}} = 26.7^\circ, \theta_{\text{min}} = 2.0^\circ$   
 $h = -9 \rightarrow 9$

$k = -17 \rightarrow 18$   
 $l = -18 \rightarrow 11$

### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.105$   
 $S = 1.02$   
3352 reflections  
201 parameters  
12 restraints  
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0323P)^2 + 1.952P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.46 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.46 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.72087 (6)	-0.07353 (3)	0.71636 (3)	0.05504 (17)	
S1	1.28221 (17)	-0.15205 (10)	0.89256 (10)	0.0835 (4)	
S2	0.3366 (2)	-0.32564 (11)	0.66375 (10)	0.1063 (6)	
N1	0.7424 (4)	-0.0555 (2)	0.5676 (2)	0.0524 (8)	
N2	0.6777 (4)	0.0641 (2)	0.6890 (2)	0.0559 (8)	
N4	0.9697 (5)	-0.1140 (3)	0.7651 (3)	0.0760 (11)	
N5	0.5583 (5)	-0.1782 (2)	0.6923 (2)	0.0684 (10)	
C1	0.7390 (5)	0.0309 (3)	0.5392 (3)	0.0515 (9)	
C2	0.7683 (5)	0.0539 (3)	0.4504 (3)	0.0628 (11)	
H2	0.7670	0.1142	0.4319	0.075*	
C3	0.7993 (6)	-0.0131 (4)	0.3901 (3)	0.0708 (13)	
H3	0.8193	0.0015	0.3303	0.085*	
C4	0.8009 (6)	-0.1007 (4)	0.4174 (3)	0.0706 (13)	
H4	0.8207	-0.1469	0.3770	0.085*	
C5	0.7719 (6)	-0.1192 (3)	0.5074 (3)	0.0629 (11)	
H5	0.7731	-0.1792	0.5268	0.075*	
C6	0.7014 (5)	0.0970 (3)	0.6107 (3)	0.0577 (10)	
C7	0.6971 (7)	0.1958 (3)	0.5876 (4)	0.0877 (15)	
H7A	0.6442	0.2284	0.6325	0.132*	
H7B	0.6279	0.2052	0.5247	0.132*	
H7C	0.8164	0.2172	0.5907	0.132*	
C11	1.1008 (6)	-0.1300 (3)	0.8180 (3)	0.0573 (10)	

C12	0.4653 (6)	-0.2395 (3)	0.6815 (3)	0.0586 (10)	
C8	0.6362 (7)	0.1164 (3)	0.7662 (3)	0.0810 (14)	0.655 (12)
H8A	0.5502	0.1631	0.7417	0.097*	0.655 (12)
H8B	0.7431	0.1451	0.8016	0.097*	0.655 (12)
C9	0.5577 (12)	0.0508 (5)	0.8318 (5)	0.074 (3)	0.655 (12)
H9A	0.5582	0.0803	0.8921	0.089*	0.655 (12)
H9B	0.4349	0.0371	0.8026	0.089*	0.655 (12)
C10	0.7978 (14)	-0.0366 (8)	0.9293 (6)	0.089 (3)	0.655 (12)
H10A	0.7586	-0.0160	0.9849	0.133*	0.655 (12)
H10B	0.8964	-0.0004	0.9196	0.133*	0.655 (12)
H10C	0.8346	-0.0986	0.9377	0.133*	0.655 (12)
N3	0.6540 (6)	-0.0290 (3)	0.8480 (2)	0.0814 (12)	0.655 (12)
H3A	0.5729	-0.0703	0.8591	0.098*	0.655 (12)
C8'	0.6362 (7)	0.1164 (3)	0.7662 (3)	0.0810 (14)	0.345 (12)
H8'A	0.5082	0.1233	0.7582	0.097*	0.345 (12)
H8'B	0.6893	0.1760	0.7673	0.097*	0.345 (12)
C9'	0.709 (2)	0.0678 (7)	0.8562 (7)	0.097 (7)	0.345 (12)
H9'A	0.6658	0.0959	0.9081	0.116*	0.345 (12)
H9'B	0.8382	0.0719	0.8702	0.116*	0.345 (12)
C10'	0.723 (2)	-0.0647 (13)	0.9393 (8)	0.066 (5)	0.345 (12)
H10D	0.6699	-0.0338	0.9854	0.099*	0.345 (12)
H10E	0.8497	-0.0565	0.9546	0.099*	0.345 (12)
H10F	0.6955	-0.1279	0.9400	0.099*	0.345 (12)
N3'	0.6540 (6)	-0.0290 (3)	0.8480 (2)	0.0814 (12)	0.345 (12)
H3'A	0.5331	-0.0302	0.8394	0.098*	0.345 (12)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.0644 (3)	0.0492 (3)	0.0518 (3)	0.0000 (2)	0.0128 (2)	0.0037 (2)
S1	0.0678 (8)	0.0948 (10)	0.0845 (9)	0.0061 (7)	0.0077 (6)	-0.0070 (7)
S2	0.1434 (14)	0.0975 (11)	0.0869 (9)	-0.0621 (10)	0.0447 (9)	-0.0190 (8)
N1	0.062 (2)	0.0456 (19)	0.0502 (18)	-0.0062 (15)	0.0126 (15)	-0.0006 (15)
N2	0.061 (2)	0.0461 (18)	0.061 (2)	0.0029 (16)	0.0128 (16)	-0.0033 (16)
N4	0.074 (3)	0.082 (3)	0.073 (3)	0.008 (2)	0.018 (2)	0.021 (2)
N5	0.082 (3)	0.055 (2)	0.069 (2)	-0.007 (2)	0.017 (2)	0.0074 (18)
C1	0.042 (2)	0.057 (2)	0.053 (2)	-0.0039 (18)	0.0023 (17)	0.0080 (19)
C2	0.055 (3)	0.071 (3)	0.059 (3)	-0.008 (2)	0.003 (2)	0.020 (2)
C3	0.064 (3)	0.100 (4)	0.048 (2)	-0.015 (3)	0.010 (2)	0.009 (3)
C4	0.068 (3)	0.092 (4)	0.054 (3)	-0.010 (3)	0.018 (2)	-0.012 (2)
C5	0.072 (3)	0.056 (3)	0.064 (3)	-0.005 (2)	0.022 (2)	-0.002 (2)
C6	0.054 (2)	0.044 (2)	0.070 (3)	-0.0030 (18)	0.002 (2)	0.004 (2)
C7	0.114 (4)	0.048 (3)	0.099 (4)	-0.001 (3)	0.015 (3)	0.008 (3)
C11	0.065 (3)	0.051 (2)	0.060 (3)	-0.005 (2)	0.024 (2)	0.003 (2)
C12	0.075 (3)	0.059 (3)	0.045 (2)	-0.002 (2)	0.022 (2)	0.007 (2)
C8	0.097 (4)	0.060 (3)	0.089 (4)	-0.004 (3)	0.025 (3)	-0.019 (3)
C9	0.081 (6)	0.080 (6)	0.062 (4)	0.033 (4)	0.018 (4)	-0.005 (4)
C10	0.111 (7)	0.092 (7)	0.059 (5)	0.030 (5)	0.010 (5)	-0.003 (4)

N3	0.092 (3)	0.097 (3)	0.057 (2)	-0.012 (3)	0.020 (2)	-0.009 (2)
C8'	0.097 (4)	0.060 (3)	0.089 (4)	-0.004 (3)	0.025 (3)	-0.019 (3)
C9'	0.096 (14)	0.151 (18)	0.044 (8)	-0.025 (12)	0.015 (8)	-0.013 (9)
C10'	0.090 (12)	0.066 (10)	0.046 (7)	0.005 (9)	0.025 (8)	0.016 (7)
N3'	0.092 (3)	0.097 (3)	0.057 (2)	-0.012 (3)	0.020 (2)	-0.009 (2)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

Zn1—N5	1.974 (4)	C5—H5	0.9300
Zn1—N4	1.986 (4)	C6—C7	1.499 (6)
Zn1—N2	2.088 (3)	C7—H7A	0.9600
Zn1—N3	2.163 (4)	C7—H7B	0.9600
Zn1—N1	2.195 (3)	C7—H7C	0.9600
S1—C11	1.605 (5)	C8—C9	1.559 (7)
S2—C12	1.601 (5)	C8—H8A	0.9700
N1—C5	1.330 (5)	C8—H8B	0.9700
N1—C1	1.342 (5)	C9—N3	1.388 (6)
N2—C6	1.273 (5)	C9—H9A	0.9700
N2—C8	1.441 (5)	C9—H9B	0.9700
N4—C11	1.154 (5)	C10—N3	1.437 (7)
N5—C12	1.145 (5)	C10—H10A	0.9600
C1—C2	1.383 (5)	C10—H10B	0.9600
C1—C6	1.490 (6)	C10—H10C	0.9600
C2—C3	1.371 (6)	N3—H3A	0.9100
C2—H2	0.9300	C9'—H9'A	0.9700
C3—C4	1.354 (6)	C9'—H9'B	0.9700
C3—H3	0.9300	C10'—H10D	0.9600
C4—C5	1.385 (6)	C10'—H10E	0.9600
C4—H4	0.9300	C10'—H10F	0.9600
N5—Zn1—N4	110.56 (16)	C6—C7—H7A	109.5
N5—Zn1—N2	131.51 (14)	C6—C7—H7B	109.5
N4—Zn1—N2	117.70 (15)	H7A—C7—H7B	109.5
N5—Zn1—N3	97.92 (16)	C6—C7—H7C	109.5
N4—Zn1—N3	99.59 (16)	H7A—C7—H7C	109.5
N2—Zn1—N3	79.09 (15)	H7B—C7—H7C	109.5
N5—Zn1—N1	95.51 (13)	N4—C11—S1	179.4 (4)
N4—Zn1—N1	96.97 (14)	N5—C12—S2	178.5 (4)
N2—Zn1—N1	74.85 (12)	N2—C8—C9	107.8 (4)
N3—Zn1—N1	153.44 (15)	N2—C8—H8A	110.2
C5—N1—C1	118.3 (3)	C9—C8—H8A	110.2
C5—N1—Zn1	127.2 (3)	N2—C8—H8B	110.2
C1—N1—Zn1	114.3 (3)	C9—C8—H8B	110.2
C6—N2—C8	124.6 (4)	H8A—C8—H8B	108.5
C6—N2—Zn1	119.6 (3)	N3—C9—C8	111.9 (5)
C8—N2—Zn1	115.5 (3)	N3—C9—H9A	109.2
C11—N4—Zn1	159.8 (4)	C8—C9—H9A	109.2
C12—N5—Zn1	177.7 (4)	N3—C9—H9B	109.2

N1—C1—C2	121.3 (4)	C8—C9—H9B	109.2
N1—C1—C6	114.3 (3)	H9A—C9—H9B	107.9
C2—C1—C6	124.4 (4)	N3—C10—H10A	109.5
C3—C2—C1	119.2 (4)	N3—C10—H10B	109.5
C3—C2—H2	120.4	H10A—C10—H10B	109.5
C1—C2—H2	120.4	N3—C10—H10C	109.5
C4—C3—C2	120.2 (4)	H10A—C10—H10C	109.5
C4—C3—H3	119.9	H10B—C10—H10C	109.5
C2—C3—H3	119.9	C9—N3—C10	119.7 (7)
C3—C4—C5	117.8 (4)	C9—N3—Zn1	109.1 (3)
C3—C4—H4	121.1	C10—N3—Zn1	114.1 (5)
C5—C4—H4	121.1	C9—N3—H3A	104.0
N1—C5—C4	123.3 (4)	C10—N3—H3A	104.0
N1—C5—H5	118.4	Zn1—N3—H3A	104.0
C4—C5—H5	118.4	H9'A—C9'—H9'B	108.1
N2—C6—C1	116.2 (3)	H10D—C10'—H10E	109.5
N2—C6—C7	124.7 (4)	H10D—C10'—H10F	109.5
C1—C6—C7	119.1 (4)	H10E—C10'—H10F	109.5

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3A···S1 <sup>i</sup>	0.91	2.66	3.551 (5)	165

Symmetry code: (i)  $x-1, y, z$ .