

1-Chloro-1-[(4-nitrophenyl)hydrazinylidene]propan-2-one

Abdullah M. Asiri,^{a,b} Abdulrahman O. Al-Youbi,^a
Mohie E. M. Zayed^a and Seik Weng Ng^{c,a*}

^aChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia, ^bThe Center of Excellence for Advanced Materials Research, King Abdul Aziz University, PO Box 8020 Jeddah, Saudi Arabia, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 29 June 2011; accepted 3 July 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.054; wR factor = 0.152; data-to-parameter ratio = 14.0.

The non-H atoms of the title compound, $\text{C}_9\text{H}_8\text{ClN}_3\text{O}_3$, lie approximately on a plane (r.m.s. deviation = 0.111 \AA), and the $\text{C}=\text{N}$ double bond has a Z configuration. In the crystal, adjacent molecules are linked by an $\text{N}-\text{H}\cdots\text{O}_{\text{carbonyl}}$ hydrogen bond, forming a chain running along [101].

Related literature

For the synthesis, see: Benincori *et al.* (1990); Sayed *et al.* (2002). For background to the title compound, see: Asiri *et al.* (2010).



Experimental

Crystal data

$\text{C}_9\text{H}_8\text{ClN}_3\text{O}_3$
 $M_r = 241.63$

Monoclinic, $P2_1/n$
 $a = 7.0628(3)\text{ \AA}$

$b = 13.4182(5)\text{ \AA}$
 $c = 11.2884(5)\text{ \AA}$
 $\beta = 95.589(4)^\circ$
 $V = 1064.72(8)\text{ \AA}^3$
 $Z = 4$

$\text{Cu } K\alpha$ radiation
 $\mu = 3.19\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.20 \times 0.10 \times 0.05\text{ mm}$

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.568$, $T_{\max} = 0.857$

4113 measured reflections
2105 independent reflections
1839 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.152$
 $S = 1.09$
2105 reflections
150 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 1.21\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.48\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2 \cdots O1 ⁱ	0.85 (4)	2.26 (4)	3.000 (3)	145 (3)
Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$				

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdulaziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5261).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Asiri, A. M., Zayed, M. E. M. & Ng, S. W. (2010). *Acta Cryst. E* **66**, o2374.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Benincori, T., Fusco, R. & Sannicolo, F. (1990). *Gazz. Chim. Ital.* **120**, 635–659.
- Sayed, S. M., Khalil, M. A., Ahmed, M. A. & Raslan, M. A. (2002). *Synth. Commun.* **32**, 481–495.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, o1963 [doi:10.1107/S1600536811026407]

1-Chloro-1-[(4-nitrophenyl)hydrazinylidene]propan-2-one

Abdullah M. Asiri, Abdulrahman O. Al-Youbi, Mohie E. M. Zayed and Seik Weng Ng

S1. Comment

We have previously reported the synthesis of ethyl (*Z*)-2-chloro-2-(2-phenylhydrazin-1-ylidene) acetate by the reaction of benzenediazonium chloride with ethyl 2-chloro-3-oxobutanoate (Asiri *et al.*, 2010). The compound is an ester. In the present study, the use of a substituted benzenediazonium chloride and the methyl ester (instead of the ethyl ester) afforded a 1-chloro-1-(arylhydrazone)-2-propanone. Such ketones are intermediates in the synthesis of pyrazoles (Sayed *et al.*, 2002) and other heterocycles (Benincori *et al.*, 1990). In the 4-nitro substituted compound (Scheme I, Fig. 1), the non-hydrogen atoms lie on a plane [r.m.s. deviation 0.111 Å] (Scheme I, Fig. 1). The C_{aryl}—N(H)—N=C(S)=O portion adopts an extended zigzag conformation. Adjacent molecules are linked by an N—H···O_{carbonyl} hydrogen bond to form a chain running [1 0 1].

S2. Experimental


To a stirred solution of methyl 2-chloro-3-oxobutanoate (1.64 g, 10 mmol) in ethanol (100 ml) was added sodium acetate trihydrate (1.30 g, 10 mmol). The mixture was chilled to 273 K and then treated with a cold solution of *p*-nitrobenzene-diazonium chloride, prepared by diazotizing *p*-nitroaniline (1.38 g, 10 mmol) dissolved in 6*M* hydrochloric acid (6 ml) with a solution of sodium nitrite (0.70 g, 10 mmol) in water (10 ml). The addition of the diazonium salt solution was carried out with rapid stirring over a period of 20 min. The reaction mixture was stirred for further 15 min. and left for 3 h in refrigerator. The resulting solid was collected by filtration and washed thoroughly with water. The crude product was crystallized from ethanol to give the corresponding hydrazoneyl chloride.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 to 0.98 Å, U_{iso}(H) 1.2 to 1.5U_{eq}(C)] and were included in the refinement in the riding model approximation.

The amino H-atom was located in a difference Fourier map, and was freely refined.

The final difference Fourier map had a peak in the vicinity of H6.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of $C_9H_8ClN_3O_3$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

1-Chloro-1-[(4-nitrophenyl)hydrazinylidene]propan-2-one

Crystal data

$C_9H_8ClN_3O_3$
 $M_r = 241.63$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 7.0628 (3)$ Å
 $b = 13.4182 (5)$ Å
 $c = 11.2884 (5)$ Å
 $\beta = 95.589 (4)$ °
 $V = 1064.72 (8)$ Å³
 $Z = 4$

$F(000) = 496$
 $D_x = 1.507 \text{ Mg m}^{-3}$
Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å
Cell parameters from 1647 reflections
 $\theta = 3.3\text{--}74.3$ °
 $\mu = 3.19 \text{ mm}^{-1}$
 $T = 100$ K
Prism, yellow
 $0.20 \times 0.10 \times 0.05$ mm

Data collection

Agilent SuperNova Dual
diffractometer with an Atlas detector
Radiation source: SuperNova (Cu) X-ray
Source
Mirror monochromator
Detector resolution: 10.4041 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(CrysAlis PRO; Agilent, 2010)

$T_{\min} = 0.568, T_{\max} = 0.857$
4113 measured reflections
2105 independent reflections
1839 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 $\theta_{\max} = 74.4$ °, $\theta_{\min} = 5.1$ °
 $h = -8 \rightarrow 5$
 $k = -16 \rightarrow 16$
 $l = -14 \rightarrow 14$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.054$$

$$wR(F^2) = 0.152$$

$$S = 1.09$$

2105 reflections

150 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0849P)^2 + 0.8946P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 1.21 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.63780 (10)	0.29860 (4)	0.39614 (6)	0.0299 (2)
O1	0.4446 (3)	0.25504 (13)	0.15943 (16)	0.0275 (4)
O2	0.8827 (3)	-0.34157 (14)	0.6658 (2)	0.0369 (5)
O3	1.0132 (3)	-0.25969 (16)	0.81885 (18)	0.0383 (5)
N1	0.6623 (3)	0.10048 (16)	0.37953 (18)	0.0213 (5)
N2	0.7385 (3)	0.09908 (17)	0.49160 (19)	0.0226 (5)
H2	0.753 (5)	0.154 (3)	0.528 (3)	0.042 (10)*
N3	0.9267 (3)	-0.26356 (18)	0.7182 (2)	0.0300 (5)
C1	0.5294 (4)	0.0846 (2)	0.1360 (2)	0.0304 (6)
H1A	0.4572	0.0912	0.0578	0.046*
H1B	0.4743	0.0311	0.1808	0.046*
H1C	0.6622	0.0686	0.1258	0.046*
C2	0.5211 (4)	0.1804 (2)	0.2026 (2)	0.0251 (6)
C3	0.6129 (4)	0.18280 (19)	0.3275 (2)	0.0224 (5)
C4	0.7818 (3)	0.00828 (19)	0.5472 (2)	0.0209 (5)
C5	0.8644 (4)	0.0084 (2)	0.6644 (2)	0.0246 (5)
H5	0.8885	0.0696	0.7054	0.030*
C6	0.9115 (4)	-0.0816 (2)	0.7210 (2)	0.0250 (5)
H6	0.9677	-0.0825	0.8009	0.030*
C7	0.8754 (4)	-0.16926 (19)	0.6594 (2)	0.0237 (5)
C8	0.7916 (4)	-0.17086 (19)	0.5427 (2)	0.0241 (5)
H8	0.7668	-0.2324	0.5026	0.029*
C9	0.7449 (4)	-0.08185 (19)	0.4860 (2)	0.0229 (5)
H9	0.6882	-0.0815	0.4061	0.027*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0424 (4)	0.0166 (4)	0.0299 (4)	0.0005 (2)	0.0002 (3)	0.0006 (2)
O1	0.0353 (10)	0.0225 (9)	0.0246 (9)	0.0050 (8)	0.0026 (8)	0.0069 (7)
O2	0.0469 (12)	0.0195 (10)	0.0448 (12)	0.0028 (9)	0.0070 (10)	0.0060 (9)
O3	0.0513 (13)	0.0354 (11)	0.0272 (10)	0.0140 (10)	-0.0015 (9)	0.0142 (9)
N1	0.0244 (10)	0.0193 (10)	0.0202 (10)	0.0008 (8)	0.0019 (8)	0.0023 (8)

N2	0.0300 (11)	0.0183 (10)	0.0195 (10)	0.0001 (9)	0.0014 (8)	-0.0007 (9)
N3	0.0329 (12)	0.0262 (12)	0.0322 (12)	0.0067 (10)	0.0099 (10)	0.0094 (10)
C1	0.0410 (15)	0.0266 (14)	0.0233 (12)	0.0013 (12)	0.0021 (11)	-0.0004 (11)
C2	0.0279 (12)	0.0229 (12)	0.0253 (13)	0.0009 (10)	0.0066 (10)	0.0058 (10)
C3	0.0265 (12)	0.0169 (11)	0.0244 (12)	0.0022 (10)	0.0062 (10)	0.0026 (10)
C4	0.0214 (11)	0.0179 (12)	0.0242 (12)	0.0006 (9)	0.0060 (9)	0.0034 (10)
C5	0.0291 (12)	0.0206 (13)	0.0245 (12)	-0.0018 (10)	0.0046 (10)	-0.0028 (10)
C6	0.0259 (12)	0.0297 (14)	0.0193 (11)	0.0032 (10)	0.0016 (9)	0.0044 (10)
C7	0.0266 (12)	0.0197 (13)	0.0254 (13)	0.0052 (10)	0.0059 (10)	0.0097 (10)
C8	0.0287 (12)	0.0186 (12)	0.0259 (13)	-0.0007 (10)	0.0071 (10)	-0.0007 (10)
C9	0.0275 (12)	0.0198 (12)	0.0214 (12)	-0.0009 (10)	0.0027 (9)	0.0002 (10)

Geometric parameters (Å, °)

C1—C3	1.737 (3)	C1—H1C	0.9800
O1—C2	1.217 (3)	C2—C3	1.494 (4)
O2—N3	1.227 (3)	C4—C5	1.393 (4)
O3—N3	1.238 (3)	C4—C9	1.405 (4)
N1—C3	1.283 (3)	C5—C6	1.391 (4)
N1—N2	1.326 (3)	C5—H5	0.9500
N2—C4	1.391 (3)	C6—C7	1.377 (4)
N2—H2	0.85 (4)	C6—H6	0.9500
N3—C7	1.458 (3)	C7—C8	1.391 (4)
C1—C2	1.493 (4)	C8—C9	1.380 (4)
C1—H1A	0.9800	C8—H8	0.9500
C1—H1B	0.9800	C9—H9	0.9500
C3—N1—N2	121.0 (2)	N2—C4—C5	118.7 (2)
N1—N2—C4	119.6 (2)	N2—C4—C9	120.7 (2)
N1—N2—H2	118 (3)	C5—C4—C9	120.6 (2)
C4—N2—H2	122 (3)	C6—C5—C4	119.6 (2)
O2—N3—O3	123.9 (2)	C6—C5—H5	120.2
O2—N3—C7	118.7 (2)	C4—C5—H5	120.2
O3—N3—C7	117.4 (2)	C7—C6—C5	119.1 (2)
C2—C1—H1A	109.5	C7—C6—H6	120.5
C2—C1—H1B	109.5	C5—C6—H6	120.5
H1A—C1—H1B	109.5	C6—C7—C8	122.1 (2)
C2—C1—H1C	109.5	C6—C7—N3	119.1 (2)
H1A—C1—H1C	109.5	C8—C7—N3	118.8 (2)
H1B—C1—H1C	109.5	C9—C8—C7	119.1 (2)
O1—C2—C1	123.0 (2)	C9—C8—H8	120.4
O1—C2—C3	119.7 (2)	C7—C8—H8	120.4
C1—C2—C3	117.3 (2)	C8—C9—C4	119.5 (2)
N1—C3—C2	119.1 (2)	C8—C9—H9	120.2
N1—C3—C11	123.7 (2)	C4—C9—H9	120.2
C2—C3—C11	117.12 (18)		
C3—N1—N2—C4	176.8 (2)	C5—C6—C7—C8	-0.6 (4)

N2—N1—C3—C2	−177.8 (2)	C5—C6—C7—N3	179.3 (2)
N2—N1—C3—Cl1	−0.4 (3)	O2—N3—C7—C6	175.2 (2)
O1—C2—C3—N1	167.5 (2)	O3—N3—C7—C6	−4.9 (4)
C1—C2—C3—N1	−12.9 (4)	O2—N3—C7—C8	−4.9 (4)
O1—C2—C3—Cl1	−10.0 (3)	O3—N3—C7—C8	175.0 (2)
C1—C2—C3—Cl1	169.63 (19)	C6—C7—C8—C9	0.8 (4)
N1—N2—C4—C5	179.0 (2)	N3—C7—C8—C9	−179.1 (2)
N1—N2—C4—C9	−0.3 (3)	C7—C8—C9—C4	−0.3 (4)
N2—C4—C5—C6	−179.0 (2)	N2—C4—C9—C8	179.2 (2)
C9—C4—C5—C6	0.3 (4)	C5—C4—C9—C8	−0.2 (4)
C4—C5—C6—C7	0.1 (4)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N2—H2···O1 ⁱ	0.85 (4)	2.26 (4)	3.000 (3)	145 (3)

Symmetry code: (i) $x+1/2, -y+1/2, z+1/2$.