

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(4-Bromo-2-[[2-(morpholin-4-yl)ethyl-imino]methyl]phenolato)dioxido-vanadium(V)

Chen-Yi Wang,* Xiang Wu, Feng Cao and Cai-Jun Yuan

Department of Chemistry, Huzhou University, Huzhou 313000, People's Republic of China

Correspondence e-mail: chen_yi_wang@163.com


Received 25 August 2011; accepted 29 August 2011

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.044; wR factor = 0.096; data-to-parameter ratio = 16.9.

In the title mononuclear dioxido vanadium(V) complex, $[\text{V}(\text{C}_{13}\text{H}_{16}\text{BrN}_2\text{O}_2)\text{O}_2]$, the V^{V} atom is five-coordinated by one phenolate O, one imine N and one morpholine N atom of the Schiff base ligand, and by two oxide O atoms, forming a distorted square-pyramidal geometry. In the crystal, weak $\text{C}-\text{H}\cdots\text{O}$ interactions and a short $\text{Br}\cdots\text{Br}$ contact [3.4597 (12) Å] are observed.

Related literature

For related Schiff base complexes that we have reported recently, see: Wang (2009, 2011); Wang & Ye (2011). For similar oxidovanadium(V) complexes, see: Xie *et al.* (2004); Gao *et al.* (2005); Hartung *et al.* (2007); Romanowski *et al.* (2009).



Experimental

Crystal data

$[\text{V}(\text{C}_{13}\text{H}_{16}\text{BrN}_2\text{O}_2)\text{O}_2]$	$V = 1468.0$ (5) Å ³
$M_r = 395.13$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 21.372$ (3) Å	$\mu = 3.41$ mm ⁻¹
$b = 6.0892$ (15) Å	$T = 298$ K
$c = 11.372$ (3) Å	$0.17 \times 0.13 \times 0.13$ mm
$\beta = 97.248$ (2)°	

Data collection

Bruker SMART CCD area-detector diffractometer	11267 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3204 independent reflections
$T_{\min} = 0.595$, $T_{\max} = 0.665$	2058 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	190 parameters
$wR(F^2) = 0.096$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.61$ e Å ⁻³
3204 reflections	$\Delta\rho_{\text{min}} = -0.62$ e Å ⁻³

Table 1

Selected bond lengths (Å).

V1—O4	1.611 (2)	V1—N1	2.142 (3)
V1—O3	1.622 (3)	V1—N2	2.159 (3)
V1—O1	1.907 (3)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C7}-\text{H7}\cdots\text{O1}^{\text{i}}$	0.93	2.53	3.241 (4)	133
$\text{C11}-\text{H11A}\cdots\text{O2}^{\text{ii}}$	0.97	2.57	3.479 (5)	156

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported financially by the Natural Science Foundation of China (No. 31071856), the Applied Research Project on Nonprofit Technology of Zhejiang Province (No. 2010C32060), the Natural Science Foundation of Zhejiang Province (No. Y407318), and the Technological Innovation Project (Sinfonietta Talent Plan) of college students in Zhejiang Province (Nos. 2010R42525 and 2011R425027).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2767).

References

- Bruker (1998). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gao, S., Huo, L.-H., Deng, Z.-P. & Zhao, H. (2005). *Acta Cryst.* **E61**, m978–m980.
- Hartung, J., Ludwig, A., Svoboda, I. & Fuess, H. (2007). *Acta Cryst.* **E63**, m1422–m1423.
- Romanowski, G., Wera, M. & Sikorski, A. (2009). *Acta Cryst.* **E65**, m190.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, C.-Y. (2009). *J. Coord. Chem.* **62**, 2860–2868.
- Wang, C.-Y. (2011). *Acta Cryst.* **E67**, m1085–m1086.
- Wang, C.-Y. & Ye, J.-Y. (2011). *Russ. J. Coord. Chem.* **37**, 235–241.
- Xie, M.-J., Ping, Y.-S., Zheng, L.-D., Hui, J.-Z. & Peng, C. (2004). *Acta Cryst.* **E60**, m1382–m1383.

supporting information

Acta Cryst. (2011). E67, m1330 [https://doi.org/10.1107/S1600536811035240]

(4-Bromo-2-[[2-(morpholin-4-yl)ethylimino]methyl]-phenolato)dioxidovanadium(V)

Chen-Yi Wang, Xiang Wu, Feng Cao and Cai-Jun Yuan

S1. Comment

As part of our investigations into new Schiff base complexes (Wang & Ye, 2011; Wang, 2009, 2011), we have synthesized the title compound, a new mononuclear dioxovanadium(V) complex (Fig. 1). The V^V atom in the complex is five-coordinated by one phenolate O, one imine N and one morpholine N atom of the Schiff base ligand, and by two oxo O atoms, forming a distorted square pyramidal coordination. The V–O and V–N bond lengths (Table 1) are typical and are comparable with those observed in other similar oxovanadium(V) complexes with Schiff bases (Xie *et al.*, 2004; Gao *et al.*, 2005; Hartung *et al.*, 2007; Romanowski *et al.*, 2009).

S2. Experimental

5-Bromosalicylaldehyde (1.0 mmol, 0.201 g), 2-morpholin-4-ylethylamine (1.0 mmol, 0.130 g) and VO(acac)₂ (1.0 mmol, 0.265 g) were dissolved in MeOH (30 ml). The mixture was stirred at room temperature for 10 min to give a yellow solution. After keeping the solution in air for a week, yellow block-shaped crystals were formed at the bottom of the vessel.

S3. Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H})$ set at $1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

(4-Bromo-2-[[2-(morpholin-4-yl)ethylimino]methyl]phenolato)dioxidovanadium(V)

Crystal data

[V(C₁₃H₁₆BrN₂O₂)O₂] $M_r = 395.13$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 21.372$ (3) Å $b = 6.0892$ (15) Å $c = 11.372$ (3) Å $\beta = 97.248$ (2)° $V = 1468.0$ (5) Å³ $Z = 4$ $F(000) = 792$ $D_x = 1.788$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2133 reflections

 $\theta = 2.7$ – 25.1 ° $\mu = 3.41$ mm⁻¹ $T = 298$ K

Block, yellow

 $0.17 \times 0.13 \times 0.13$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scansAbsorption correction: multi-scan
(SADABS; Sheldrick, 1996) $T_{\min} = 0.595$, $T_{\max} = 0.665$

11267 measured reflections

3204 independent reflections

2058 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.051$ $\theta_{\text{max}} = 27.0$ °, $\theta_{\text{min}} = 2.9$ ° $h = -27 \rightarrow 27$ $k = -7 \rightarrow 7$ $l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.044$ $wR(F^2) = 0.096$ $S = 1.02$

3204 reflections

190 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0418P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.61$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.62$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
V1	0.28158 (3)	0.31216 (10)	0.45502 (5)	0.02930 (17)
Br1	0.01304 (2)	-0.30291 (9)	0.10991 (4)	0.06124 (19)
N1	0.23304 (13)	0.4015 (4)	0.2854 (2)	0.0255 (6)
N2	0.34664 (13)	0.5238 (4)	0.3783 (2)	0.0253 (7)

O1	0.21270 (13)	0.1134 (5)	0.4562 (2)	0.0482 (7)
O2	0.47871 (12)	0.5987 (5)	0.3519 (3)	0.0527 (8)
O3	0.33864 (12)	0.1419 (4)	0.4953 (2)	0.0442 (7)
O4	0.27296 (13)	0.4752 (4)	0.5634 (2)	0.0473 (7)
C1	0.15474 (16)	0.1138 (6)	0.2635 (3)	0.0290 (8)
C2	0.17008 (17)	0.0235 (6)	0.3773 (3)	0.0329 (9)
C3	0.13930 (18)	-0.1665 (6)	0.4067 (3)	0.0407 (10)
H3	0.1501	-0.2298	0.4809	0.049*
C4	0.09347 (19)	-0.2617 (6)	0.3282 (4)	0.0431 (10)
H4	0.0730	-0.3876	0.3496	0.052*
C5	0.07759 (18)	-0.1697 (7)	0.2164 (3)	0.0401 (10)
C6	0.10726 (17)	0.0147 (6)	0.1835 (3)	0.0372 (9)
H6	0.0962	0.0748	0.1085	0.045*
C7	0.18586 (17)	0.3060 (6)	0.2271 (3)	0.0307 (8)
H7	0.1702	0.3667	0.1542	0.037*
C8	0.25689 (16)	0.5992 (6)	0.2292 (3)	0.0301 (8)
H8A	0.2228	0.7031	0.2088	0.036*
H8B	0.2741	0.5587	0.1573	0.036*
C9	0.30743 (16)	0.7010 (5)	0.3164 (3)	0.0304 (8)
H9A	0.3337	0.7968	0.2751	0.036*
H9B	0.2883	0.7882	0.3737	0.036*
C10	0.38229 (17)	0.3950 (6)	0.2964 (3)	0.0351 (9)
H10A	0.3534	0.3487	0.2283	0.042*
H10B	0.3994	0.2640	0.3371	0.042*
C11	0.43528 (19)	0.5240 (7)	0.2543 (3)	0.0494 (12)
H11A	0.4573	0.4323	0.2031	0.059*
H11B	0.4181	0.6492	0.2084	0.059*
C12	0.44707 (19)	0.7346 (7)	0.4274 (4)	0.0459 (11)
H12A	0.4307	0.8634	0.3837	0.055*
H12B	0.4770	0.7837	0.4934	0.055*
C13	0.39376 (17)	0.6159 (6)	0.4741 (3)	0.0362 (9)
H13A	0.4109	0.4971	0.5252	0.043*
H13B	0.3725	0.7166	0.5219	0.043*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
V1	0.0394 (4)	0.0279 (4)	0.0198 (3)	-0.0050 (3)	0.0006 (3)	0.0003 (3)
Br1	0.0583 (3)	0.0709 (4)	0.0544 (3)	-0.0337 (3)	0.0067 (2)	-0.0207 (2)
N1	0.0354 (17)	0.0205 (16)	0.0202 (15)	-0.0021 (13)	0.0015 (13)	0.0013 (12)
N2	0.0338 (17)	0.0198 (16)	0.0215 (14)	-0.0002 (13)	0.0006 (12)	-0.0009 (12)
O1	0.0541 (18)	0.0570 (19)	0.0305 (15)	-0.0253 (15)	-0.0059 (13)	0.0126 (14)
O2	0.0331 (16)	0.068 (2)	0.0572 (18)	-0.0001 (15)	0.0048 (14)	0.0167 (17)
O3	0.0503 (17)	0.0344 (16)	0.0459 (16)	0.0026 (13)	-0.0017 (13)	0.0139 (13)
O4	0.0660 (19)	0.0461 (18)	0.0318 (14)	-0.0093 (15)	0.0135 (13)	-0.0113 (13)
C1	0.034 (2)	0.025 (2)	0.029 (2)	-0.0045 (16)	0.0059 (16)	-0.0027 (16)
C2	0.033 (2)	0.034 (2)	0.033 (2)	-0.0049 (18)	0.0085 (17)	-0.0014 (17)
C3	0.045 (2)	0.045 (3)	0.033 (2)	-0.011 (2)	0.0064 (18)	0.0057 (19)

C4	0.048 (3)	0.033 (2)	0.051 (3)	-0.0123 (19)	0.018 (2)	-0.003 (2)
C5	0.037 (2)	0.045 (3)	0.039 (2)	-0.0175 (19)	0.0094 (18)	-0.019 (2)
C6	0.039 (2)	0.041 (3)	0.031 (2)	-0.0065 (19)	0.0039 (17)	-0.0050 (18)
C7	0.037 (2)	0.037 (2)	0.0181 (17)	0.0017 (18)	0.0026 (15)	-0.0014 (17)
C8	0.037 (2)	0.024 (2)	0.0287 (19)	0.0028 (16)	0.0010 (16)	0.0062 (16)
C9	0.033 (2)	0.022 (2)	0.035 (2)	0.0019 (16)	0.0016 (16)	-0.0003 (17)
C10	0.045 (2)	0.030 (2)	0.030 (2)	0.0120 (18)	0.0030 (17)	0.0002 (17)
C11	0.047 (3)	0.065 (3)	0.039 (2)	0.011 (2)	0.017 (2)	0.010 (2)
C12	0.041 (2)	0.043 (3)	0.051 (3)	-0.008 (2)	-0.008 (2)	0.007 (2)
C13	0.041 (2)	0.038 (2)	0.027 (2)	-0.0040 (19)	-0.0072 (17)	-0.0010 (17)

Geometric parameters (Å, °)

V1—O4	1.611 (2)	C4—C5	1.392 (5)
V1—O3	1.622 (3)	C4—H4	0.9300
V1—O1	1.907 (3)	C5—C6	1.365 (5)
V1—N1	2.142 (3)	C6—H6	0.9300
V1—N2	2.159 (3)	C7—H7	0.9300
Br1—C5	1.899 (4)	C8—C9	1.504 (5)
N1—C7	1.275 (4)	C8—H8A	0.9700
N1—C8	1.483 (4)	C8—H8B	0.9700
N2—C9	1.487 (4)	C9—H9A	0.9700
N2—C13	1.497 (4)	C9—H9B	0.9700
N2—C10	1.497 (4)	C10—C11	1.505 (5)
O1—C2	1.314 (4)	C10—H10A	0.9700
O2—C12	1.423 (5)	C10—H10B	0.9700
O2—C11	1.428 (4)	C11—H11A	0.9700
C1—C2	1.406 (5)	C11—H11B	0.9700
C1—C6	1.410 (5)	C12—C13	1.502 (5)
C1—C7	1.433 (5)	C12—H12A	0.9700
C2—C3	1.392 (5)	C12—H12B	0.9700
C3—C4	1.368 (5)	C13—H13A	0.9700
C3—H3	0.9300	C13—H13B	0.9700
O4—V1—O3	109.41 (14)	N1—C7—C1	126.0 (3)
O4—V1—O1	102.89 (13)	N1—C7—H7	117.0
O3—V1—O1	98.38 (13)	C1—C7—H7	117.0
O4—V1—N1	116.30 (13)	N1—C8—C9	107.9 (3)
O3—V1—N1	132.70 (12)	N1—C8—H8A	110.1
O1—V1—N1	83.17 (11)	C9—C8—H8A	110.1
O4—V1—N2	94.75 (12)	N1—C8—H8B	110.1
O3—V1—N2	89.77 (12)	C9—C8—H8B	110.1
O1—V1—N2	156.74 (10)	H8A—C8—H8B	108.4
N1—V1—N2	75.39 (10)	N2—C9—C8	109.1 (3)
C7—N1—C8	116.0 (3)	N2—C9—H9A	109.9
C7—N1—V1	127.9 (2)	C8—C9—H9A	109.9
C8—N1—V1	116.1 (2)	N2—C9—H9B	109.9
C9—N2—C13	111.1 (3)	C8—C9—H9B	109.9

C9—N2—C10	112.7 (3)	H9A—C9—H9B	108.3
C13—N2—C10	107.5 (3)	N2—C10—C11	112.7 (3)
C9—N2—V1	105.7 (2)	N2—C10—H10A	109.0
C13—N2—V1	109.74 (19)	C11—C10—H10A	109.0
C10—N2—V1	110.0 (2)	N2—C10—H10B	109.0
C2—O1—V1	136.7 (2)	C11—C10—H10B	109.0
C12—O2—C11	110.0 (3)	H10A—C10—H10B	107.8
C2—C1—C6	119.5 (3)	O2—C11—C10	111.1 (3)
C2—C1—C7	121.5 (3)	O2—C11—H11A	109.4
C6—C1—C7	119.0 (3)	C10—C11—H11A	109.4
O1—C2—C3	119.3 (3)	O2—C11—H11B	109.4
O1—C2—C1	121.8 (3)	C10—C11—H11B	109.4
C3—C2—C1	118.9 (3)	H11A—C11—H11B	108.0
C4—C3—C2	121.1 (4)	O2—C12—C13	111.9 (3)
C4—C3—H3	119.4	O2—C12—H12A	109.2
C2—C3—H3	119.4	C13—C12—H12A	109.2
C3—C4—C5	119.8 (4)	O2—C12—H12B	109.2
C3—C4—H4	120.1	C13—C12—H12B	109.2
C5—C4—H4	120.1	H12A—C12—H12B	107.9
C6—C5—C4	120.9 (4)	N2—C13—C12	113.1 (3)
C6—C5—Br1	120.1 (3)	N2—C13—H13A	109.0
C4—C5—Br1	119.0 (3)	C12—C13—H13A	109.0
C5—C6—C1	119.7 (4)	N2—C13—H13B	109.0
C5—C6—H6	120.1	C12—C13—H13B	109.0
C1—C6—H6	120.1	H13A—C13—H13B	107.8

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C7—H7 \cdots O1 ⁱ	0.93	2.53	3.241 (4)	133
C11—H11A \cdots O2 ⁱⁱ	0.97	2.57	3.479 (5)	156

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+1, y-1/2, -z+1/2$.