

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dicaesium diaquabis(methylene-diphosphonato- κ^2O,O')cobaltate(II)

Kina van der Merwe,* Hendrik G. Visser and Johan A. Venter

Department of Chemistry, University of the Free State, PO Box 339, Bloemfontein 9330, South Africa

Correspondence e-mail: vandermerwe@gmail.com


Received 24 August 2011; accepted 30 August 2011

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{Cs}-\text{O}) = 0.003$ Å; R factor = 0.015; wR factor = 0.044; data-to-parameter ratio = 14.4.

The asymmetric unit of the title compound, $\text{Cs}_2[\text{Co}(\text{CH}_4\text{O}_6\text{P}_2)_2(\text{H}_2\text{O})_2]$, is comprised of one bidentate methylenediphosphonate ligand and one water molecule which are coordinated to the Co^{II} atom, as well as a caesium counter-cation. The Co atom occupies a special position on a crystallographic inversion center. The caesium ion is octahedrally coordinated by six O atoms with Cs—O distances ranging from 3.119 (2) to 3.296 (2) Å. A three-dimensional network is formed through $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For related structures, see: Fleisch (1991); Neville-Webbe *et al.* (2002); Van der Merwe *et al.* (2010). For bond lengths and bond angles in related structures, see: Bao *et al.* (2003); Cao *et al.* (2007); Gong *et al.* (2006); Van der Merwe *et al.* (2009); Visser *et al.* (2010); Yin *et al.* (2003).



Experimental

Crystal data

$\text{Cs}_2[\text{Co}(\text{CH}_4\text{O}_6\text{P}_2)_2(\text{H}_2\text{O})_2]$
 $M_r = 708.75$
 Triclinic, $P\bar{1}$
 $a = 7.333$ (5) Å
 $b = 7.412$ (5) Å
 $c = 7.666$ (5) Å

$\alpha = 74.621$ (5)°
 $\beta = 83.064$ (5)°
 $\gamma = 86.496$ (5)°
 $V = 398.6$ (5) Å³
 $Z = 1$
 Mo $K\alpha$ radiation

$\mu = 5.96$ mm⁻¹
 $T = 293$ K

0.38 × 0.07 × 0.05 mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.210$, $T_{\text{max}} = 0.755$

4203 measured reflections
 1904 independent reflections
 1827 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$
 $wR(F^2) = 0.044$
 $S = 0.75$
 1904 reflections
 132 parameters
 11 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.47$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.48$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1A}\cdots\text{O4}^{\text{ii}}$	0.92 (2)	1.94 (2)	2.860 (3)	173 (3)
$\text{O5}-\text{H5B}\cdots\text{O4}^{\text{ii}}$	0.85 (2)	1.69 (3)	2.518 (3)	164 (7)

Symmetry codes: (i) $x, y, z + 1$; (ii) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 2005); software used to prepare material for publication: WinGX (Farrugia, 1999).

The University of the Free State and Professor A. Roodt are gratefully acknowledged for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2322).

References

- Bao, S., Zheng, L., Liu, Y., Xu, W. & Feng, S. (2003). *Inorg. Chem.* **42**, 5037–5039.
- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *SAINTE-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cao, D., Li, Y. & Zheng, L. (2007). *Inorg. Chem.* **46**, 7571–7578.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Fleisch, H. (1991). *Drugs*, **42**, 919–944.
- Gong, Y., Tang, W., Hou, W., Zha, Z. & Hu, C. (2006). *Inorg. Chem.* **45**, 4987–4995.
- Neville-Webbe, H. L., Holen, I. & Coleman, R. E. (2002). *Cancer Treat. Rev.* **28**, 305–319.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Van der Merwe, K. A., Visser, H. G. & Venter, J. A. (2009). *Acta Cryst.* **E65**, m1394.
- Van der Merwe, K., Visser, H. G. & Venter, J. A. (2010). *Acta Cryst.* **E66**, m1011–m1012.
- Visser, H. G., Venter, J. A. & Van der Merwe, K. A. (2010). *Acta Cryst.* **E66**, m159.
- Yin, P., Gao, S., Zheng, L. & Xin, X. (2003). *Chem. Mater.* **15**, 3233–3236.

supporting information

Acta Cryst. (2011). E67, m1354 [https://doi.org/10.1107/S1600536811035355]

Dicaesium diaquabis(methylenediphosphonato- κ^2O,O')cobaltate(II)**Kina van der Merwe, Hendrik G. Visser and Johan A. Venter****S1. Comment**

This work is part of an ongoing investigation aimed at synthesizing and characterizing new methylene diphosphonate complexes and expanding on our knowledge of the interactions of the methylene diphosphonate ligand with various metal centers. (Van der Merwe *et al.* (2009) & Van der Merwe *et al.* (2010)).

Methylene diphosphonates ($O_3PCH_2PO_3$) has a diversified coordination capability with metal ions, due to the single methyl group which divides the two phosphonate groups. The formation of a stable six-membered ring comprised of $M-O-P-C-P-O$ is favoured (Bao *et al.* (2003)). Bisphosphonates adhere strongly to hydroxyapatite crystals and constrain their formation and dissolution (Fleisch (1991)). This physicochemical *in vivo* effect may result in the prevention of soft tissue calcification or even prevent normal calcification. The bis(phosphonic acid) has a high affinity for bone surfaces and it is also non-hydrolyzable (Neville-Webbe *et al.* (2002)).

The asymmetric unit of the title compound, $Cs_2[Co(CH_4O_6P_2)_2(H_2O)_2]$, is comprised of one bidentate methylene diphosphonate ligand and one water molecule which are coordinated to the Co^{II} atom, as well as a non-coordinated caesium cation. The Co atom occupies a special position on a crystallographic inversion center. The caesium ion is octahedrally coordinated to six oxygen atoms with $Cs-O$ distances ranging from 3.119 (2) to 3.296 (2) Å. The two methylene diphosphonate ligands chelate to the central cobalt metal *via* four oxygen atoms ($O2/O2'$ and $O7/O7'$) from the phosphonate groups. This leads to the formation of two six-membered rings.

The Co^{II} metal center has a slightly distorted octahedral geometry with $O-Co-O$ angles ranging between 85.35 (8)° and 94.65 (8)°. The $Co-O$ bond lengths vary between 2.0761 (18) and 2.1272 (19) Å. These distances correspond to literature values (Bao *et al.* (2003); Cao *et al.* (2007); Gong *et al.* (2006); Van der Merwe *et al.* (2009); Visser *et al.* (2010); Yin *et al.* (2003)).

A three-dimensional network is provided by $O-H-O$ hydrogen bonds (Table 2).

S2. Experimental

$[Co(NH_3)_6]Cl_3$ (0,1700 g, 0,714 mmol) was dissolved in distilled water (10 cm^3) and the pH of the solution was lowered to 1.87 using hydrochloric acid. The solution was heated for 30 minutes at 313.15 K. Methylene diphosphonate (0.251 g, 1.43 mmol) was dissolved in distilled water (7 cm^3) and the pH of the solution was elevated to 1.93 using caesium chloride. Both solutions were combined and the pH was adjusted to 2.04, the pink solution was heated for 3 h at 353.15 K. Pink crystals, suitable for X-ray diffraction, was obtained. (Yield: 7.2%)

S3. Refinement

All H atoms were located from difference Fourier maps and were refined isotropically without further restraints. The highest residual electron density was located 0.88 Å from P1.




Figure 1

Representation of the title compound, showing the numbering scheme and displacement ellipsoids drawn at the 50% probability level. [Symmetry code: (i) $1 - x, -y, 1 - z$].

Dicaesium diaquabis(methylenediphosphonato- κ^2O,O')cobaltate(II)

Crystal data

$\text{Cs}_2[\text{Co}(\text{CH}_4\text{O}_6\text{P}_2)_2(\text{H}_2\text{O})_2]$

$M_r = 708.75$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.333\ (5)\ \text{\AA}$

$b = 7.412\ (5)\ \text{\AA}$

$c = 7.666\ (5)\ \text{\AA}$

$\alpha = 74.621\ (5)^\circ$

$\beta = 83.064\ (5)^\circ$

$\gamma = 86.496\ (5)^\circ$

$V = 398.6\ (5)\ \text{\AA}^3$

$Z = 1$

$F(000) = 253$

$D_x = 2.953\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3160 reflections

$\theta = 2.8\text{--}28.4^\circ$

$\mu = 5.96\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Needle, pink

$0.38 \times 0.07 \times 0.05\ \text{mm}$

Data collection

Bruker APEXII CCD

diffractometer

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.210$, $T_{\max} = 0.755$

4203 measured reflections

1904 independent reflections

1827 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 28^\circ$, $\theta_{\min} = 3.7^\circ$

$h = -9 \rightarrow 9$

$k = -9 \rightarrow 9$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.015$
 $wR(F^2) = 0.044$
 $S = 0.75$
 1904 reflections
 132 parameters
 11 restraints

H atoms treated by a mixture of independent
 and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0412P)^2 + 0.4899P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cs1	0.284577 (17)	0.954858 (18)	0.692901 (17)	0.00965 (6)	
Co1	0.5	0.5	0.5	0.00595 (9)	
P1	0.26392 (7)	0.34252 (8)	0.22866 (8)	0.00590 (11)	
P2	0.19616 (7)	0.74778 (8)	0.25147 (8)	0.00589 (11)	
O1	0.2886 (2)	0.4437 (2)	0.7201 (2)	0.0112 (3)	
O2	0.3857 (2)	0.3210 (2)	0.3786 (2)	0.0088 (3)	
O3	0.0667 (2)	0.2706 (2)	0.3122 (2)	0.0086 (3)	
O4	0.3336 (2)	0.2396 (2)	0.0861 (2)	0.0086 (3)	
O5	0.1822 (2)	0.9384 (2)	0.1036 (2)	0.0109 (3)	
O6	0.0107 (2)	0.7058 (2)	0.3621 (2)	0.0102 (3)	
O7	0.3541 (2)	0.7427 (2)	0.3623 (2)	0.0086 (3)	
C1	0.2397 (3)	0.5858 (3)	0.1121 (3)	0.0071 (4)	
H1A	0.313 (4)	0.383 (4)	0.837 (3)	0.023 (9)*	
H1B	0.185 (4)	0.410 (6)	0.704 (6)	0.063 (15)*	
H2	0.048 (5)	0.287 (5)	0.417 (3)	0.035 (10)*	
H3	0.145 (3)	0.599 (4)	0.033 (4)	0.018 (8)*	
H4	0.357 (3)	0.616 (4)	0.046 (4)	0.015 (8)*	
H5A	0.156 (9)	0.935 (9)	0.002 (5)	0.022*	0.5
H5B	0.239 (8)	1.030 (7)	0.116 (9)	0.022*	0.5

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cs1	0.01010 (9)	0.00910 (9)	0.00854 (9)	0.00056 (5)	-0.00040 (5)	-0.00068 (6)
Co1	0.00540 (19)	0.00593 (19)	0.0067 (2)	-0.00035 (15)	-0.00109 (15)	-0.00174 (16)
P1	0.0061 (2)	0.0053 (2)	0.0063 (3)	-0.00119 (19)	-0.0007 (2)	-0.0015 (2)
P2	0.0060 (2)	0.0060 (3)	0.0057 (3)	-0.00059 (19)	-0.0006 (2)	-0.0014 (2)
O1	0.0084 (8)	0.0160 (9)	0.0088 (8)	-0.0027 (7)	-0.0002 (6)	-0.0019 (7)
O2	0.0097 (7)	0.0077 (7)	0.0092 (8)	-0.0008 (6)	-0.0028 (6)	-0.0015 (6)
O3	0.0076 (7)	0.0110 (8)	0.0077 (8)	-0.0033 (6)	0.0007 (6)	-0.0032 (6)

O4	0.0098 (7)	0.0072 (7)	0.0096 (8)	-0.0012 (6)	0.0008 (6)	-0.0040 (6)
O5	0.0173 (8)	0.0063 (7)	0.0093 (8)	-0.0012 (6)	-0.0063 (7)	-0.0002 (6)
O6	0.0072 (7)	0.0139 (8)	0.0096 (8)	-0.0017 (6)	0.0012 (6)	-0.0040 (6)
O7	0.0096 (7)	0.0067 (7)	0.0098 (8)	-0.0003 (6)	-0.0032 (6)	-0.0016 (6)
C1	0.0087 (9)	0.0066 (10)	0.0056 (10)	-0.0009 (8)	-0.0011 (8)	-0.0005 (8)

Geometric parameters (Å, °)

Cs1—O5 ⁱ	3.119 (3)	P2—O6	1.5158 (18)
Cs1—O3 ⁱⁱ	3.165 (2)	P2—O5	1.5663 (19)
Cs1—O2 ⁱⁱⁱ	3.162 (2)	P2—C1	1.799 (2)
Cs1—O2 ^{iv}	3.173 (2)	O1—H1A	0.922 (17)
Cs1—O6 ^v	3.192 (2)	O1—H1B	0.846 (19)
Cs1—O4 ⁱⁱⁱ	3.485 (2)	O2—Cs1 ⁱⁱⁱ	3.162 (2)
Cs1—O7 ^{vi}	3.490 (2)	O2—Cs1 ^{vii}	3.173 (2)
Co1—O2 ⁱⁱⁱ	2.0761 (18)	O3—Cs1 ⁱⁱ	3.165 (2)
Co1—O2	2.0761 (18)	O3—H2	0.840 (19)
Co1—O1 ⁱⁱⁱ	2.1209 (19)	O4—Cs1 ⁱⁱⁱ	3.485 (2)
Co1—O1	2.1209 (19)	O5—Cs1 ^{viii}	3.119 (3)
Co1—O7 ⁱⁱⁱ	2.1272 (19)	O5—H5A	0.83 (2)
Co1—O7	2.1272 (19)	O5—H5B	0.85 (2)
P1—O2	1.5087 (18)	O6—Cs1 ^v	3.192 (2)
P1—O4	1.5158 (18)	O7—Cs1 ^{vi}	3.490 (2)
P1—O3	1.5732 (18)	C1—H3	0.963 (14)
P1—C1	1.796 (3)	C1—H4	0.951 (14)
P2—O7	1.5099 (18)		
O5 ⁱ —Cs1—O3 ⁱⁱ	91.51 (5)	O7—Co1—Cs1 ⁱⁱⁱ	128.78 (6)
O5 ⁱ —Cs1—O2 ⁱⁱⁱ	113.53 (4)	O2 ⁱⁱⁱ —Co1—Cs1 ^{vi}	45.80 (6)
O3 ⁱⁱ —Cs1—O2 ⁱⁱⁱ	103.25 (7)	O2—Co1—Cs1 ^{vi}	134.20 (6)
O5 ⁱ —Cs1—O2 ^{iv}	125.59 (5)	O1 ⁱⁱⁱ —Co1—Cs1 ^{vi}	60.18 (5)
O3 ⁱⁱ —Cs1—O2 ^{iv}	120.31 (5)	O1—Co1—Cs1 ^{vi}	119.82 (5)
O2 ⁱⁱⁱ —Cs1—O2 ^{iv}	101.31 (5)	O7 ⁱⁱⁱ —Co1—Cs1 ^{vi}	125.13 (6)
O5 ⁱ —Cs1—O6 ^v	82.95 (5)	O7—Co1—Cs1 ^{vi}	54.87 (6)
O3 ⁱⁱ —Cs1—O6 ^v	81.36 (7)	Cs1 ⁱⁱⁱ —Co1—Cs1 ^{vi}	123.10 (4)
O2 ⁱⁱⁱ —Cs1—O6 ^v	162.47 (4)	O2 ⁱⁱⁱ —Co1—Cs1 ^{vii}	134.20 (6)
O2 ^{iv} —Cs1—O6 ^v	62.52 (5)	O2—Co1—Cs1 ^{vii}	45.80 (6)
O5 ⁱ —Cs1—O4 ⁱⁱⁱ	73.40 (5)	O1 ⁱⁱⁱ —Co1—Cs1 ^{vii}	119.82 (5)
O3 ⁱⁱ —Cs1—O4 ⁱⁱⁱ	124.24 (6)	O1—Co1—Cs1 ^{vii}	60.18 (5)
O2 ⁱⁱⁱ —Cs1—O4 ⁱⁱⁱ	44.75 (5)	O7 ⁱⁱⁱ —Co1—Cs1 ^{vii}	54.87 (6)
O2 ^{iv} —Cs1—O4 ⁱⁱⁱ	111.44 (5)	O7—Co1—Cs1 ^{vii}	125.13 (6)
O6 ^v —Cs1—O4 ⁱⁱⁱ	144.68 (4)	Cs1 ⁱⁱⁱ —Co1—Cs1 ^{vii}	56.90 (4)
O5 ⁱ —Cs1—O7 ^{vi}	94.14 (4)	Cs1 ^{vi} —Co1—Cs1 ^{vii}	180
O3 ⁱⁱ —Cs1—O7 ^{vi}	170.61 (4)	O2—P1—O4	114.76 (10)
O2 ⁱⁱⁱ —Cs1—O7 ^{vi}	81.34 (7)	O2—P1—O3	109.86 (10)
O2 ^{iv} —Cs1—O7 ^{vi}	50.37 (5)	O4—P1—O3	107.73 (9)
O6 ^v —Cs1—O7 ^{vi}	91.89 (7)	O2—P1—C1	109.55 (10)
O4 ⁱⁱⁱ —Cs1—O7 ^{vi}	64.74 (6)	O4—P1—C1	107.05 (11)

O5 ⁱ —Cs1—C1 ⁱ	44.75 (6)	O3—P1—C1	107.62 (10)
O3 ⁱⁱ —Cs1—C1 ⁱ	72.92 (5)	O2—P1—Cs1 ⁱⁱⁱ	51.16 (7)
O2 ⁱⁱⁱ —Cs1—C1 ⁱ	78.47 (6)	O4—P1—Cs1 ⁱⁱⁱ	63.66 (7)
O2 ^{iv} —Cs1—C1 ⁱ	165.97 (5)	O3—P1—Cs1 ⁱⁱⁱ	124.40 (8)
O6 ^v —Cs1—C1 ⁱ	118.91 (5)	C1—P1—Cs1 ⁱⁱⁱ	127.77 (8)
O4 ⁱⁱⁱ —Cs1—C1 ⁱ	58.52 (5)	O2—P1—Cs1 ^{vii}	49.06 (7)
O7 ^{vi} —Cs1—C1 ⁱ	116.25 (5)	O4—P1—Cs1 ^{vii}	103.57 (8)
O5 ⁱ —Cs1—O1 ^{iv}	74.68 (4)	O3—P1—Cs1 ^{vii}	68.88 (7)
O3 ⁱⁱ —Cs1—O1 ^{iv}	126.32 (5)	C1—P1—Cs1 ^{vii}	148.54 (8)
O2 ⁱⁱⁱ —Cs1—O1 ^{iv}	130.04 (5)	Cs1 ⁱⁱⁱ —P1—Cs1 ^{vii}	61.70 (2)
O2 ^{iv} —Cs1—O1 ^{iv}	50.97 (5)	O7—P2—O6	114.83 (11)
O6 ^v —Cs1—O1 ^{iv}	46.06 (5)	O7—P2—O5	111.69 (10)
O4 ⁱⁱⁱ —Cs1—O1 ^{iv}	101.41 (4)	O6—P2—O5	109.44 (10)
O7 ^{vi} —Cs1—O1 ^{iv}	48.72 (4)	O7—P2—C1	110.57 (10)
C1 ⁱ —Cs1—O1 ^{iv}	118.60 (6)	O6—P2—C1	108.09 (10)
O5 ⁱ —Cs1—O3 ^{iv}	125.42 (5)	O5—P2—C1	101.32 (11)
O3 ⁱⁱ —Cs1—O3 ^{iv}	78.65 (6)	O7—P2—Cs1 ^v	124.05 (8)
O2 ⁱⁱⁱ —Cs1—O3 ^{iv}	121.02 (5)	O5—P2—Cs1 ^v	65.52 (7)
O2 ^{iv} —Cs1—O3 ^{iv}	42.33 (5)	C1—P2—Cs1 ^v	125.05 (8)
O6 ^v —Cs1—O3 ^{iv}	42.63 (4)	O7—P2—Cs1 ^{viii}	118.54 (7)
O4 ⁱⁱⁱ —Cs1—O3 ^{iv}	152.49 (4)	O6—P2—Cs1 ^{viii}	125.77 (7)
O7 ^{vi} —Cs1—O3 ^{iv}	91.97 (5)	C1—P2—Cs1 ^{viii}	61.03 (8)
C1 ⁱ —Cs1—O3 ^{iv}	148.87 (5)	Cs1 ^v —P2—Cs1 ^{viii}	94.916 (19)
O1 ^{iv} —Cs1—O3 ^{iv}	69.55 (5)	Co1—O1—Cs1 ^{vii}	89.88 (6)
O5 ⁱ —Cs1—O5 ^v	55.64 (5)	Co1—O1—H1A	122 (2)
O3 ⁱⁱ —Cs1—O5 ^v	56.50 (6)	Cs1 ^{vii} —O1—H1A	81 (2)
O2 ⁱⁱⁱ —Cs1—O5 ^v	153.15 (4)	Co1—O1—H1B	120 (3)
O2 ^{iv} —Cs1—O5 ^v	104.38 (5)	Cs1 ^{vii} —O1—H1B	66 (3)
O6 ^v —Cs1—O5 ^v	41.87 (4)	H1A—O1—H1B	108 (3)
O4 ⁱⁱⁱ —Cs1—O5 ^v	128.47 (5)	P1—O2—Co1	135.98 (10)
O7 ^{vi} —Cs1—O5 ^v	121.54 (5)	P1—O2—Cs1 ⁱⁱⁱ	107.03 (9)
C1 ⁱ —Cs1—O5 ^v	78.50 (5)	Co1—O2—Cs1 ⁱⁱⁱ	103.97 (7)
O1 ^{iv} —Cs1—O5 ^v	73.92 (4)	P1—O2—Cs1 ^{vii}	109.88 (8)
O3 ^{iv} —Cs1—O5 ^v	75.37 (5)	Co1—O2—Cs1 ^{vii}	106.22 (8)
O5 ⁱ —Cs1—P1 ⁱⁱⁱ	94.28 (4)	Cs1 ⁱⁱⁱ —O2—Cs1 ^{vii}	78.69 (5)
O3 ⁱⁱ —Cs1—P1 ⁱⁱⁱ	115.59 (6)	P1—O3—Cs1 ⁱⁱ	150.94 (9)
O2 ⁱⁱⁱ —Cs1—P1 ⁱⁱⁱ	21.82 (3)	P1—O3—Cs1 ^{vii}	87.66 (8)
O2 ^{iv} —Cs1—P1 ⁱⁱⁱ	106.97 (4)	Cs1 ⁱⁱ —O3—Cs1 ^{vii}	101.35 (6)
O6 ^v —Cs1—P1 ⁱⁱⁱ	162.96 (3)	P1—O3—H2	108 (3)
O4 ⁱⁱⁱ —Cs1—P1 ⁱⁱⁱ	22.94 (3)	Cs1 ⁱⁱ —O3—H2	101 (3)
O7 ^{vi} —Cs1—P1 ⁱⁱⁱ	71.49 (6)	Cs1 ^{vii} —O3—H2	59 (3)
C1 ⁱ —Cs1—P1 ⁱⁱⁱ	67.80 (4)	P1—O4—Cs1 ⁱⁱⁱ	93.39 (8)
O1 ^{iv} —Cs1—P1 ⁱⁱⁱ	116.98 (4)	P2—O5—Cs1 ^{viii}	120.07 (9)
O3 ^{iv} —Cs1—P1 ⁱⁱⁱ	138.50 (3)	P2—O5—Cs1 ^v	91.93 (8)
O5 ^v —Cs1—P1 ⁱⁱⁱ	145.78 (4)	Cs1 ^{viii} —O5—Cs1 ^v	124.36 (5)
O2 ⁱⁱⁱ —Co1—O2	180	P2—O5—H5A	118 (5)
O2 ⁱⁱⁱ —Co1—O1 ⁱⁱⁱ	90.76 (8)	Cs1 ^v —O5—H5A	99 (5)
O2—Co1—O1 ⁱⁱⁱ	89.24 (8)	P2—O5—H5B	117 (5)

O2 ⁱⁱⁱ —Co1—O1	89.24 (8)	Cs1 ^{viii} —O5—H5B	103 (5)
O2—Co1—O1	90.76 (8)	Cs1 ^v —O5—H5B	100 (5)
O1 ⁱⁱⁱ —Co1—O1	180	H5A—O5—H5B	121 (6)
O2 ⁱⁱⁱ —Co1—O7 ⁱⁱⁱ	94.65 (8)	P2—O6—Cs1 ^v	115.52 (9)
O2—Co1—O7 ⁱⁱⁱ	85.35 (8)	P2—O7—Co1	126.62 (10)
O1 ⁱⁱⁱ —Co1—O7 ⁱⁱⁱ	91.47 (7)	P2—O7—Cs1 ^{vi}	131.62 (9)
O1—Co1—O7 ⁱⁱⁱ	88.53 (7)	Co1—O7—Cs1 ^{vi}	95.23 (7)
O2 ⁱⁱⁱ —Co1—O7	85.35 (8)	P1—C1—P2	116.79 (13)
O2—Co1—O7	94.65 (8)	P1—C1—Cs1 ^{viii}	149.47 (11)
O1 ⁱⁱⁱ —Co1—O7	88.53 (7)	P2—C1—Cs1 ^{viii}	93.21 (10)
O1—Co1—O7	91.47 (7)	P1—C1—H3	108.1 (19)
O7 ⁱⁱⁱ —Co1—O7	180.00 (9)	P2—C1—H3	110.1 (19)
O2 ⁱⁱⁱ —Co1—Cs1 ⁱⁱⁱ	132.78 (6)	Cs1 ^{viii} —C1—H3	62.8 (19)
O2—Co1—Cs1 ⁱⁱⁱ	47.22 (6)	P1—C1—H4	104.3 (19)
O1 ⁱⁱⁱ —Co1—Cs1 ⁱⁱⁱ	63.15 (5)	P2—C1—H4	105.3 (19)
O1—Co1—Cs1 ⁱⁱⁱ	116.85 (5)	Cs1 ^{viii} —C1—H4	59.3 (18)
O7 ⁱⁱⁱ —Co1—Cs1 ⁱⁱⁱ	51.22 (6)	H3—C1—H4	112 (3)
O2 ⁱⁱⁱ —Co1—O1—Cs1 ^{vii}	−144.21 (6)	C1—P2—O5—Cs1 ^{viii}	−8.70 (12)
O2—Co1—O1—Cs1 ^{vii}	35.79 (6)	Cs1 ^v —P2—O5—Cs1 ^{viii}	−132.22 (9)
O7 ⁱⁱⁱ —Co1—O1—Cs1 ^{vii}	−49.55 (6)	O7—P2—O5—Cs1 ^v	−118.75 (9)
O7—Co1—O1—Cs1 ^{vii}	130.45 (6)	O6—P2—O5—Cs1 ^v	9.54 (9)
Cs1 ⁱⁱⁱ —Co1—O1—Cs1 ^{vii}	−5.22 (5)	C1—P2—O5—Cs1 ^v	123.53 (8)
Cs1 ^{vi} —Co1—O1—Cs1 ^{vii}	180	Cs1 ^{viii} —P2—O5—Cs1 ^v	132.22 (9)
O4—P1—O2—Co1	130.04 (13)	O7—P2—O6—Cs1 ^v	114.18 (10)
O3—P1—O2—Co1	−108.42 (14)	O5—P2—O6—Cs1 ^v	−12.34 (11)
C1—P1—O2—Co1	9.60 (17)	C1—P2—O6—Cs1 ^v	−121.88 (10)
Cs1 ⁱⁱⁱ —P1—O2—Co1	132.87 (17)	Cs1 ^{viii} —P2—O6—Cs1 ^v	−55.00 (10)
Cs1 ^{vii} —P1—O2—Co1	−143.29 (18)	O6—P2—O7—Co1	80.65 (14)
O4—P1—O2—Cs1 ⁱⁱⁱ	−2.83 (11)	O5—P2—O7—Co1	−154.00 (11)
O3—P1—O2—Cs1 ⁱⁱⁱ	118.71 (9)	C1—P2—O7—Co1	−41.97 (15)
C1—P1—O2—Cs1 ⁱⁱⁱ	−123.27 (9)	Cs1 ^v —P2—O7—Co1	131.64 (9)
Cs1 ^{vii} —P1—O2—Cs1 ⁱⁱⁱ	83.84 (8)	Cs1 ^{viii} —P2—O7—Co1	−109.34 (10)
O4—P1—O2—Cs1 ^{vii}	−86.67 (11)	O6—P2—O7—Cs1 ^{vi}	−135.06 (11)
O3—P1—O2—Cs1 ^{vii}	34.87 (11)	O5—P2—O7—Cs1 ^{vi}	−9.70 (14)
C1—P1—O2—Cs1 ^{vii}	152.89 (9)	C1—P2—O7—Cs1 ^{vi}	102.32 (13)
Cs1 ⁱⁱⁱ —P1—O2—Cs1 ^{vii}	−83.84 (8)	Cs1 ^v —P2—O7—Cs1 ^{vi}	−84.06 (12)
O1 ⁱⁱⁱ —Co1—O2—P1	−80.87 (15)	Cs1 ^{viii} —P2—O7—Cs1 ^{vi}	34.96 (13)
O1—Co1—O2—P1	99.13 (15)	O2 ⁱⁱⁱ —Co1—O7—P2	−168.93 (12)
O7 ⁱⁱⁱ —Co1—O2—P1	−172.41 (15)	O2—Co1—O7—P2	11.07 (12)
O7—Co1—O2—P1	7.59 (15)	O1 ⁱⁱⁱ —Co1—O7—P2	100.19 (13)
Cs1 ⁱⁱⁱ —Co1—O2—P1	−133.77 (17)	O1—Co1—O7—P2	−79.81 (13)
Cs1 ^{vi} —Co1—O2—P1	−35.83 (17)	Cs1 ⁱⁱⁱ —Co1—O7—P2	47.08 (14)
Cs1 ^{vii} —Co1—O2—P1	144.17 (17)	Cs1 ^{vi} —Co1—O7—P2	154.02 (14)
O1 ⁱⁱⁱ —Co1—O2—Cs1 ⁱⁱⁱ	52.90 (7)	Cs1 ^{vii} —Co1—O7—P2	−25.98 (14)
O1—Co1—O2—Cs1 ⁱⁱⁱ	−127.10 (7)	O2 ⁱⁱⁱ —Co1—O7—Cs1 ^{vi}	37.05 (6)
O7 ⁱⁱⁱ —Co1—O2—Cs1 ⁱⁱⁱ	−38.64 (6)	O2—Co1—O7—Cs1 ^{vi}	−142.95 (6)
O7—Co1—O2—Cs1 ⁱⁱⁱ	141.36 (6)	O1 ⁱⁱⁱ —Co1—O7—Cs1 ^{vi}	−53.82 (6)

Cs1 ^{vi} —Co1—O2—Cs1 ⁱⁱⁱ	97.94 (6)	O1—Co1—O7—Cs1 ^{vi}	126.18 (6)
Cs1 ^{vii} —Co1—O2—Cs1 ⁱⁱⁱ	-82.06 (6)	Cs1 ⁱⁱⁱ —Co1—O7—Cs1 ^{vi}	-106.94 (4)
O1 ⁱⁱⁱ —Co1—O2—Cs1 ^{vii}	134.96 (7)	Cs1 ^{vii} —Co1—O7—Cs1 ^{vi}	180
O1—Co1—O2—Cs1 ^{vii}	-45.04 (7)	O2—P1—C1—P2	-44.78 (16)
O7 ⁱⁱⁱ —Co1—O2—Cs1 ^{vii}	43.42 (6)	O4—P1—C1—P2	-169.80 (12)
O7—Co1—O2—Cs1 ^{vii}	-136.58 (6)	O3—P1—C1—P2	74.63 (14)
Cs1 ⁱⁱⁱ —Co1—O2—Cs1 ^{vii}	82.06 (6)	Cs1 ⁱⁱⁱ —P1—C1—P2	-100.25 (12)
Cs1 ^{vi} —Co1—O2—Cs1 ^{vii}	180	Cs1 ^{vii} —P1—C1—P2	-3.5 (2)
O2—P1—O3—Cs1 ⁱⁱ	-136.98 (17)	O2—P1—C1—Cs1 ^{viii}	123.58 (19)
O4—P1—O3—Cs1 ⁱⁱ	-11.3 (2)	O4—P1—C1—Cs1 ^{viii}	-1.4 (2)
C1—P1—O3—Cs1 ⁱⁱ	103.8 (2)	O3—P1—C1—Cs1 ^{viii}	-117.01 (19)
Cs1 ⁱⁱⁱ —P1—O3—Cs1 ⁱⁱ	-81.1 (2)	Cs1 ⁱⁱⁱ —P1—C1—Cs1 ^{viii}	68.1 (2)
Cs1 ^{vii} —P1—O3—Cs1 ⁱⁱ	-109.40 (19)	Cs1 ^{vii} —P1—C1—Cs1 ^{viii}	164.84 (9)
O2—P1—O3—Cs1 ^{vii}	-27.58 (9)	O7—P2—C1—P1	62.05 (15)
O4—P1—O3—Cs1 ^{vii}	98.08 (9)	O6—P2—C1—P1	-64.42 (15)
C1—P1—O3—Cs1 ^{vii}	-146.80 (8)	O5—P2—C1—P1	-179.41 (12)
Cs1 ⁱⁱⁱ —P1—O3—Cs1 ^{vii}	28.30 (7)	Cs1 ^v —P2—C1—P1	-111.48 (11)
O2—P1—O4—Cs1 ⁱⁱⁱ	2.46 (9)	Cs1 ^{viii} —P2—C1—P1	174.11 (15)
O3—P1—O4—Cs1 ⁱⁱⁱ	-120.24 (8)	O7—P2—C1—Cs1 ^{viii}	-112.06 (8)
C1—P1—O4—Cs1 ⁱⁱⁱ	124.27 (8)	O6—P2—C1—Cs1 ^{viii}	121.47 (8)
Cs1 ^{vii} —P1—O4—Cs1 ⁱⁱⁱ	-48.42 (5)	O5—P2—C1—Cs1 ^{viii}	6.48 (9)
O7—P2—O5—Cs1 ^{viii}	109.03 (10)	Cs1 ^v —P2—C1—Cs1 ^{viii}	74.41 (7)
O6—P2—O5—Cs1 ^{viii}	-122.68 (10)		

Symmetry codes: (i) $x, y, z+1$; (ii) $-x, -y+1, -z+1$; (iii) $-x+1, -y+1, -z+1$; (iv) $x, y+1, z$; (v) $-x, -y+2, -z+1$; (vi) $-x+1, -y+2, -z+1$; (vii) $x, y-1, z$; (viii) $x, y, z-1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1A \cdots O4 ⁱ	0.92 (2)	1.94 (2)	2.860 (3)	173 (3)
O5—H5B \cdots O4 ^{iv}	0.85 (2)	1.69 (3)	2.518 (3)	164 (7)

Symmetry codes: (i) $x, y, z+1$; (iv) $x, y+1, z$.