

Bis[μ -3,5-bis(2-pyridyl)pyrazolato]bis-(hydrogensulfato)dicopper(II) methanol disolvate

Akio Mishima,^a Akira Fuyuhiro,^b Hitoshi Kumagai^c and Satoshi Kawata^{a*}

^aDepartment of Chemistry, Faculty of Science, Fukuoka University, Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan, ^bDepartment of Chemistry, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan, and ^cInstitute for Molecular Science, 38 Nishigounaka Okazaki, Aichi 444-8585, Japan
Correspondence e-mail: kawata@fukuoka-u.ac.jp


Received 10 August 2011; accepted 21 September 2011

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(C-C) = 0.002$ Å;
 R factor = 0.024; wR factor = 0.068; data-to-parameter ratio = 15.3.

The title compound, $[Cu_2(C_{13}H_9N_4)_2(HSO_4)_2] \cdot 2CH_3OH$, consists of discrete centrosymmetric dinuclear complex molecules and methanol solvent molecules. The Cu^{II} atom shows a square-pyramidal coordination geometry and is bonded to four N atoms of the two bis-chelating 3,5-bis(2-pyridyl)-pyrazolate ions ($bpypz^-$) and one O atom of the hydrogensulfate ion. The $bpypz^-$ ligands in the complex molecule are virtually coplanar [dihedral angle between the mean ligand planes = 0.000(1) $^\circ$] with the Cu^{II} atom deviating in opposite directions from their best plane by 0.2080 (12) Å. $\pi-\pi$ stacking interactions between the pyridyl and pyrazole rings [centroid–centroid distance = 3.391 (3) Å] and strong O–H···O hydrogen bonds between the hydrogensulfate ligands and the methanol molecules assemble the molecules into a one-dimensional polymeric structure extending along the a axis. The methanol molecule acts both as an acceptor and a donor in the hydrogen bonding.

Related literature

For metal complexes of 3,5-bis(2-pyridyl)pyrazole, see: Munakata *et al.* (1995); Nakano *et al.* (2004); Du *et al.* (2005); Yoneda, Adachi, Hayami *et al.* (2006); Yoneda, Adachi, Nishio *et al.* (2006); Ishikawa *et al.* (2008, 2010). For an example of a coordinated hydrogensulfate ion, see: Dragancea *et al.* (2008).



Experimental

Crystal data

$[Cu_2(C_{13}H_9N_4)_2(HSO_4)_2] \cdot 2CH_3OH$	$V = 1525.16 (12)$ Å ³
$M_r = 827.82$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 6.0909 (3)$ Å	$\mu = 1.61$ mm ⁻¹
$b = 16.0581 (6)$ Å	$T = 200$ K
$c = 15.6579 (7)$ Å	$0.35 \times 0.04 \times 0.03$ mm
$\beta = 95.2044 (14)$ °	

Data collection

Rigaku R-Axis RAPID diffractometer	16136 measured reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Rigaku, 1995)	2470 independent reflections
$T_{min} = 0.926$, $T_{max} = 0.953$	3262 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$	227 parameters
$wR(F^2) = 0.068$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.39$ e Å ⁻³
3470 reflections	$\Delta\rho_{\text{min}} = -0.51$ e Å ⁻³

Table 1

Selected geometric parameters (Å, °).

Cu1–O1	2.2696 (13)	Cu1–N3 ⁱ	1.9507 (13)
Cu1–N1	2.0865 (13)	Cu1–N4 ⁱ	2.0924 (14)
Cu1–N2	1.9558 (13)		

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku, 2002); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalStructure* (Rigaku, 2010); software used to prepare material for publication: *CrystalStructure*.

This work was supported by funds (No. 101501) from the Central Research Institute of Fukuoka University and Grant-in-Aids for Science Research (No. 22550067) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2397).

References

- Dragancea, D., Addison, A. W., Zeller, M., Thompson, L. K., Hoole, D., Revenco, M. D. & Hunter, A. D. (2008). *Eur. J. Inorg. Chem.* pp. 2530–2536.
- Du, M., Chen, S.-T., Guo, Y.-M., Bu, X.-H. & Ribas, J. (2005). *J. Mol. Struct.*, **737**, 17–21.
- Ishikawa, R., Fuyuhiro, A., Hayami, S., Inoue, K. & Kawata, S. (2008). *J. Mol. Struct.*, **892**, 220–224.
- Ishikawa, R., Nakano, M., Fuyuhiro, A., Takeuchi, T., Kimura, S., Kashiwagi, T., Hagiwara, M., Kindo, K., Kaizaki, S. & Kawata, S. (2010). *Chem. Eur. J.* **16**, 11139–11144.
- Munakata, M., Wu, L. P., Yamamoto, M., Kuroda-Sowa, T., Maekawa, M., Kawata, S. & Kitagawa, S. (1995). *J. Chem. Soc. Dalton Trans.* pp. 4099–4106.
- Nakano, K., Suemura, N., Kawata, S., Fuyuhiro, A., Yagi, T., Nasu, S., Morimoto, S. & Kaizaki, S. (2004). *Dalton Trans.* pp. 982–988.
- Rigaku (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2002). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2010). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yoneda, K., Adachi, K., Hayami, S., Maeda, Y., Katada, M., Fuyuhiro, A., Kawata, S. & Kaizaki, S. (2006). *Chem. Commun.* pp. 45–47.
- Yoneda, K., Adachi, K., Nishio, K., Yamasaki, M., Fuyuhiro, A., Katada, M., Kaizaki, S. & Kawata, S. (2006). *Angew. Chem. Int. Ed.* **45**, 5459–5461.

supporting information

Acta Cryst. (2011). E67, m1523–m1524 [doi:10.1107/S1600536811038700]

Bis[μ -3,5-bis(2-pyridyl)pyrazolato]bis(hydrogensulfato)dicopper(II) methanol disolvate

Akio Mishima, Akira Fuyuhiro, Hitoshi Kumagai and Satoshi Kawata

S1. Comment


3,5-Bis(2-pyridyl)pyrazole[Hbypz] is a versatile ligand in the construction of a series of mononuclear, dinuclear and polynuclear complexes (Munakata *et al.*, 1995; Du *et al.*, 2005; Yoneda *et al.*, 2006; Ishikawa *et al.*, 2010). The dinuclear complexes show the structure where two bpypz²⁻ ions are bridging two metal ions with the axial coordination sites. This kind of dinuclear complexes with transition metal ions were reported previously (Nakano *et al.*, 2004; Yoneda *et al.*, 2006; Ishikawa *et al.*, 2008). The title compound consists of the Cu^{II} dinuclear complex and two methanol solvent molecules. In the dinuclear complex, four N donors from two deprotonated tetradentate bridging bpypz²⁻ ligands form the basal plane (Table 1). The copper(II) ion is penta-coordinated and it is in a slightly deformed square-pyramidal coordination environment with the τ value of 0.001 (Fig. 1). The apical position is occupied by the hydrogensulfate ion. An uncommon feature is that a sulfate ion is actually bound as the hydrogensulfate ion as there are only a few instances of unidentately coordinated hydrogensulfate ion (Dragancea *et al.*, 2008). The $\pi\cdots\pi$ stacking interactions between pyridyl and pyrazole rings [centroid-centroid distance 3.391 (3) Å] and strong hydrogen bonds between the hydrogensulfate ligands and the methanol molecules assemble molecules into a one dimensional polymeric structure extended along the a axis (Table 2, Fig. 2). The methanol molecule acts both as an acceptor and a donor in O-H \cdots O hydrogen bond.

S2. Experimental


A methanolic solution of CuSO₄.6H₂O (5ml, 10 mmol L⁻¹) was transferred to a glass tube, and then a methanolic solution of Hbypz (5ml, 10 mmol L⁻¹) was poured into the glass tube without mixing the solutions. Green crystals began to form at ambient temperature within one week (yield 84%). Elemental analysis: calcd (%) for C₂₈H₂₈Cu₂N₈O₁₀S₂: C 40.62, H 3.41, N 13.54; found: C 40.45, H 3.28, N 13.60.

S3. Refinement


The C-bound hydrogen atoms in the bpypz²⁻ ion and the methyl group of the methanol molecule were placed at calculated positions, C—H 0.950 Å and 0.980 Å respectively, and were treated as riding on their parent atoms with $U_{\text{iso}}(\text{H})$ set to 1.2 $U_{\text{eq}}(\text{C})$. The O-bound hydrogen atoms in the hydrogensulfate ion and the methanol molecule were located in a difference Fourier map. The H-atom coordinates were fixed. The distances were O2—H10 0.90 Å and O5—H11 0.96 Å.

**Figure 1**

ORTEP drawing for the title complex showing 50% probability displacement ellipsoids. Symmetry code for unlabelled atoms: -x, -y+1, -z+1.

**Figure 2**


A fragment of one-dimensional chain structure of the title compound. Hydrogen bonds are shown as dashed lines.

**Figure 3**

Crystal structure for the title compound viewed along the a axis (a) and the b axis (b). The C-bound hydrogen atoms have been omitted.

Bis[μ -3,5-bis(2-pyridyl)pyrazolato]bis(hydrogensulfato)dicopper(II) methanol disolvate

Crystal data



$$M_r = 827.82$$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$$a = 6.0909 (3) \text{ \AA}$$

$$b = 16.0581 (6) \text{ \AA}$$

$$c = 15.6579 (7) \text{ \AA}$$

$$\beta = 95.2044 (14)^\circ$$

$$V = 1525.16 (12) \text{ \AA}^3$$

$$Z = 2$$

$$F(000) = 844.00$$

$$D_x = 1.803 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71075 \text{ \AA}$

Cell parameters from 14158 reflections

$$\theta = 3.4\text{--}27.5^\circ$$

$$\mu = 1.61 \text{ mm}^{-1}$$

$$T = 200 \text{ K}$$

Column, green

$$0.35 \times 0.04 \times 0.03 \text{ mm}$$

Data collection

Rigaku R-AXIS RAPID
diffractometer

Detector resolution: 10.000 pixels mm^{-1}

ω scans

Absorption correction: multi-scan
(*ABSCOR*; Rigaku, 1995)

$$T_{\min} = 0.926, T_{\max} = 0.953$$

16136 measured reflections

2470 independent reflections

3262 reflections with $F^2 > 2.0\sigma(F^2)$

$$R_{\text{int}} = 0.021$$

$$\theta_{\max} = 27.4^\circ$$

$$h = -7 \rightarrow 7$$

$$k = -20 \rightarrow 20$$

$$l = -20 \rightarrow 20$$

Refinement

Refinement on F^2

$$R[F^2 > 2\sigma(F^2)] = 0.024$$

$$wR(F^2) = 0.068$$

$$S = 1.08$$

3470 reflections

227 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0383P)^2 + 0.881P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.39 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.51 \text{ e \AA}^{-3}$$

*Special details***Geometry.** ENTER SPECIAL DETAILS OF THE MOLECULAR GEOMETRY

Refinement. Refinement was performed using all reflections. The weighted *R*-factor (*wR*) and goodness of fit (*S*) are based on F^2 . *R*-factor (gt) are based on *F*. The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating *R*-factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.28031 (3)	0.520598 (11)	0.578370 (12)	0.01532 (7)
S1	0.20037 (6)	0.55399 (2)	0.78731 (2)	0.01893 (10)
O1	0.1195 (2)	0.56027 (8)	0.69710 (8)	0.0266 (3)
O2	0.4401 (2)	0.58901 (8)	0.78864 (8)	0.0293 (3)
O3	0.2141 (3)	0.46998 (8)	0.81989 (9)	0.0292 (3)
O4	0.0797 (3)	0.60921 (9)	0.84007 (9)	0.0356 (4)
O5	0.7222 (2)	0.55846 (10)	0.91528 (9)	0.0331 (3)
N1	0.4667 (3)	0.42870 (8)	0.64544 (9)	0.0172 (3)
N2	0.0927 (2)	0.42369 (8)	0.55066 (9)	0.0175 (3)
N3	-0.1001 (2)	0.41178 (8)	0.50440 (9)	0.0172 (3)
N4	-0.4729 (3)	0.37148 (8)	0.42106 (9)	0.0180 (3)
C1	0.6627 (3)	0.43350 (10)	0.69186 (11)	0.0222 (4)
C2	0.7564 (3)	0.36790 (11)	0.73994 (11)	0.0243 (4)
C3	0.6436 (3)	0.29342 (11)	0.74028 (11)	0.0249 (4)
C4	0.4411 (3)	0.28646 (10)	0.69234 (11)	0.0221 (4)
C5	0.3577 (3)	0.35468 (10)	0.64594 (10)	0.0168 (3)
C6	0.1491 (3)	0.35246 (9)	0.59261 (10)	0.0170 (3)
C7	-0.0117 (3)	0.29197 (9)	0.57300 (10)	0.0186 (3)
C8	-0.1656 (3)	0.33282 (9)	0.51668 (10)	0.0167 (3)
C9	-0.3750 (3)	0.30969 (9)	0.47033 (10)	0.0169 (3)
C10	-0.4679 (3)	0.23107 (10)	0.47581 (11)	0.0226 (4)
C11	-0.6675 (3)	0.21423 (11)	0.42906 (12)	0.0262 (4)
C12	-0.7683 (3)	0.27685 (11)	0.37925 (12)	0.0256 (4)
C13	-0.6672 (3)	0.35401 (11)	0.37704 (11)	0.0232 (4)
C14	0.7361 (4)	0.47382 (13)	0.94479 (15)	0.0361 (5)
H1	0.7415	0.4846	0.6918	0.0267*
H2	0.8955	0.3741	0.7720	0.0292*
H3	0.7036	0.2475	0.7729	0.0299*
H4	0.3610	0.2357	0.6914	0.0266*
H5	-0.0154	0.2363	0.5932	0.0223*
H6	-0.3957	0.1894	0.5111	0.0272*
H7	-0.7334	0.1608	0.4313	0.0314*
H8	-0.9054	0.2671	0.3469	0.0308*
H9	-0.7386	0.3966	0.3428	0.0278*
H10	0.5306	0.5733	0.8344	0.0351*
H11	0.8529	0.5697	0.8873	0.0397*
H12	0.8658	0.4471	0.9240	0.0434*
H13	0.6029	0.4436	0.9229	0.0434*
H14	0.7489	0.4730	1.0076	0.0434*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.01457 (11)	0.01325 (11)	0.01755 (11)	-0.00117 (6)	-0.00178 (7)	0.00169 (6)
S1	0.02020 (19)	0.01756 (19)	0.01917 (19)	-0.00183 (14)	0.00250 (15)	-0.00195 (14)
O1	0.0213 (6)	0.0362 (7)	0.0219 (6)	0.0049 (6)	-0.0005 (5)	-0.0002 (5)
O2	0.0235 (6)	0.0335 (7)	0.0299 (7)	-0.0093 (6)	-0.0023 (5)	0.0060 (6)
O3	0.0320 (7)	0.0207 (6)	0.0344 (8)	-0.0032 (5)	0.0007 (6)	0.0053 (5)
O4	0.0447 (8)	0.0269 (7)	0.0380 (8)	-0.0005 (6)	0.0189 (7)	-0.0097 (6)
O5	0.0244 (7)	0.0382 (8)	0.0362 (8)	-0.0030 (6)	-0.0003 (6)	0.0022 (6)
N1	0.0175 (6)	0.0153 (6)	0.0187 (7)	0.0016 (5)	0.0003 (5)	-0.0006 (5)
N2	0.0169 (6)	0.0155 (6)	0.0195 (7)	-0.0008 (5)	-0.0028 (5)	0.0018 (5)
N3	0.0171 (6)	0.0146 (6)	0.0191 (7)	-0.0022 (5)	-0.0022 (5)	0.0017 (5)
N4	0.0166 (6)	0.0173 (6)	0.0202 (7)	-0.0014 (5)	0.0014 (5)	0.0007 (5)
C1	0.0189 (8)	0.0202 (8)	0.0269 (9)	0.0003 (6)	-0.0019 (7)	-0.0028 (7)
C2	0.0204 (8)	0.0268 (9)	0.0244 (9)	0.0040 (7)	-0.0052 (7)	-0.0028 (7)
C3	0.0255 (9)	0.0247 (9)	0.0233 (9)	0.0054 (7)	-0.0043 (7)	0.0042 (7)
C4	0.0236 (8)	0.0190 (8)	0.0232 (8)	0.0006 (7)	-0.0016 (7)	0.0035 (6)
C5	0.0182 (7)	0.0164 (7)	0.0161 (7)	0.0013 (6)	0.0022 (6)	-0.0003 (6)
C6	0.0182 (7)	0.0154 (7)	0.0174 (7)	0.0015 (6)	0.0012 (6)	0.0021 (6)
C7	0.0205 (8)	0.0149 (7)	0.0203 (8)	-0.0004 (6)	0.0009 (6)	0.0022 (6)
C8	0.0183 (7)	0.0141 (7)	0.0178 (7)	-0.0015 (6)	0.0026 (6)	0.0006 (6)
C9	0.0175 (7)	0.0160 (7)	0.0173 (7)	-0.0006 (6)	0.0030 (6)	-0.0011 (6)
C10	0.0238 (8)	0.0172 (7)	0.0267 (9)	-0.0021 (7)	0.0010 (7)	0.0006 (6)
C11	0.0254 (9)	0.0208 (8)	0.0322 (10)	-0.0082 (7)	0.0020 (7)	-0.0031 (7)
C12	0.0186 (8)	0.0290 (9)	0.0286 (9)	-0.0070 (7)	-0.0020 (7)	-0.0023 (7)
C13	0.0185 (8)	0.0248 (8)	0.0254 (8)	-0.0021 (7)	-0.0027 (7)	0.0026 (7)
C14	0.0367 (11)	0.0384 (12)	0.0331 (11)	-0.0026 (9)	0.0014 (8)	0.0057 (9)

Geometric parameters (\AA , $^\circ$)

Cu1—O1	2.2696 (13)	C6—C7	1.394 (2)
Cu1—N1	2.0865 (13)	C7—C8	1.392 (2)
Cu1—N2	1.9558 (13)	C8—C9	1.457 (2)
Cu1—N3 ⁱ	1.9507 (13)	C9—C10	1.389 (3)
Cu1—N4 ⁱ	2.0924 (14)	C10—C11	1.387 (3)
S1—O1	1.4566 (13)	C11—C12	1.382 (3)
S1—O2	1.5630 (13)	C12—C13	1.385 (3)
S1—O3	1.4420 (14)	O2—H10	0.900
S1—O4	1.4557 (16)	O5—H11	0.960
O5—C14	1.436 (3)	C1—H1	0.950
N1—C1	1.342 (2)	C2—H2	0.950
N1—C5	1.362 (2)	C3—H3	0.950
N2—N3	1.3368 (18)	C4—H4	0.950
N2—C6	1.348 (2)	C7—H5	0.950
N3—C8	1.348 (2)	C10—H6	0.950
N4—C9	1.361 (2)	C11—H7	0.950
N4—C13	1.344 (2)	C12—H8	0.950

C1—C2	1.387 (3)	C13—H9	0.950
C2—C3	1.380 (3)	C14—H12	0.980
C3—C4	1.389 (3)	C14—H13	0.980
C4—C5	1.385 (3)	C14—H14	0.980
C5—C6	1.457 (2)		
O1—Cu1—N1	92.40 (5)	N2—C6—C7	109.97 (13)
O1—Cu1—N2	96.79 (6)	C5—C6—C7	134.63 (14)
O1—Cu1—N3 ⁱ	97.44 (6)	C6—C7—C8	103.33 (13)
O1—Cu1—N4 ⁱ	92.72 (5)	N3—C8—C7	110.04 (13)
N1—Cu1—N2	80.16 (6)	N3—C8—C9	115.19 (13)
N1—Cu1—N3 ⁱ	167.37 (6)	C7—C8—C9	134.77 (14)
N1—Cu1—N4 ⁱ	107.72 (6)	N4—C9—C8	114.53 (13)
N2—Cu1—N3 ⁱ	90.77 (6)	N4—C9—C10	122.53 (14)
N2—Cu1—N4 ⁱ	167.42 (6)	C8—C9—C10	122.94 (14)
N3 ⁱ —Cu1—N4 ⁱ	79.82 (6)	C9—C10—C11	119.12 (15)
O1—S1—O2	102.78 (7)	C10—C11—C12	118.63 (16)
O1—S1—O3	114.30 (8)	C11—C12—C13	119.29 (16)
O1—S1—O4	111.38 (8)	N4—C13—C12	123.13 (16)
O2—S1—O3	107.92 (8)	N1—C1—H1	118.240
O2—S1—O4	107.00 (8)	C2—C1—H1	118.243
O3—S1—O4	112.66 (9)	C1—C2—H2	120.658
Cu1—O1—S1	129.93 (8)	C3—C2—H2	120.663
Cu1—N1—C1	130.43 (11)	C2—C3—H3	120.467
Cu1—N1—C5	112.14 (10)	C4—C3—H3	120.467
C1—N1—C5	117.20 (14)	C3—C4—H4	120.496
Cu1—N2—N3	134.51 (11)	C5—C4—H4	120.484
Cu1—N2—C6	116.71 (10)	C6—C7—H5	128.336
N3—N2—C6	108.36 (13)	C8—C7—H5	128.337
Cu1 ⁱ —N3—N2	133.96 (11)	C9—C10—H6	120.442
Cu1 ⁱ —N3—C8	117.54 (10)	C11—C10—H6	120.440
N2—N3—C8	108.31 (13)	C10—C11—H7	120.683
Cu1 ⁱ —N4—C9	112.64 (10)	C12—C11—H7	120.692
Cu1 ⁱ —N4—C13	129.97 (12)	C11—C12—H8	120.351
C9—N4—C13	117.30 (14)	C13—C12—H8	120.364
N1—C1—C2	123.52 (15)	N4—C13—H9	118.439
C1—C2—C3	118.68 (16)	C12—C13—H9	118.430
C2—C3—C4	119.07 (16)	O5—C14—H12	109.477
C3—C4—C5	119.02 (15)	O5—C14—H13	109.477
N1—C5—C4	122.52 (14)	O5—C14—H14	109.472
N1—C5—C6	114.75 (14)	H12—C14—H13	109.472
C4—C5—C6	122.73 (15)	H12—C14—H14	109.469
N2—C6—C5	115.40 (13)	H13—C14—H14	109.460
O1—Cu1—N1—C1	-85.61 (12)	Cu1—N2—C6—C5	-6.86 (18)
O1—Cu1—N1—C5	88.62 (9)	Cu1—N2—C6—C7	173.75 (9)
N1—Cu1—O1—S1	26.68 (10)	N3—N2—C6—C5	179.56 (12)
O1—Cu1—N2—N3	88.20 (13)	N3—N2—C6—C7	0.17 (17)

O1—Cu1—N2—C6	−83.23 (10)	C6—N2—N3—Cu1 ⁱ	−174.97 (13)
N2—Cu1—O1—S1	107.06 (10)	C6—N2—N3—C8	−0.31 (17)
O1—Cu1—N3 ⁱ —N2 ⁱ	−87.66 (13)	Cu1 ⁱ —N3—C8—C7	176.01 (9)
O1—Cu1—N3 ⁱ —C8 ⁱ	86.62 (10)	Cu1 ⁱ —N3—C8—C9	−4.37 (18)
N3 ⁱ —Cu1—O1—S1	−161.27 (10)	N2—N3—C8—C7	0.34 (17)
O1—Cu1—N4 ⁱ —C9 ⁱ	−92.61 (9)	N2—N3—C8—C9	179.96 (12)
O1—Cu1—N4 ⁱ —C13 ⁱ	83.70 (12)	Cu1 ⁱ —N4—C9—C8	3.47 (16)
N4 ⁱ —Cu1—O1—S1	−81.19 (10)	Cu1 ⁱ —N4—C9—C10	−176.54 (10)
N1—Cu1—N2—N3	179.47 (14)	Cu1 ⁱ —N4—C13—C12	175.54 (10)
N1—Cu1—N2—C6	8.03 (9)	C9—N4—C13—C12	−0.6 (3)
N2—Cu1—N1—C1	177.90 (13)	C13—N4—C9—C8	−179.71 (13)
N2—Cu1—N1—C5	−7.86 (9)	C13—N4—C9—C10	0.3 (3)
N1—Cu1—N4 ⁱ —C9 ⁱ	174.00 (8)	N1—C1—C2—C3	0.2 (3)
N1—Cu1—N4 ⁱ —C13 ⁱ	−9.69 (14)	C1—C2—C3—C4	0.3 (3)
N4 ⁱ —Cu1—N1—C1	8.01 (14)	C2—C3—C4—C5	−0.3 (3)
N4 ⁱ —Cu1—N1—C5	−177.75 (8)	C3—C4—C5—N1	−0.0 (3)
N2—Cu1—N3 ⁱ —N2 ⁱ	9.29 (14)	C3—C4—C5—C6	178.73 (14)
N2—Cu1—N3 ⁱ —C8 ⁱ	−176.42 (10)	N1—C5—C6—N2	−0.2 (2)
N3 ⁱ —Cu1—N2—N3	−9.38 (14)	N1—C5—C6—C7	178.96 (15)
N3 ⁱ —Cu1—N2—C6	179.19 (10)	C4—C5—C6—N2	−179.10 (14)
N3 ⁱ —Cu1—N4 ⁱ —C9 ⁱ	4.46 (9)	C4—C5—C6—C7	0.1 (3)
N3 ⁱ —Cu1—N4 ⁱ —C13 ⁱ	−179.23 (13)	N2—C6—C7—C8	0.03 (17)
N4 ⁱ —Cu1—N3 ⁱ —N2 ⁱ	−179.10 (14)	C5—C6—C7—C8	−179.20 (17)
N4 ⁱ —Cu1—N3 ⁱ —C8 ⁱ	−4.82 (9)	C6—C7—C8—N3	−0.22 (17)
O2—S1—O1—Cu1	47.83 (11)	C6—C7—C8—C9	−179.74 (16)
O3—S1—O1—Cu1	−68.82 (12)	N3—C8—C9—N4	0.3 (2)
O4—S1—O1—Cu1	162.08 (10)	N3—C8—C9—C10	−179.68 (13)
Cu1—N1—C1—C2	173.45 (10)	C7—C8—C9—N4	179.81 (17)
Cu1—N1—C5—C4	−174.59 (10)	C7—C8—C9—C10	−0.2 (3)
Cu1—N1—C5—C6	6.54 (16)	N4—C9—C10—C11	0.4 (3)
C1—N1—C5—C4	0.5 (3)	C8—C9—C10—C11	−179.63 (14)
C1—N1—C5—C6	−178.39 (13)	C9—C10—C11—C12	−0.7 (3)
C5—N1—C1—C2	−0.5 (3)	C10—C11—C12—C13	0.4 (3)
Cu1—N2—N3—Cu1 ⁱ	13.1 (3)	C11—C12—C13—N4	0.3 (3)
Cu1—N2—N3—C8	−172.25 (11)		

Symmetry code: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

D—H···A	D—H	H···A	D···A	D—H···A
O2—H10···O5	0.90	1.66	2.5509 (18)	170
O5—H11···O4 ⁱⁱ	0.96	1.74	2.694 (2)	169

Symmetry code: (ii) $x+1, y, z$.