

μ_4 -Orthothiocarbonato-tetrakis[tri-carbonyliron(I)](2 Fe—Fe)

Yao-Cheng Shi,^{a*} Huan-Ren Cheng,^a Li-Min Yuan^b and Qian-Kun Li^c

^aCollege of Chemistry and Chemical Engineering, Yangzhou University, Yangzhou 225002, People's Republic of China, ^bTesting Center, Yangzhou University, Yangzhou 225009, People's Republic of China, and ^cHubei Research Institute of Geophysics Survey and Design, Wuhan 430056, People's Republic of China
Correspondence e-mail: ycshi@yzu.edu.cn


Received 28 September 2011; accepted 11 October 2011

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{O}-\text{C}) = 0.004$ Å; R factor = 0.026; wR factor = 0.070; data-to-parameter ratio = 17.2.

The fused bis-butterfly-shaped title compound, $[\text{Fe}_4(\text{CS}_4)(\text{CO})_{12}]$, possesses an orthothiocarbonato (CS_4^{4-}) ligand that acts as a bridge between two $\text{Fe}_2(\text{CO})_6$ units. A short intramolecular $\text{S} \cdots \text{S}$ contact [2.6984 (8) and 2.6977 (8) Å] occurs in each $\text{S}_2\text{Fe}_2(\text{CO})_6$ fragment.

Related literature

For general background to related complexes, see: Mathur *et al.* (2009). For uses of $R_3\text{P}/\text{CS}_2$ in coordination chemistry and organometallic chemistry, see: Galindo *et al.* (1999). For the synthesis of butterfly $\text{S}_2\text{Fe}_2(\text{CO})_6$ complexes, see: Song (2005). For related structures, see: Shaver *et al.* (1979); Ortega-Alfaro *et al.* (2004).



Experimental

Crystal data

$[\text{Fe}_4(\text{CS}_4)(\text{CO})_{12}]$ $\gamma = 110.0857$ (12) $^\circ$
 $M_r = 699.81$ $V = 1142.2$ (2) Å³
 Triclinic, $P\bar{1}$ $Z = 2$
 $a = 9.0875$ (9) Å Mo $K\alpha$ radiation
 $b = 10.9002$ (11) Å $\mu = 2.91$ mm⁻¹
 $c = 12.6448$ (13) Å $T = 296$ K
 $\alpha = 101.8859$ (12) $^\circ$ $0.15 \times 0.12 \times 0.11$ mm
 $\beta = 92.4964$ (12) $^\circ$

Data collection

Bruker SMART APEX CCD 10006 measured reflections
 diffractometer 5128 independent reflections
 Absorption correction: multi-scan 4237 reflections with $I > 2\sigma(I)$
 (SADABS; Sheldrick, 2004) $R_{\text{int}} = 0.025$
 $T_{\text{min}} = 0.658$, $T_{\text{max}} = 0.721$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$ 298 parameters
 $wR(F^2) = 0.070$ 6 restraints
 $S = 1.04$ $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 5128 reflections $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Selected geometric parameters (Å, $^\circ$).

C13—S1	1.827 (2)	Fe2—S1	2.2723 (6)
C13—S2	1.8300 (19)	Fe2—S2	2.2685 (7)
C13—S3	1.830 (2)	Fe3—S3	2.2676 (6)
C13—S4	1.837 (2)	Fe3—S4	2.2680 (6)
Fe1—S1	2.2730 (6)	Fe3—Fe4	2.5007 (5)
Fe1—S2	2.2688 (7)	Fe4—S3	2.2712 (7)
Fe1—Fe2	2.4949 (5)	Fe4—S4	2.2626 (6)
S1—C13—S2	95.10 (10)	S3—C13—S4	94.73 (9)

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT-Plus* (Bruker, 2003); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009) and *WinGX* (Farrugia, 1999); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank the Natural Science Foundation of China (No. 20572091) and the Natural Science Foundation of Jiangsu Province (No. 05KJB150151) for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5239).

References

- Bruker (2002). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2003). *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Burla, M. C., Caliendo, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Galindo, A., Miguel, D. & Perez, J. (1999). *Coord. Chem. Rev.* **193–195**, 643–690.
 Mathur, P., Boodida, S., Ji, R. S. & Mobin, S. M. (2009). *J. Organomet. Chem.* **694**, 3043–3045.
 Ortega-Alfaro, M. C., Hernández, N., Cerna, I., López-Cortés, J. G., Gómez, E., Toscano, R. A. & Alvarez-Toledano, C. (2004). *J. Organomet. Chem.* **689**, 885–893.
 Shaver, A., Fitzpatrick, P. J., Steliou, K. & Butler, I. S. (1979). *J. Am. Chem. Soc.* **101**, 1313–1315.
 Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Song, L.-C. (2005). *Acc. Chem. Res.* **38**, 21–28.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2011). E67, m1534 [doi:10.1107/S1600536811041936]

 μ_4 -Orthothiocarbonato-tetrakis[tricarbonyliron(I)](2 Fe—Fe)

Yao-Cheng Shi, Huan-Ren Cheng, Li-Min Yuan and Qian-Kun Li


S1. Comment

The activation and cleavage of selected bonds of small molecules by transition metal complexes is one of the challenging subjects of recent researches. CS₂ has been shown to undergo a variety of reactions with transition metals, including insertion and disproportionation, and there is a growing interest in the activation of CS₂ from catalytic and biological points of view. The cleavage of the C—S bonds is often observed in various transition metal complexes in which chemistry has been explored for the hydrosulfurization of fossil products. In these complexes, the S²⁻ ion derived from the C—S bond scission functions as a bridging ligand to link metal ions and metal cluster fragments and is generally of use in various cluster growth processes (Mathur *et al.*, 2009).

Interestingly, the reaction of Et₃P/CS₂ and Fe₃(CO)₁₂ in THF under inert atmosphere at room temperature leads to the formation of a novel complex (Scheme 1). The molecular structure of the novel complex (Fig. 1) consists of two butterfly Fe₂(CO)₆ units connected by a bridging CS₄ ligand in axial C—S bond fashions similar to the related complex Fe₂(CO)₆(μ -S)₂CH₂ (Shaver *et al.*, 1979). The Fe—Fe bond lengths are 2.4949 (5) and 2.5007 (5) Å and close to 2.485 (1) Å in Fe₂(CO)₆(μ -S)₂CH₂, but slightly shorter than 2.511 (1) Å in the complex Fe₂(CO)₆(μ -SCH₃)₂ (Table 1) (Ortega-Alfaro *et al.*, 2004), the corresponding C—S bond lengths are 1.827 (2), 1.830 (2) and 1.830 (2), 1.837 (2)^o, respectively, which are longer than those in the complex Fe₂(CO)₆(μ -SCH₃)₂. For each S₂Fe₂(CO)₆ butterfly core, the S—C—S bond angle is 95.10 (10) and 94.73 (9)^o and close to 94.55 (3)^o in Fe₂(CO)₆(μ -S)₂CH₂ (Table 1). As compared with 2.744 (1)–2.773 (1) Å in Fe₂(CO)₆(μ -SCH₃)₂, the S⋯S distance (2.6984 (8) and 2.6977 (8) Å) indicates an intramolecular short contact in each S₂Fe₂(CO)₆ butterfly core.

S2. Experimental

A THF solution of Et₃P/CS₂ (1 mmol) and Fe₃(CO)₁₂ (1 mmol) under inert atmosphere is stirred for 24 h at room temperature. After removal of the solvent, the mixture was purified by chromatography on silica gel with dichloromethane-petroleum ether (*v/v*, 1:3) as eluant to give the red-orange solid. Single crystals were grown from ether solution of the title compound.

**Figure 1**

The molecule of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

μ_4 -Orthothiocarbonato-tetrakis[tricarbonyliron(I)](2 Fe—Fe)

Crystal data

[Fe₄(CS₄)(CO)₁₂]

$M_r = 699.81$

Triclinic, $P\bar{1}$

$a = 9.0875$ (9) Å

$b = 10.9002$ (11) Å

$c = 12.6448$ (13) Å

$\alpha = 101.8859$ (12)°

$\beta = 92.4964$ (12)°

$\gamma = 110.0857$ (12)°

$V = 1142.2$ (2) Å³

$Z = 2$

$F(000) = 684$

$D_x = 2.035$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4237 reflections

$\theta = 1.7$ – 27.5 °

$\mu = 2.91$ mm⁻¹

$T = 296$ K

Prism, red

$0.15 \times 0.12 \times 0.11$ mm

Data collection

Bruker SMART APEX CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and φ scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2004)

$T_{\min} = 0.658$, $T_{\max} = 0.721$

10006 measured reflections

5128 independent reflections

4237 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 1.7$ °

$h = -11 \rightarrow 11$

$k = -14 \rightarrow 14$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.026$

$wR(F^2) = 0.070$

$S = 1.04$

5128 reflections

298 parameters

6 restraints

Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

$$w = 1/[\sigma^2(F_o^2) + (0.0332P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6561 (3)	0.7935 (2)	0.77984 (18)	0.0440 (5)
C2	0.8526 (3)	0.8863 (3)	0.6347 (2)	0.0506 (6)
C3	0.9578 (3)	0.9654 (3)	0.8400 (2)	0.0495 (6)
C4	1.1219 (3)	0.7705 (3)	0.5832 (2)	0.0576 (7)
C5	1.2260 (3)	0.8567 (3)	0.7869 (2)	0.0499 (6)
C6	1.1533 (3)	0.5863 (3)	0.6883 (2)	0.0520 (6)
C7	0.7715 (3)	0.4656 (3)	1.0044 (2)	0.0518 (6)
C8	0.6869 (3)	0.2015 (3)	0.9062 (2)	0.0571 (7)
C9	0.4644 (3)	0.2955 (2)	0.94124 (17)	0.0461 (6)
C10	0.5449 (3)	0.1072 (3)	0.6532 (2)	0.0533 (6)
C11	0.3306 (3)	0.2117 (3)	0.69401 (19)	0.0538 (6)
C12	0.5320 (3)	0.3000 (2)	0.54991 (19)	0.0461 (5)
C13	0.7801 (2)	0.5349 (2)	0.75626 (15)	0.0326 (4)
Fe1	0.85275 (4)	0.80727 (3)	0.74597 (2)	0.03532 (9)
Fe2	1.06610 (4)	0.71578 (3)	0.70503 (2)	0.03858 (9)
Fe3	0.64765 (4)	0.34558 (3)	0.88425 (2)	0.03604 (9)
Fe4	0.54035 (4)	0.27424 (3)	0.68685 (2)	0.03649 (9)
O1	0.5347 (2)	0.7883 (2)	0.80095 (16)	0.0652 (5)
O2	0.8523 (3)	0.9343 (2)	0.56303 (17)	0.0808 (7)
O3	1.0252 (3)	1.0659 (2)	0.89976 (17)	0.0801 (6)
O4	1.1562 (3)	0.8060 (3)	0.50561 (17)	0.0906 (8)
O5	1.3234 (2)	0.9495 (2)	0.83931 (18)	0.0764 (6)
O6	1.2101 (3)	0.5078 (2)	0.67522 (19)	0.0802 (6)
O7	0.8437 (3)	0.5364 (2)	1.08183 (16)	0.0849 (7)
O8	0.7108 (3)	0.1096 (2)	0.9194 (2)	0.0923 (7)
O9	0.3472 (2)	0.2622 (2)	0.97500 (15)	0.0716 (6)
O10	0.5481 (3)	0.0019 (2)	0.63331 (18)	0.0840 (7)
O11	0.1982 (3)	0.1726 (3)	0.69931 (18)	0.0901 (7)
O12	0.5245 (2)	0.3122 (2)	0.46352 (14)	0.0717 (6)
S1	0.92568 (6)	0.68363 (5)	0.84778 (4)	0.03454 (12)

S2	0.80928 (7)	0.60300 (6)	0.63482 (4)	0.03789 (13)
S3	0.80213 (6)	0.37272 (5)	0.74843 (4)	0.03674 (12)
S4	0.57807 (6)	0.47963 (5)	0.79267 (4)	0.03420 (12)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0464 (14)	0.0398 (13)	0.0478 (12)	0.0174 (11)	0.0032 (10)	0.0116 (10)
C2	0.0482 (15)	0.0460 (14)	0.0586 (14)	0.0159 (12)	0.0062 (12)	0.0168 (12)
C3	0.0446 (14)	0.0430 (14)	0.0568 (14)	0.0113 (12)	0.0098 (11)	0.0103 (12)
C4	0.0577 (17)	0.0620 (17)	0.0653 (17)	0.0292 (14)	0.0259 (14)	0.0241 (14)
C5	0.0368 (14)	0.0484 (15)	0.0656 (16)	0.0140 (12)	0.0142 (12)	0.0163 (13)
C6	0.0452 (15)	0.0523 (15)	0.0623 (15)	0.0179 (12)	0.0168 (12)	0.0189 (13)
C7	0.0536 (16)	0.0528 (15)	0.0469 (13)	0.0150 (13)	0.0004 (11)	0.0155 (12)
C8	0.0601 (17)	0.0545 (16)	0.0633 (16)	0.0241 (14)	0.0094 (13)	0.0218 (13)
C9	0.0510 (15)	0.0453 (14)	0.0360 (11)	0.0114 (12)	0.0063 (10)	0.0065 (10)
C10	0.0601 (17)	0.0402 (14)	0.0502 (14)	0.0095 (12)	0.0063 (12)	0.0057 (11)
C11	0.0481 (16)	0.0526 (16)	0.0440 (13)	0.0020 (13)	0.0040 (11)	0.0040 (11)
C12	0.0436 (14)	0.0401 (13)	0.0434 (13)	0.0046 (11)	0.0000 (10)	0.0049 (10)
C13	0.0291 (11)	0.0319 (10)	0.0341 (10)	0.0089 (9)	0.0031 (8)	0.0057 (8)
Fe1	0.03283 (18)	0.03237 (17)	0.03938 (17)	0.00973 (13)	0.00397 (13)	0.00912 (13)
Fe2	0.03253 (18)	0.03943 (18)	0.04416 (18)	0.01127 (14)	0.01067 (13)	0.01241 (14)
Fe3	0.03703 (18)	0.03452 (17)	0.03534 (16)	0.01093 (14)	0.00442 (13)	0.00903 (13)
Fe4	0.03603 (18)	0.03052 (17)	0.03559 (16)	0.00580 (13)	0.00263 (13)	0.00313 (13)
O1	0.0438 (11)	0.0676 (13)	0.0955 (14)	0.0303 (10)	0.0199 (10)	0.0237 (11)
O2	0.0919 (17)	0.0883 (16)	0.0771 (13)	0.0311 (14)	0.0146 (12)	0.0530 (13)
O3	0.0750 (15)	0.0484 (12)	0.0894 (15)	0.0052 (11)	0.0037 (12)	-0.0119 (11)
O4	0.112 (2)	0.112 (2)	0.0796 (14)	0.0537 (16)	0.0579 (14)	0.0566 (15)
O5	0.0456 (12)	0.0589 (13)	0.1025 (16)	0.0024 (10)	0.0007 (11)	0.0014 (12)
O6	0.0805 (16)	0.0736 (15)	0.1125 (17)	0.0495 (13)	0.0371 (13)	0.0347 (13)
O7	0.0857 (16)	0.0882 (16)	0.0530 (11)	0.0075 (13)	-0.0226 (11)	0.0045 (11)
O8	0.1114 (19)	0.0722 (15)	0.1227 (19)	0.0545 (15)	0.0225 (15)	0.0475 (15)
O9	0.0573 (13)	0.0795 (15)	0.0636 (12)	0.0077 (11)	0.0265 (10)	0.0115 (11)
O10	0.1131 (19)	0.0391 (11)	0.0960 (16)	0.0283 (12)	0.0166 (14)	0.0056 (11)
O11	0.0473 (13)	0.1058 (19)	0.0861 (15)	-0.0053 (12)	0.0126 (11)	0.0119 (14)
O12	0.0788 (14)	0.0732 (14)	0.0428 (10)	0.0033 (11)	-0.0061 (9)	0.0147 (9)
S1	0.0314 (3)	0.0334 (3)	0.0332 (2)	0.0058 (2)	0.0010 (2)	0.0063 (2)
S2	0.0398 (3)	0.0375 (3)	0.0317 (3)	0.0086 (2)	0.0031 (2)	0.0077 (2)
S3	0.0338 (3)	0.0338 (3)	0.0430 (3)	0.0135 (2)	0.0070 (2)	0.0071 (2)
S4	0.0298 (3)	0.0309 (3)	0.0400 (3)	0.0097 (2)	0.0045 (2)	0.0062 (2)

Geometric parameters (Å, °)

C1—O1	1.131 (3)	C10—Fe4	1.798 (3)
C1—Fe1	1.819 (3)	C11—O11	1.140 (3)
C2—O2	1.136 (3)	C11—Fe4	1.804 (3)
C2—Fe1	1.795 (2)	C12—O12	1.129 (3)
C3—O3	1.142 (3)	C12—Fe4	1.813 (2)

C3—Fe1	1.796 (3)	C13—S1	1.827 (2)
C4—O4	1.145 (3)	C13—S2	1.8300 (19)
C4—Fe2	1.795 (3)	C13—S3	1.830 (2)
C5—O5	1.142 (3)	C13—S4	1.837 (2)
C5—Fe2	1.795 (3)	Fe1—S1	2.2730 (6)
C6—O6	1.130 (3)	Fe1—S2	2.2688 (7)
C6—Fe2	1.824 (3)	Fe1—Fe2	2.4949 (5)
C7—O7	1.127 (3)	Fe2—S1	2.2723 (6)
C7—Fe3	1.818 (3)	Fe2—S2	2.2685 (7)
C8—O8	1.138 (3)	Fe3—S3	2.2676 (6)
C8—Fe3	1.796 (3)	Fe3—S4	2.2680 (6)
C9—O9	1.134 (3)	Fe3—Fe4	2.5007 (5)
C9—Fe3	1.798 (3)	Fe4—S3	2.2712 (7)
C10—O10	1.135 (3)	Fe4—S4	2.2626 (6)
S1...S2	2.6984 (8)	S3...S4	2.6977 (8)
O1—C1—Fe1	178.3 (2)	C6—Fe2—Fe1	154.51 (8)
O2—C2—Fe1	178.8 (2)	S2—Fe2—Fe1	56.649 (18)
O3—C3—Fe1	179.6 (3)	S1—Fe2—Fe1	56.723 (17)
O4—C4—Fe2	179.2 (3)	C8—Fe3—C9	91.66 (12)
O5—C5—Fe2	177.0 (2)	C8—Fe3—C7	97.13 (12)
O6—C6—Fe2	177.6 (2)	C9—Fe3—C7	98.48 (11)
O7—C7—Fe3	176.7 (2)	C8—Fe3—S3	93.97 (9)
O8—C8—Fe3	179.5 (3)	C9—Fe3—S3	155.51 (7)
O9—C9—Fe3	178.5 (2)	C7—Fe3—S3	104.42 (8)
O10—C10—Fe4	179.2 (3)	C8—Fe3—S4	158.63 (9)
O11—C11—Fe4	179.5 (3)	C9—Fe3—S4	93.86 (8)
O12—C12—Fe4	178.0 (2)	C7—Fe3—S4	102.43 (8)
S1—C13—S3	118.10 (10)	S3—Fe3—S4	72.99 (2)
S1—C13—S2	95.10 (10)	C8—Fe3—Fe4	102.35 (9)
S3—C13—S2	117.13 (11)	C9—Fe3—Fe4	98.88 (7)
S1—C13—S4	116.95 (11)	C7—Fe3—Fe4	153.39 (8)
S3—C13—S4	94.73 (9)	S3—Fe3—Fe4	56.635 (18)
S2—C13—S4	116.62 (10)	S4—Fe3—Fe4	56.394 (17)
C2—Fe1—C3	92.20 (12)	C10—Fe4—C11	92.11 (13)
C2—Fe1—C1	97.54 (11)	C10—Fe4—C12	97.98 (11)
C3—Fe1—C1	96.81 (11)	C11—Fe4—C12	97.54 (11)
C2—Fe1—S2	93.44 (9)	C10—Fe4—S4	157.10 (8)
C3—Fe1—S2	158.89 (8)	C11—Fe4—S4	93.60 (9)
C1—Fe1—S2	102.58 (8)	C12—Fe4—S4	103.21 (8)
C2—Fe1—S1	156.74 (8)	C10—Fe4—S3	94.03 (9)
C3—Fe1—S1	94.58 (8)	C11—Fe4—S3	157.73 (8)
C1—Fe1—S1	103.69 (7)	C12—Fe4—S3	102.76 (8)
S2—Fe1—S1	72.90 (2)	S4—Fe4—S3	73.03 (2)
C2—Fe1—Fe2	100.11 (8)	C10—Fe4—Fe3	100.52 (8)
C3—Fe1—Fe2	102.33 (8)	C11—Fe4—Fe3	101.33 (8)
C1—Fe1—Fe2	153.30 (7)	C12—Fe4—Fe3	152.93 (8)

S2—Fe1—Fe2	56.636 (18)	S4—Fe4—Fe3	56.600 (16)
S1—Fe1—Fe2	56.695 (17)	S3—Fe4—Fe3	56.499 (16)
C5—Fe2—C4	91.26 (13)	C13—S1—Fe2	88.02 (6)
C5—Fe2—C6	100.64 (12)	C13—S1—Fe1	87.12 (6)
C4—Fe2—C6	96.91 (11)	Fe2—S1—Fe1	66.582 (18)
C5—Fe2—S2	153.65 (8)	C13—S2—Fe2	88.06 (7)
C4—Fe2—S2	93.88 (9)	C13—S2—Fe1	87.17 (7)
C6—Fe2—S2	104.35 (9)	Fe2—S2—Fe1	66.715 (19)
C5—Fe2—S1	93.23 (8)	C13—S3—Fe3	87.77 (7)
C4—Fe2—S1	157.61 (9)	C13—S3—Fe4	87.52 (7)
C6—Fe2—S1	103.77 (8)	Fe3—S3—Fe4	66.867 (19)
S2—Fe2—S1	72.92 (2)	C13—S4—Fe4	87.60 (7)
C5—Fe2—Fe1	97.00 (8)	C13—S4—Fe3	87.58 (6)
C4—Fe2—Fe1	100.96 (8)	Fe4—S4—Fe3	67.006 (19)
