

Hexyl (*E*)-3-(3,4-dihydroxyphenyl)-acrylate

Jun Wang,^a Shuang-Shuang Gu,^a Jing Li^a and Fu-An Wu^{a,b*}

^aSchool of Biological and Chemical Engineering, Jiangsu University of Science and Technology, Zhenjiang 212018, People's Republic of China, and ^bSericultural Research Institute, Chinese Academy of Agricultural Sciences, Zhenjiang 212018, People's Republic of China
Correspondence e-mail: fuword@163.com


Received 13 October 2011; accepted 30 November 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.061; wR factor = 0.178; data-to-parameter ratio = 15.2.

The title molecule, $C_{15}H_{20}O_4$, has an *E* conformation about its $\text{C}=\text{C}$ bond and is almost planar (r.m.s. deviation of all non-H atoms = 0.04 \AA). The crystal structure features $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For general background to caffeic acid and its derivatives, see: Buzzi *et al.* (2009); Uwai *et al.* (2008). For details of the synthesis, see: Feng *et al.* (2011); Son *et al.* (2011). For related structures, see: Xia *et al.* (2004, 2006). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

| | |
|-----------------------------|--|
| $C_{15}H_{20}O_4$ | $\gamma = 96.84(3)^\circ$ |
| $M_r = 264.31$ | $V = 710.6(2)\text{ \AA}^3$ |
| Triclinic, $P\bar{1}$ | $Z = 2$ |
| $a = 5.2920(11)\text{ \AA}$ | Mo $K\alpha$ radiation |
| $b = 10.689(2)\text{ \AA}$ | $\mu = 0.09\text{ mm}^{-1}$ |
| $c = 12.732(3)\text{ \AA}$ | $T = 293\text{ K}$ |
| $\alpha = 95.45(3)^\circ$ | $0.20 \times 0.10 \times 0.10\text{ mm}$ |
| $\beta = 92.76(3)^\circ$ | |

Data collection

| | |
|---|--|
| Enraf–Nonius CAD-4 diffractometer | 2608 independent reflections |
| Absorption correction: ψ scan (North <i>et al.</i> , 1968) | 1515 reflections with $I > 2\sigma(I)$ |
| $T_{\min} = 0.983$, $T_{\max} = 0.991$ | $R_{\text{int}} = 0.022$ |
| 2912 measured reflections | 3 standard reflections every 200 reflections |
| | intensity decay: 1% |

Refinement

| | |
|---------------------------------|---|
| $R[F^2 > 2\sigma(F^2)] = 0.061$ | 172 parameters |
| $wR(F^2) = 0.178$ | H-atom parameters constrained |
| $S = 1.00$ | $\Delta\rho_{\text{max}} = 0.19\text{ e \AA}^{-3}$ |
| 2608 reflections | $\Delta\rho_{\text{min}} = -0.28\text{ e \AA}^{-3}$ |

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|----------------------------------|--------------|--------------------|-------------|----------------------|
| O1—H1A \cdots O2 ⁱ | 0.85 | 2.07 | 2.857 (2) | 154 |
| O2—H2A \cdots O3 ⁱⁱ | 0.82 | 1.97 | 2.786 (3) | 173 |
| C5—H5A \cdots O3 ⁱⁱ | 0.93 | 2.54 | 3.243 (3) | 133 |

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

This work was sponsored by the Qing Lan Project of Jiangsu Province, the Natural Science Foundation of Jiangsu Province (BK2009213), the College Natural Science Research Project of Jiangsu Province (08KJB530002), the Science and Technology Support Program of Jiangsu Province (BE2010419), the Start Project for Introducing Talent of Jiangsu University of Science and Technology (35211002), the Pre-research for NSFC Project of Jiangsu University of Science and Technology (33201002), and the earmarked fund for Modern Agro-industry Technology Research System (CARS-22).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AA2034).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Buzzi, F. de C., Franzoi, C. L., Antonini, G., Fracasso, M., Filho, V. C., Yunes, R. A. & Niero, R. (2009). *Eur. J. Med. Chem.* **44**, 4596–4602.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Feng, Y., Zhang, A., Li, J. & He, B. (2011). *Bioresource Technol.* **102**, 3607–3609.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Son, S. M., Kimura, H. & Kusakabe, K. (2011). *Bioresource Technol.* **102**, 2130–2132.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Uwai, K., Osanai, Y., Imaizumi, T., Kanno, S., Takeshita, M. & Ishikawa, M. (2008). *Bioorg. Med. Chem.* **16**, 7795–7803.
- Xia, C.-N., Hu, W.-X. & Rao, G.-W. (2004). *Acta Cryst. E* **60**, o913–o914.
- Xia, C.-N., Hu, W.-X. & Zhou, W. (2006). *Acta Cryst. E* **62**, o3900–o3901.

supporting information

Acta Cryst. (2012). E68, o78 [doi:10.1107/S1600536811051671]

Hexyl (*E*)-3-(3,4-dihydroxyphenyl)acrylate

Jun Wang, Shuang-Shuang Gu, Jing Li and Fu-An Wu

S1. Comment

Caffeic acid and its derivatives are widely distributed in medicinal plants and are therefore present in human plasma in a diet dependent concentration. These compounds are known to show a variety of biological effects such as anti-tumor, anti-oxidant, and anti-inflammatory activities (Uwai *et al.*, 2008; Buzzi *et al.*, 2009). In order to investigate their properties better, we synthesize a series of caffeic acid esters. The title compound, hexyl (*E*)-3-(3,4-dihydroxyphenyl)acrylate (I) was obtained earlier (Feng *et al.*, 2011; Son *et al.*, 2011). We report herein the crystal structure of the title compound.

The molecule of (I) has an *E* configuration (Fig. 1). All non-H atoms of (I) are almost coplanar, with a root mean square deviation from the least-squares plane of 0.04 Å°. The bond lengths and angles are within normal ranges (Allen *et al.*, 1987), they are in very good agreement with those found in similar caffeic acid structures (Xia *et al.*, 2004; Xia *et al.*, 2006).


In the crystal structure, intermolecular O—H···O interactions (Table 1) link the molecules into ribbons parallel to the (112) plane (Fig. 2), this may be effective in the stabilization of the structure. On the other hand, the intramolecular O—H···O H-bond also contribute to the stability of the molecular configuration (Table 1).

S2. Experimental

Esterification of caffeic acid with hexyl alcohol was performed in a column (inner diameter = 15 mm, length = 200 mm). Cation exchange resin CD-552 particles (5 g), molecular sieve (5 g) and glass beads of 2 mm in diameter were packed into the middle of the reactor. In a reaction mixture tank, caffeic acid (8.95 g) was mixed with 100 ml of hexyl alcohol. The reaction mixture was supplied to the reaction column at 10.0 ml/h. The reaction continued at 90°C for 24 h. The mixture was evaporated to dryness and followed by the addition of ethanol and extracted with dichloromethane three times. The dichloromethane extract was evaporated to give a solid residue. The residue was recrystallized from ethanol/petroleum ether (1:1) to give the title compound as brown crystals (4.9 g, 54.7%).

S3. Refinement

The H atoms were placed in calculated positions (O—H = 0.82 Å° and C—H = 0.93–0.97 Å°) and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5 U_{\text{eq}}(\text{C}, \text{O})$.

**Figure 1**

The molecular structure of the title molecule, with the atom numbering scheme. Thermal displacement ellipsoids are drawn at 30% probability level.




Figure 2

A partial packing diagram. Hydrogen bonds are shown as dashed lines.

Hexyl (*E*-3-(3,4-dihydroxyphenyl)acrylate*Crystal data*

$C_{15}H_{20}O_4$
 $M_r = 264.31$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 5.2920$ (11) Å
 $b = 10.689$ (2) Å
 $c = 12.732$ (3) Å
 $\alpha = 95.45$ (3)°
 $\beta = 92.76$ (3)°
 $\gamma = 96.84$ (3)°
 $V = 710.6$ (2) Å³

$Z = 2$
 $F(000) = 284$
 $D_x = 1.235$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 25 reflections
 $\theta = 9\text{--}13^\circ$
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
Block, brown
 $0.20 \times 0.10 \times 0.10$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator
 $\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.983$, $T_{\max} = 0.991$
2912 measured reflections

2608 independent reflections
1515 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 25.4^\circ$, $\theta_{\min} = 1.6^\circ$
 $h = 0\text{--}6$
 $k = -12\text{--}12$
 $l = -15\text{--}15$
3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.061$

$wR(F^2) = 0.178$

$S = 1.00$

2608 reflections

172 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.085P)^2 + 0.096P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.19$ e Å⁻³

$\Delta\rho_{\min} = -0.28$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|------------|--------------|--------------|----------------------------------|
| O1 | 0.7392 (4) | 0.09426 (17) | 0.38828 (16) | 0.0649 (6) |
| H1A | 0.6881 | 0.0408 | 0.4304 | 0.078* |
| C1 | 0.5906 (5) | 0.3938 (2) | 0.3065 (2) | 0.0504 (7) |
| H1B | 0.6496 | 0.4508 | 0.2597 | 0.060* |
| O2 | 0.3463 (4) | 0.13532 (16) | 0.51361 (14) | 0.0564 (6) |
| H2A | 0.2394 | 0.1617 | 0.5516 | 0.085* |
| C2 | 0.7038 (5) | 0.2853 (2) | 0.3144 (2) | 0.0528 (8) |
| H2B | 0.8390 | 0.2700 | 0.2730 | 0.063* |
| C3 | 0.6192 (5) | 0.1991 (2) | 0.3832 (2) | 0.0463 (7) |
| O3 | 0.0120 (4) | 0.75695 (18) | 0.36537 (16) | 0.0685 (7) |
| O4 | 0.2768 (4) | 0.81988 (16) | 0.24479 (15) | 0.0571 (6) |
| C4 | 0.4190 (5) | 0.2232 (2) | 0.4456 (2) | 0.0434 (7) |
| C5 | 0.3053 (5) | 0.3312 (2) | 0.43703 (19) | 0.0440 (7) |
| H5A | 0.1696 | 0.3462 | 0.4783 | 0.053* |
| C6 | 0.3883 (5) | 0.4188 (2) | 0.36794 (19) | 0.0420 (6) |
| C7 | 0.2635 (5) | 0.5329 (2) | 0.36356 (19) | 0.0443 (7) |
| H7A | 0.1264 | 0.5394 | 0.4059 | 0.053* |
| C8 | 0.3218 (5) | 0.6278 (2) | 0.3067 (2) | 0.0505 (7) |
| H8A | 0.4555 | 0.6237 | 0.2622 | 0.061* |
| C9 | 0.1864 (5) | 0.7388 (2) | 0.3105 (2) | 0.0466 (7) |
| C10 | 0.1571 (6) | 0.9344 (2) | 0.2397 (2) | 0.0518 (7) |
| H10A | 0.1845 | 0.9868 | 0.3067 | 0.062* |
| H10B | -0.0249 | 0.9137 | 0.2235 | 0.062* |
| C11 | 0.2783 (6) | 1.0026 (2) | 0.1537 (2) | 0.0525 (7) |
| H11A | 0.2586 | 0.9468 | 0.0884 | 0.063* |
| H11B | 0.4594 | 1.0242 | 0.1720 | 0.063* |
| C12 | 0.1614 (5) | 1.1224 (2) | 0.1367 (2) | 0.0506 (7) |
| H12A | -0.0210 | 1.1008 | 0.1222 | 0.061* |
| H12B | 0.1877 | 1.1792 | 0.2015 | 0.061* |
| C13 | 0.2692 (6) | 1.1911 (3) | 0.0477 (2) | 0.0584 (8) |
| H13A | 0.2456 | 1.1339 | -0.0168 | 0.070* |
| H13B | 0.4512 | 1.2140 | 0.0629 | 0.070* |
| C14 | 0.1498 (7) | 1.3098 (3) | 0.0292 (3) | 0.0744 (10) |
| H14A | -0.0338 | 1.2884 | 0.0198 | 0.089* |
| H14B | 0.1860 | 1.3702 | 0.0915 | 0.089* |
| C15 | 0.2443 (8) | 1.3718 (3) | -0.0660 (3) | 0.0952 (13) |
| H15A | 0.1629 | 1.4464 | -0.0732 | 0.143* |
| H15B | 0.2044 | 1.3136 | -0.1284 | 0.143* |
| H15C | 0.4256 | 1.3947 | -0.0569 | 0.143* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|-------------|-------------|-------------|-------------|-------------|-------------|
| O1 | 0.0696 (14) | 0.0486 (12) | 0.0879 (15) | 0.0312 (10) | 0.0301 (12) | 0.0239 (10) |
| C1 | 0.0538 (17) | 0.0444 (15) | 0.0573 (17) | 0.0095 (13) | 0.0182 (14) | 0.0156 (13) |

| | | | | | | |
|-----|-------------|-------------|-------------|-------------|-------------|-------------|
| O2 | 0.0697 (13) | 0.0446 (11) | 0.0651 (12) | 0.0252 (10) | 0.0277 (10) | 0.0239 (9) |
| C2 | 0.0500 (17) | 0.0478 (16) | 0.0664 (19) | 0.0191 (14) | 0.0213 (15) | 0.0108 (14) |
| C3 | 0.0475 (16) | 0.0373 (14) | 0.0569 (17) | 0.0139 (12) | 0.0091 (13) | 0.0056 (12) |
| O3 | 0.0834 (16) | 0.0585 (13) | 0.0778 (14) | 0.0343 (11) | 0.0419 (12) | 0.0294 (11) |
| O4 | 0.0708 (13) | 0.0429 (11) | 0.0678 (13) | 0.0232 (10) | 0.0280 (11) | 0.0254 (10) |
| C4 | 0.0489 (16) | 0.0362 (14) | 0.0478 (15) | 0.0100 (12) | 0.0100 (13) | 0.0085 (12) |
| C5 | 0.0484 (16) | 0.0406 (15) | 0.0470 (15) | 0.0153 (12) | 0.0148 (13) | 0.0077 (12) |
| C6 | 0.0485 (16) | 0.0359 (14) | 0.0441 (14) | 0.0104 (12) | 0.0083 (12) | 0.0084 (11) |
| C7 | 0.0486 (17) | 0.0403 (15) | 0.0477 (16) | 0.0126 (13) | 0.0137 (13) | 0.0083 (12) |
| C8 | 0.0581 (18) | 0.0464 (16) | 0.0533 (17) | 0.0174 (14) | 0.0213 (14) | 0.0150 (13) |
| C9 | 0.0539 (17) | 0.0393 (15) | 0.0509 (16) | 0.0126 (13) | 0.0110 (14) | 0.0134 (12) |
| C10 | 0.0632 (19) | 0.0366 (14) | 0.0618 (18) | 0.0199 (13) | 0.0177 (15) | 0.0136 (13) |
| C11 | 0.0647 (19) | 0.0437 (15) | 0.0538 (16) | 0.0150 (14) | 0.0165 (14) | 0.0133 (13) |
| C12 | 0.0589 (18) | 0.0412 (15) | 0.0556 (17) | 0.0119 (13) | 0.0149 (14) | 0.0124 (13) |
| C13 | 0.072 (2) | 0.0495 (17) | 0.0578 (18) | 0.0131 (15) | 0.0152 (16) | 0.0156 (14) |
| C14 | 0.098 (3) | 0.0557 (19) | 0.077 (2) | 0.0170 (18) | 0.018 (2) | 0.0281 (17) |
| C15 | 0.134 (4) | 0.072 (2) | 0.083 (3) | 0.002 (2) | 0.007 (2) | 0.037 (2) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|-----------|-----------|---------------|-----------|
| O1—C3 | 1.357 (3) | C8—H8A | 0.9300 |
| O1—H1A | 0.8500 | C10—C11 | 1.499 (3) |
| C1—C2 | 1.377 (4) | C10—H10A | 0.9700 |
| C1—C6 | 1.392 (3) | C10—H10B | 0.9700 |
| C1—H1B | 0.9300 | C11—C12 | 1.516 (3) |
| O2—C4 | 1.371 (3) | C11—H11A | 0.9700 |
| O2—H2A | 0.8200 | C11—H11B | 0.9700 |
| C2—C3 | 1.382 (3) | C12—C13 | 1.507 (3) |
| C2—H2B | 0.9300 | C12—H12A | 0.9700 |
| C3—C4 | 1.389 (3) | C12—H12B | 0.9700 |
| O3—C9 | 1.207 (3) | C13—C14 | 1.516 (4) |
| O4—C9 | 1.326 (3) | C13—H13A | 0.9700 |
| O4—C10 | 1.449 (3) | C13—H13B | 0.9700 |
| C4—C5 | 1.375 (3) | C14—C15 | 1.514 (4) |
| C5—C6 | 1.392 (3) | C14—H14A | 0.9700 |
| C5—H5A | 0.9300 | C14—H14B | 0.9700 |
| C6—C7 | 1.459 (3) | C15—H15A | 0.9600 |
| C7—C8 | 1.317 (3) | C15—H15B | 0.9600 |
| C7—H7A | 0.9300 | C15—H15C | 0.9600 |
| C8—C9 | 1.455 (3) | | |
| C3—O1—H1A | 118.8 | O4—C10—H10B | 110.4 |
| C2—C1—C6 | 120.5 (2) | C11—C10—H10B | 110.4 |
| C2—C1—H1B | 119.8 | H10A—C10—H10B | 108.6 |
| C6—C1—H1B | 119.8 | C10—C11—C12 | 112.0 (2) |
| C4—O2—H2A | 109.5 | C10—C11—H11A | 109.2 |
| C1—C2—C3 | 120.9 (2) | C12—C11—H11A | 109.2 |
| C1—C2—H2B | 119.6 | C10—C11—H11B | 109.2 |

| | | | |
|--------------|------------|-----------------|------------|
| C3—C2—H2B | 119.6 | C12—C11—H11B | 109.2 |
| O1—C3—C2 | 118.3 (2) | H11A—C11—H11B | 107.9 |
| O1—C3—C4 | 122.3 (2) | C13—C12—C11 | 113.9 (2) |
| C2—C3—C4 | 119.3 (2) | C13—C12—H12A | 108.8 |
| C9—O4—C10 | 117.5 (2) | C11—C12—H12A | 108.8 |
| O2—C4—C5 | 123.6 (2) | C13—C12—H12B | 108.8 |
| O2—C4—C3 | 116.7 (2) | C11—C12—H12B | 108.8 |
| C5—C4—C3 | 119.7 (2) | H12A—C12—H12B | 107.7 |
| C4—C5—C6 | 121.6 (2) | C12—C13—C14 | 114.1 (3) |
| C4—C5—H5A | 119.2 | C12—C13—H13A | 108.7 |
| C6—C5—H5A | 119.2 | C14—C13—H13A | 108.7 |
| C5—C6—C1 | 118.1 (2) | C12—C13—H13B | 108.7 |
| C5—C6—C7 | 119.2 (2) | C14—C13—H13B | 108.7 |
| C1—C6—C7 | 122.7 (2) | H13A—C13—H13B | 107.6 |
| C8—C7—C6 | 127.7 (2) | C15—C14—C13 | 113.5 (3) |
| C8—C7—H7A | 116.1 | C15—C14—H14A | 108.9 |
| C6—C7—H7A | 116.1 | C13—C14—H14A | 108.9 |
| C7—C8—C9 | 122.9 (2) | C15—C14—H14B | 108.9 |
| C7—C8—H8A | 118.5 | C13—C14—H14B | 108.9 |
| C9—C8—H8A | 118.5 | H14A—C14—H14B | 107.7 |
| O3—C9—O4 | 122.9 (2) | C14—C15—H15A | 109.5 |
| O3—C9—C8 | 125.4 (2) | C14—C15—H15B | 109.5 |
| O4—C9—C8 | 111.7 (2) | H15A—C15—H15B | 109.5 |
| O4—C10—C11 | 106.6 (2) | C14—C15—H15C | 109.5 |
| O4—C10—H10A | 110.4 | H15A—C15—H15C | 109.5 |
| C11—C10—H10A | 110.4 | H15B—C15—H15C | 109.5 |
| | | | |
| C6—C1—C2—C3 | 0.1 (4) | C5—C6—C7—C8 | 177.1 (3) |
| C1—C2—C3—O1 | -179.5 (3) | C1—C6—C7—C8 | -2.1 (5) |
| C1—C2—C3—C4 | -0.7 (4) | C6—C7—C8—C9 | -178.6 (3) |
| O1—C3—C4—O2 | -0.1 (4) | C10—O4—C9—O3 | -0.3 (4) |
| C2—C3—C4—O2 | -178.9 (2) | C10—O4—C9—C8 | 179.3 (2) |
| O1—C3—C4—C5 | 179.9 (3) | C7—C8—C9—O3 | 0.9 (5) |
| C2—C3—C4—C5 | 1.1 (4) | C7—C8—C9—O4 | -178.8 (3) |
| O2—C4—C5—C6 | 179.0 (2) | C9—O4—C10—C11 | -175.1 (2) |
| C3—C4—C5—C6 | -0.9 (4) | O4—C10—C11—C12 | 177.5 (2) |
| C4—C5—C6—C1 | 0.4 (4) | C10—C11—C12—C13 | -177.4 (2) |
| C4—C5—C6—C7 | -178.9 (2) | C11—C12—C13—C14 | 179.0 (3) |
| C2—C1—C6—C5 | 0.0 (4) | C12—C13—C14—C15 | -175.2 (3) |
| C2—C1—C6—C7 | 179.3 (3) | | |

Hydrogen-bond geometry (\AA , $^\circ$)

| $D\text{—H}\cdots A$ | $D\text{—H}$ | $H\cdots A$ | $D\cdots A$ | $D\text{—H}\cdots A$ |
|---------------------------------------|--------------|-------------|-------------|----------------------|
| O1—H1A ⁱ —O2 | 0.85 | 2.41 | 2.733 (2) | 103 |
| O1—H1A ⁱ —O2 ^j | 0.85 | 2.07 | 2.857 (2) | 154 |
| O2—H2A ⁱ —O3 ⁱⁱ | 0.82 | 1.97 | 2.786 (3) | 173 |

| | | | | |
|---------------------------|------|------|-----------|-----|
| C5—H5A···O3 ⁱⁱ | 0.93 | 2.54 | 3.243 (3) | 133 |
| C7—H7A···O3 | 0.93 | 2.56 | 2.874 (3) | 100 |

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, -y+1, -z+1$.