

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Nonaaquaprasedymium triiodide–thiourea (1/2)

Taisia A. Antonenko,^a Lyudmila Yu. Alikberova^a and Dmitry V. Albov^{b*}

^aDepartment of Inorganic Chemistry, M. V. Lomonosov Moscow State University of Fine Chemical Technologies, 86 Vernadskogo Av., Moscow 119571, Russian Federation, and ^bChemistry Department, Moscow State University, Leninskiye Gory, Moscow 119992, Russian Federation

Correspondence e-mail: dmitryalbov@yandex.ru


Received 21 November 2011; accepted 19 December 2011

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{N}-\text{C}) = 0.016$ Å; H-atom completeness 31%; R factor = 0.040; wR factor = 0.100; data-to-parameter ratio = 23.6.

The title compound, $[\text{Pr}(\text{H}_2\text{O})_9]\text{I}_3 \cdot 2\text{CS}(\text{NH}_2)_2$, an adduct of nona-aquaprasedymium triiodide with two thiourea molecules, is composed from $[\text{Pr}(\text{H}_2\text{O})_9]^{3+}$ cations (polyhedron: monocapped tetragonal antiprism), noncoordinated thiourea molecules and iodide anions. The components are evidently connected by hydrogen bonds but in the presence of heavy atoms water H atoms have not been located. The complex cation and one of the two independent iodide anions are located on a twofold axis.

Related literature

For related compounds, see: Romanenko *et al.* (1980, 1981a,b, 1985, 1986); Antonenko *et al.* (2011). For applications of similar complexes, see: Suponitsky *et al.* (1988). For titration methods, see: Patrovsky (1959); Kolthoff & Belcher (1957).



Experimental

Crystal data

 $[\text{Pr}(\text{H}_2\text{O})_9]\text{I}_3 \cdot 2\text{CH}_4\text{N}_2\text{S}$ $M_r = 836.00$

Monoclinic, $C2/c$
 $a = 24.934$ (18) Å
 $b = 8.439$ (3) Å
 $c = 14.143$ (8) Å
 $\beta = 124.68$ (5)°
 $V = 2447$ (3) Å³

$Z = 4$
 Ag $K\alpha$ radiation
 $\lambda = 0.56085$ Å
 $\mu = 3.16$ mm⁻¹
 $T = 295$ K
 $0.20 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\text{min}} = 0.405$, $T_{\text{max}} = 0.592$

2309 measured reflections
 2309 independent reflections
 1827 reflections with $I > 2\sigma(I)$
 1 standard reflection every 120 min
 intensity decay: 2%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.100$
 $S = 0.99$
 2309 reflections

98 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.88$ e Å⁻³

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2319).

References

- Antonenko, T. A., Alikberova, L. Yu., Albov, D. V. & Rukk, N. S. (2011). *Russ. J. Coord. Chem.* **37**, 785–790.
- Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Kolthoff, I. M. & Belcher, R. N. Y. (1957). *Volumetric Analysis*. Vol. 3, pp. 387–389. New York: Interscience.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Patrovsky, V. (1959). *Coll. Czech. Chem. Commun.* **24**, 3305–3308.
- Romanenko, G. V., Podberezskaya, N. V. & Bakakin, V. V. (1981a). *J. Struct. Chem.* **22**, 299–301.
- Romanenko, G. V., Podberezskaya, N. V. & Bakakin, V. V. (1986). *J. Struct. Chem.* **27**, 321–324.
- Romanenko, G. V., Podberezskaya, N. V., Bakakin, V. V. & Sakharova, Yu. G. (1980). *J. Struct. Chem.* **21**, 348–351.
- Romanenko, G. V., Podberezskaya, N. V., Bakakin, V. V. & Sakharova, Yu. G. (1981b). *J. Struct. Chem.* **22**, 740–743.
- Romanenko, G. V., Podberezskaya, N. V., Bakakin, V. V. & Sakharova, Yu. G. (1985). *J. Struct. Chem.* **26**, 743–748.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Suponitsky, Yu. L., Kuz'micheva, G. M. & Eliseev, A. A. (1988). *Russ. Chem. Rev.* **57**(3), 209–220.

supporting information

Acta Cryst. (2012). E68, m110 [doi:10.1107/S1600536811054663]

Nonaaquapraseodymium triiodide–thiourea (1/2)

Taisia A. Antonenko, Lyudmila Yu. Alikberova and Dmitry V. Albov

S1. Comment

The structure investigation of the interaction products of metal salts with thiourea $\text{CS}(\text{NH}_2)_2$ is very promising, because it allows to predict the possible ways of thermal decomposition of these compounds yielding oxide, sulfide and oxosulfide derivatives (Suponitsky *et al.*, 1988). The lanthanide derivatives are very promising objects of research in this regard, since the corresponding sulfides and oxosulfides are used as the activators of materials with the luminescent properties. To the present time the structure of only some thiourea derivatives of lanthanide salts has been studied in details. Systematic investigation of the previously synthesized thiourea derivatives allowed us to conclude that there are two isostructural series of lanthanide acetates (La–Pr and Nd–Lu) (Romanenko *et al.*, 1980; Romanenko *et al.*, 1981a; Romanenko *et al.*, 1986), and three isostructural series of lanthanide propionates (La–Pr, Nd–Tm, Yb–Lu) (Romanenko *et al.*, 1981b; Romanenko *et al.*, 1985). It was established the existence of complex cations in the structures, involving coordinated water molecules as well as bidentate and bridging acetate or propionate ions. It was noted that thiourea is not included into the internal sphere of complexes. The information about the synthesis and structure of thiourea derivatives of lanthanide halides is much smaller. We have obtained the compounds of thiourea with LnI_3 ($\text{Ln} = \text{Eu}, \text{Ho}, \text{Er}$) at room temperature (Antonenko *et al.*, 2011). X–ray data have been demonstrated that in the solid state these compounds are composed from $[\text{Ln}(\text{H}_2\text{O})_9]^{3+}$ cations (polyhedron: monocapped tetragonal antiprism), non–coordinated thiourea molecules and iodide–ions.

Herein we report the structure of thiourea adduct of nonaaquapraseodymium triiodide **I** (Fig. 1). In the solid state **I** is composed from $[\text{Pr}(\text{H}_2\text{O})_9]^{3+}$ cations (polyhedron: monocapped tetragonal antiprism), thiourea molecules and iodide anions. All mentioned species are evidently connected with H–bonds but in the presence of heavy atoms water H atoms have not been located and thus can not be discussed. The complex cation and one of the two independent iodide anions are located on a twofold axis. There is no coordination of thiourea by the lanthanide atom as through the atom S, and through the atom N as well as in the cases of compounds which have been obtained previously (Antonenko *et al.*, 2011). A packing diagram is shown in Fig. 2.

S2. Experimental

The synthesis of title compound was carried out at room temperature by mixing $\text{PrI}_3 \cdot 9\text{H}_2\text{O}$ and $\text{CS}(\text{NH}_2)_2$ at a molar ratio 1:1.7. Few drops of water were added to the reaction mixture to the formation of clear solution. After 30 days the light green crystals were identified from it. These crystals are hygroscopic; they are decomposed by water with the release of the initial thiourea. The crystals of title compound suitable for X–ray analysis were dried over the alkali in the desiccator. By complexometric titration with 0.1 M *Edta* and reverse iodometric titration with 0.1 M $\text{Na}_2\text{S}_2\text{O}_3$ (Patrovsky, 1959; Kolthoff & Belcher, 1957) we established that the molar ratio of nonaaquapraseodymium triiodide and thiourea in this compound is 1:2.

S3. Refinement

In the presence of heavy atoms water H atoms could not be located. The hydrogen atoms bound to N atoms were placed in calculated positions with $N-H = 0.86\text{\AA}$ and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$.




Figure 1

Crystal structure of **I** with the atom numbering scheme. Displacement ellipsoids are shown at the 50% probability level. Only independent iodide anions are shown. Symmetry code: (i) $1-x, y, 1/2-z$.




Figure 2

Crystal packing of **I**, a view along the a axis.

Nonaquapraseodymium triiodide–thiourea (1/2)

Crystal data

$[\text{Pr}(\text{H}_2\text{O})_9]\text{I}_3 \cdot 2\text{CH}_4\text{N}_2\text{S}$
 $M_r = 836.00$
 Monoclinic, $C2/c$
 Hall symbol: $-C 2yc$

$a = 24.934 (18) \text{\AA}$
 $b = 8.439 (3) \text{\AA}$
 $c = 14.143 (8) \text{\AA}$
 $\beta = 124.68 (5)^\circ$

$V = 2447 (3) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 1552$
 $D_x = 2.269 \text{ Mg m}^{-3}$
 Ag $K\alpha$ radiation, $\lambda = 0.56085 \text{ \AA}$
 Cell parameters from 25 reflections

$\theta = 12\text{--}13^\circ$
 $\mu = 3.16 \text{ mm}^{-1}$
 $T = 295 \text{ K}$
 Prism, light green
 $0.20 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Enraf–Nonius CAD-4
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 non-profiled ω scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.405$, $T_{\max} = 0.592$
 2309 measured reflections

2309 independent reflections
 1827 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.000$
 $\theta_{\max} = 20.0^\circ$, $\theta_{\min} = 1.6^\circ$
 $h = -30 \rightarrow 24$
 $k = 0 \rightarrow 10$
 $l = 0 \rightarrow 17$
 1 standard reflections every 120 min
 intensity decay: 2%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.100$
 $S = 0.99$
 2309 reflections
 98 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0486P)^2 + 1.3974P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.24 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.88 \text{ e \AA}^{-3}$

Special details

Experimental. North *et al.*, 1968. Number of ψ -scan sets used was 5. Theta correction was applied. Averaged transmission function was used. Fourier smoothing - Window value 3.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pr1	0.5000	0.80167 (6)	0.2500	0.03210 (15)
O1	0.5000	1.1003 (7)	0.2500	0.0419 (15)
O2	0.5919 (2)	0.6295 (6)	0.2854 (4)	0.0547 (13)
O3	0.5369 (3)	0.6201 (7)	0.4163 (4)	0.0619 (14)
O4	0.5386 (3)	0.9090 (6)	0.1328 (5)	0.0604 (14)
O5	0.6059 (2)	0.9151 (7)	0.4141 (5)	0.0705 (17)
I1	0.2500	0.7500	0.0000	0.0930 (4)
I2	0.36775 (2)	0.71928 (6)	0.45157 (4)	0.04941 (17)

S1	0.57937 (10)	0.7171 (2)	0.66733 (17)	0.0466 (4)
C1	0.6621 (4)	0.7453 (8)	0.7467 (7)	0.0489 (18)
N11	0.6976 (4)	0.6708 (17)	0.7270 (10)	0.160 (6)
H11A	0.7387	0.6912	0.7657	0.192*
H11B	0.6812	0.5988	0.6747	0.192*
N12	0.6906 (4)	0.8508 (17)	0.8263 (9)	0.170 (6)
H12A	0.7319	0.8661	0.8622	0.204*
H12B	0.6683	0.9061	0.8437	0.204*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pr1	0.0272 (2)	0.0302 (2)	0.0343 (3)	0.000	0.0148 (2)	0.000
O1	0.051 (4)	0.031 (3)	0.052 (4)	0.000	0.034 (3)	0.000
O2	0.050 (3)	0.051 (3)	0.058 (3)	0.013 (2)	0.028 (3)	-0.010 (3)
O3	0.075 (4)	0.058 (3)	0.040 (3)	0.021 (3)	0.026 (3)	0.019 (3)
O4	0.081 (4)	0.046 (3)	0.079 (4)	-0.017 (3)	0.060 (3)	-0.010 (3)
O5	0.039 (3)	0.058 (3)	0.075 (4)	0.006 (3)	0.009 (3)	-0.031 (3)
I1	0.0320 (4)	0.1182 (9)	0.0991 (8)	-0.0120 (4)	0.0196 (5)	0.0195 (6)
I2	0.0496 (3)	0.0431 (3)	0.0541 (3)	0.0015 (2)	0.0286 (3)	-0.0034 (2)
S1	0.0478 (10)	0.0444 (10)	0.0513 (10)	0.0068 (8)	0.0304 (9)	0.0000 (8)
C1	0.060 (5)	0.048 (4)	0.052 (4)	0.002 (4)	0.040 (4)	-0.004 (3)
N11	0.053 (5)	0.244 (15)	0.160 (10)	-0.019 (8)	0.047 (6)	-0.129 (11)
N12	0.075 (6)	0.240 (15)	0.156 (10)	-0.030 (8)	0.042 (7)	-0.151 (11)

Geometric parameters (Å, °)

Pr1—O3 ⁱ	2.503 (5)	Pr1—O1	2.520 (6)
Pr1—O3	2.503 (5)	S1—C1	1.713 (10)
Pr1—O5	2.511 (5)	C1—N11	1.240 (12)
Pr1—O5 ⁱ	2.511 (5)	C1—N12	1.287 (11)
Pr1—O2 ⁱ	2.512 (5)	N11—H11A	0.8600
Pr1—O2	2.512 (5)	N11—H11B	0.8600
Pr1—O4	2.512 (5)	N12—H12A	0.8600
Pr1—O4 ⁱ	2.512 (5)	N12—H12B	0.8600
O3 ⁱ —Pr1—O3	104.5 (3)	O5—Pr1—O4 ⁱ	81.32 (19)
O3 ⁱ —Pr1—O5	137.14 (18)	O5 ⁱ —Pr1—O4 ⁱ	82.9 (2)
O3—Pr1—O5	74.6 (2)	O2 ⁱ —Pr1—O4 ⁱ	72.02 (19)
O3 ⁱ —Pr1—O5 ⁱ	74.6 (2)	O2—Pr1—O4 ⁱ	136.54 (17)
O3—Pr1—O5 ⁱ	137.14 (18)	O4—Pr1—O4 ⁱ	137.7 (2)
O5—Pr1—O5 ⁱ	135.2 (3)	O3 ⁱ —Pr1—O1	127.76 (13)
O3 ⁱ —Pr1—O2 ⁱ	69.67 (18)	O3—Pr1—O1	127.76 (13)
O3—Pr1—O2 ⁱ	68.84 (18)	O5—Pr1—O1	67.59 (13)
O5—Pr1—O2 ⁱ	139.9 (2)	O5 ⁱ —Pr1—O1	67.59 (13)
O5 ⁱ —Pr1—O2 ⁱ	71.06 (17)	O2 ⁱ —Pr1—O1	125.34 (13)
O3 ⁱ —Pr1—O2	68.84 (18)	O2—Pr1—O1	125.34 (13)
O3—Pr1—O2	69.67 (18)	O4—Pr1—O1	68.86 (12)

O5—Pr1—O2	71.06 (17)	O4 ⁱ —Pr1—O1	68.86 (12)
O5 ⁱ —Pr1—O2	139.9 (2)	N11—C1—N12	115.9 (9)
O2 ⁱ —Pr1—O2	109.3 (3)	N11—C1—S1	122.0 (7)
O3 ⁱ —Pr1—O4	71.02 (17)	N12—C1—S1	122.0 (7)
O3—Pr1—O4	140.07 (18)	C1—N11—H11A	120.0
O5—Pr1—O4	82.9 (2)	C1—N11—H11B	120.0
O5 ⁱ —Pr1—O4	81.32 (19)	H11A—N11—H11B	120.0
O2 ⁱ —Pr1—O4	136.54 (17)	C1—N12—H12A	120.0
O2—Pr1—O4	72.02 (19)	C1—N12—H12B	120.0
O3 ⁱ —Pr1—O4 ⁱ	140.07 (18)	H12A—N12—H12B	120.0
O3—Pr1—O4 ⁱ	71.02 (17)		

Symmetry code: (i) $-x+1, y, -z+1/2$.