

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(3-Chloro-4-methylphenyl)maleamic acid

 U. Chaithanya,^a Sabine Foro^b and B. Thimme Gowda^{a*}
^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 21 February 2012; accepted 21 February 2012

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.105; data-to-parameter ratio = 14.1.

In the title compound, $\text{C}_{11}\text{H}_{10}\text{ClNO}_3$, the dihedral angle between the benzene ring and the amide group is $6.6(10)^\circ$ and an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond occurs. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, generating $C(7)$ zigzag chains.

Related literature

For our studies on the effects of substituents on the structures and other aspects of N -(aryl)-amides, see: Gowda *et al.* (2000, 2003, 2007); Chaithanya *et al.* (2012). For N -chloroarylamides, see: Jyothi & Gowda (2004). For N -bromoarylsulfonamides, see: Usha & Gowda (2006).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{10}\text{ClNO}_3$
 $M_r = 239.65$
 Monoclinic, $P2_1/c$
 $a = 9.005(1)$ Å
 $b = 13.491(2)$ Å
 $c = 8.757(1)$ Å
 $\beta = 97.91(1)^\circ$

$V = 1053.7(2)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.35$ mm⁻¹
 $T = 293$ K
 $0.48 \times 0.40 \times 0.34$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)
 $T_{\min} = 0.849$, $T_{\max} = 0.890$
 4069 measured reflections
 2147 independent reflections
 1798 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.105$
 $S = 1.04$
 2147 reflections
 152 parameters
 2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.87 (2)	2.11 (2)	2.9546 (19)	164 (2)
$\text{O3}-\text{H3O}\cdots\text{O1}$	0.87 (2)	1.62 (2)	2.4885 (17)	173 (2)

 Symmetry code: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the University Grants Commission, Government of India, New Delhi, for a UGC-BSR one-time Grant to Faculty.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2391).

References

- Chaithanya, U., Foro, S. & Gowda, B. T. (2012). *Acta Cryst.* **E68**, o785.
 Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o1975–o1976.
 Gowda, B. T., Kumar, B. H. A. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 721–728.
 Gowda, B. T., Usha, K. M. & Jayalakshmi, K. L. (2003). *Z. Naturforsch. Teil A*, **58**, 801–806.
 Jyothi, K. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **59**, 64–68.
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Usha, K. M. & Gowda, B. T. (2006). *J. Chem. Sci.* **118**, 351–359.

supporting information

Acta Cryst. (2012). E68, o889 [doi:10.1107/S1600536812007842]

N*-(3-Chloro-4-methylphenyl)maleamic acid*U. Chaithanya, Sabine Foro and B. Thimme Gowda****S1. Comment**

As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Gowda *et al.*, 2000, 2003, 2007; Chaithanya *et al.*, 2012), *N*-chloroarylsulfonamides (Jyothi & Gowda, 2004) and *N*-bromoaryl-sulfonamides (Usha & Gowda, 2006), in the present work, the crystal structure of *N*-(3-chloro-4-methylphenyl)maleamic acid has been determined (Fig. 1). The conformations of the N—H and the C=O bonds in the amide segment are *anti* to each other. The conformation of the N—H bond is also *anti* to the *meta*-chloro atom. Further, the conformation of the amide C=O is *anti* to the H atom on the adjacent —CH group, while the carboxyl C=O of the acid segment is *syn* to the adjacent —CH group. Furthermore, the C=O and O—H bond of the acid group are in relatively rare *anti* position to each other, due to the donation of hydrogen bond to the amide by the carboxyl group, in contrast to the more general *syn* conformation observed in *N*-(3-chloro-4-methylphenyl)-succinamic acid (I) (Chaithanya *et al.*, 2012).

The dihedral angle between the phenyl ring and the amide group in the title compound is 6.55 (99)°, compared to the values of 40.58 (22)° and 44.93 (27)° in the two derivatives of (I).

In the structure, the pairs of O—H...O and N—H...O intermolecular hydrogen bonds pack the molecules into chains (Table 1, Fig.2).


S2. Experimental

Maleic anhydride (0.025 mol) in toluene (25 mL) was treated dropwise with 3-chloro-4-methylaniline (0.025 mol) also in toluene (20 mL) with constant stirring. The resulting mixture was stirred for about 30 min and set aside for an additional 30 min at room temperature for the completion of reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 3-chloro-4-methylaniline. The resultant solid *N*-(3-chloro-4-methylphenyl)-maleamic acid was filtered under suction and washed thoroughly with water to remove the unreacted maleic anhydride and maleic acid. It was recrystallised to constant melting point from ethanol. The purity of the compound was checked and characterized by its infrared spectra.

Prism like pale yellow single crystals of the title compound used in X-ray diffraction studies were grown in an ethanol solution by slow evaporation of the solvent (0.5 g in about 30 mL of ethanol) at room temperature.

S3. Refinement

The H atoms of the NH group and the OH group were located in a difference map and later restrained to the distance N—H = 0.86 (2) Å and O—H = 0.82 (2) Å, respectively. The other H atoms were positioned with idealized geometry using a riding model with the aromatic C—H = 0.93 Å and methylene C—H = 0.97 Å. All H atoms were refined with isotropic displacement parameters set at 1.2 U_{eq} (C-aromatic, N) and 1.5 U_{eq} (C-methyl).

**Figure 1**

Molecular structure of the title compound showing the atom labelling scheme with displacement ellipsoids drawn at the 50% probability level. Intramolecular hydrogen bond is shown.




Figure 2

Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

***N*-(3-Chloro-4-methylphenyl)maleamic acid**

Crystal data

$C_{11}H_{10}ClNO_3$

$M_r = 239.65$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.005\ (1)\ \text{\AA}$

$b = 13.491\ (2)\ \text{\AA}$

$c = 8.757\ (1)\ \text{\AA}$

$\beta = 97.91\ (1)^\circ$

$V = 1053.7\ (2)\ \text{\AA}^3$

$Z = 4$

$F(000) = 496$

$D_x = 1.511\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1538 reflections

$\theta = 2.8\text{--}27.9^\circ$

$\mu = 0.35\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, yellow

$0.48 \times 0.40 \times 0.34\ \text{mm}$

Data collection

Oxford Diffraction Xcalibur
diffractometer with a Sapphire CCD detector
Radiation source: fine-focus sealed tube
Graphite monochromator
Rotation method data acquisition using ω and
phi scans
Absorption correction: multi-scan
(*CrysAlis RED*; Oxford Diffraction, 2009)
 $T_{\min} = 0.849$, $T_{\max} = 0.890$

4069 measured reflections
2147 independent reflections
1798 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$
 $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.8^\circ$
 $h = -11 \rightarrow 8$
 $k = -16 \rightarrow 14$
 $l = -7 \rightarrow 10$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.105$
 $S = 1.04$
2147 reflections
152 parameters
2 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0582P)^2 + 0.2934P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. *CrysAlis RED* (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.66874 (17)	0.04762 (11)	0.39541 (18)	0.0309 (3)
C2	0.59803 (18)	0.12392 (11)	0.30833 (19)	0.0359 (4)
H2	0.6327	0.1887	0.3217	0.043*
C3	0.47494 (18)	0.10217 (11)	0.20120 (19)	0.0355 (4)
C4	0.41627 (17)	0.00764 (12)	0.17621 (18)	0.0332 (4)
C5	0.49161 (19)	-0.06669 (12)	0.2641 (2)	0.0398 (4)
H5	0.4571	-0.1315	0.2503	0.048*
C6	0.61566 (18)	-0.04879 (12)	0.3714 (2)	0.0381 (4)
H6	0.6637	-0.1009	0.4275	0.046*
C7	0.85614 (17)	0.14437 (12)	0.56761 (18)	0.0324 (3)
C8	0.98127 (18)	0.13165 (12)	0.69471 (19)	0.0351 (4)
H8	1.0081	0.0665	0.7197	0.042*
C9	1.06040 (18)	0.20092 (13)	0.77789 (19)	0.0378 (4)

H9	1.1357	0.1758	0.8509	0.045*
C10	1.05105 (18)	0.31142 (12)	0.77605 (19)	0.0366 (4)
C11	0.27987 (18)	-0.01421 (14)	0.0625 (2)	0.0434 (4)
H11A	0.2960	0.0081	-0.0380	0.052*
H11B	0.1949	0.0196	0.0930	0.052*
H11C	0.2614	-0.0843	0.0598	0.052*
N1	0.79351 (15)	0.06004 (10)	0.51139 (16)	0.0338 (3)
H1N	0.827 (2)	0.0067 (12)	0.560 (2)	0.041*
O1	0.81351 (14)	0.22693 (9)	0.51661 (15)	0.0483 (3)
O2	1.13005 (15)	0.35781 (10)	0.87433 (15)	0.0507 (4)
O3	0.95988 (16)	0.35732 (9)	0.67213 (17)	0.0537 (4)
H3O	0.911 (2)	0.3136 (16)	0.611 (2)	0.064*
C11	0.38990 (6)	0.19890 (3)	0.09113 (6)	0.0620 (2)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0306 (7)	0.0240 (7)	0.0350 (8)	-0.0015 (6)	-0.0065 (6)	-0.0020 (6)
C2	0.0396 (8)	0.0203 (7)	0.0427 (9)	-0.0025 (6)	-0.0119 (7)	-0.0008 (6)
C3	0.0381 (8)	0.0234 (7)	0.0407 (9)	0.0018 (6)	-0.0099 (7)	-0.0006 (7)
C4	0.0324 (8)	0.0264 (8)	0.0375 (8)	-0.0021 (6)	-0.0064 (6)	-0.0040 (6)
C5	0.0425 (9)	0.0218 (8)	0.0503 (10)	-0.0058 (6)	-0.0112 (7)	-0.0008 (7)
C6	0.0421 (9)	0.0219 (8)	0.0453 (9)	-0.0008 (6)	-0.0118 (7)	0.0024 (7)
C7	0.0323 (7)	0.0259 (7)	0.0359 (8)	-0.0011 (6)	-0.0064 (6)	-0.0018 (6)
C8	0.0372 (8)	0.0264 (8)	0.0378 (9)	0.0012 (6)	-0.0083 (7)	0.0009 (6)
C9	0.0374 (8)	0.0341 (9)	0.0368 (8)	0.0010 (7)	-0.0125 (7)	0.0008 (7)
C10	0.0370 (8)	0.0322 (8)	0.0376 (8)	-0.0037 (7)	-0.0051 (7)	-0.0042 (7)
C11	0.0395 (9)	0.0356 (9)	0.0497 (10)	-0.0037 (7)	-0.0135 (8)	-0.0050 (8)
N1	0.0354 (7)	0.0229 (6)	0.0385 (7)	-0.0005 (5)	-0.0115 (6)	0.0022 (5)
O1	0.0517 (7)	0.0246 (6)	0.0584 (8)	-0.0014 (5)	-0.0286 (6)	0.0011 (5)
O2	0.0559 (8)	0.0382 (7)	0.0511 (8)	-0.0093 (6)	-0.0174 (6)	-0.0103 (6)
O3	0.0593 (8)	0.0273 (6)	0.0638 (9)	-0.0037 (6)	-0.0298 (6)	-0.0009 (6)
C11	0.0708 (4)	0.0261 (2)	0.0743 (4)	0.00032 (19)	-0.0426 (3)	0.0058 (2)

Geometric parameters (Å, °)

C1—C2	1.383 (2)	C7—N1	1.334 (2)
C1—C6	1.392 (2)	C7—C8	1.481 (2)
C1—N1	1.4170 (19)	C8—C9	1.330 (2)
C2—C3	1.382 (2)	C8—H8	0.9300
C2—H2	0.9300	C9—C10	1.493 (2)
C3—C4	1.386 (2)	C9—H9	0.9300
C3—C11	1.7364 (16)	C10—O2	1.212 (2)
C4—C5	1.384 (2)	C10—O3	1.296 (2)
C4—C11	1.500 (2)	C11—H11A	0.9600
C5—C6	1.378 (2)	C11—H11B	0.9600
C5—H5	0.9300	C11—H11C	0.9600
C6—H6	0.9300	N1—H1N	0.869 (15)

C7—O1	1.2410 (19)	O3—H3O	0.871 (16)
C2—C1—C6	119.32 (14)	N1—C7—C8	114.71 (13)
C2—C1—N1	124.51 (13)	C9—C8—C7	128.69 (15)
C6—C1—N1	116.16 (13)	C9—C8—H8	115.7
C3—C2—C1	118.80 (14)	C7—C8—H8	115.7
C3—C2—H2	120.6	C8—C9—C10	132.10 (15)
C1—C2—H2	120.6	C8—C9—H9	113.9
C2—C3—C4	123.73 (14)	C10—C9—H9	113.9
C2—C3—C11	117.95 (12)	O2—C10—O3	120.32 (16)
C4—C3—C11	118.31 (12)	O2—C10—C9	118.75 (16)
C5—C4—C3	115.58 (14)	O3—C10—C9	120.92 (14)
C5—C4—C11	121.39 (14)	C4—C11—H11A	109.5
C3—C4—C11	123.03 (14)	C4—C11—H11B	109.5
C6—C5—C4	122.79 (15)	H11A—C11—H11B	109.5
C6—C5—H5	118.6	C4—C11—H11C	109.5
C4—C5—H5	118.6	H11A—C11—H11C	109.5
C5—C6—C1	119.74 (15)	H11B—C11—H11C	109.5
C5—C6—H6	120.1	C7—N1—C1	128.23 (13)
C1—C6—H6	120.1	C7—N1—H1N	115.1 (13)
O1—C7—N1	122.59 (14)	C1—N1—H1N	116.2 (13)
O1—C7—C8	122.70 (14)	C10—O3—H3O	108.9 (16)
C6—C1—C2—C3	0.9 (3)	C2—C1—C6—C5	-1.5 (3)
N1—C1—C2—C3	-178.44 (16)	N1—C1—C6—C5	177.89 (15)
C1—C2—C3—C4	0.6 (3)	O1—C7—C8—C9	3.8 (3)
C1—C2—C3—C11	-178.99 (13)	N1—C7—C8—C9	-176.65 (18)
C2—C3—C4—C5	-1.5 (3)	C7—C8—C9—C10	1.3 (3)
C11—C3—C4—C5	178.12 (13)	C8—C9—C10—O2	173.92 (19)
C2—C3—C4—C11	178.21 (16)	C8—C9—C10—O3	-5.7 (3)
C11—C3—C4—C11	-2.2 (2)	O1—C7—N1—C1	-3.5 (3)
C3—C4—C5—C6	0.9 (3)	C8—C7—N1—C1	177.01 (15)
C11—C4—C5—C6	-178.86 (17)	C2—C1—N1—C7	6.7 (3)
C4—C5—C6—C1	0.6 (3)	C6—C1—N1—C7	-172.69 (16)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N \cdots O2 ⁱ	0.87 (2)	2.11 (2)	2.9546 (19)	164 (2)
O3—H3O \cdots O1	0.87 (2)	1.62 (2)	2.4885 (17)	173 (2)

Symmetry code: (i) $-x+2, y-1/2, -z+3/2$.