

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5 α -Hydroxyeudesm-4(15),11(13)-dien-8 β ,12-olide

Xue Gao* and Gang Chen

Research Center of Medical Chemistry and Chemical Biology, Chongqing Technology and Business University, Chongqing 400067, People's Republic of China
Correspondence e-mail: gaoxue@ctbu.edu.cn


Received 8 March 2012; accepted 22 March 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å;
 R factor = 0.036; wR factor = 0.084; data-to-parameter ratio = 8.0.

The title compound, $\text{C}_{15}\text{H}_{20}\text{O}_3$, a sesquiterpene lactone, was isolated from the aerial parts of *Carpesium minus* Hemsl. (Compositae). The molecule is composed of three rings, with the two cyclohexane rings in chair conformations and the cyclopentane ring adopting a twist conformation. The A/B ring junction is *trans*-fused. The absolute configuration shown has been arbitrarily assigned. In the crystal, molecules are linked into [100] chains by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the isolation and biological activity of the title compound, see: Lee *et al.* (2002); Yang *et al.* (2002); Li *et al.* (2011). For conformational analysis, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{20}\text{O}_3$
 $M_r = 248.31$
Monoclinic, $P2_1$
 $a = 7.893$ (2) Å
 $b = 7.034$ (2) Å
 $c = 12.166$ (4) Å
 $\beta = 101.154$ (3)°
 $V = 662.7$ (3) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 296$ K
0.23 × 0.20 × 0.19 mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2006)
 $T_{\min} = 0.981$, $T_{\max} = 0.984$
3673 measured reflections
1323 independent reflections
1159 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.084$
 $S = 1.08$
1323 reflections
165 parameters
1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.12$ e Å⁻³
 $\Delta\rho_{\min} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|---|--------------|--------------------|-------------|----------------------|
| $\text{O1}-\text{H1A}\cdots\text{O3}^i$ | 0.82 | 2.06 | 2.868 (2) | 168 |

Symmetry code: (i) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the Ministry of Education of Chongqing (grant No. KJ100719) and the Innovative Research Team Development Program of the University of Chongqing (grant No. KJTD201020).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RN2102).

References

- Bruker (2006). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
Lee, J. S., Min, B. S., Lee, S. M., Na, M. K., Kwon, B. M., Lee, C. O., Kim, Y. H. & Bae, K. H. (2002). *Planta Med.* **68**, 745–747.
Li, X. W., Weng, L., Gao, X., Zhao, Y., Pang, F., Liu, J. H., Zhang, H. F. & Hu, J. F. (2011). *Bioorg. Med. Chem. Lett.* **21**, 366–372.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Yang, C., Shi, Y. P. & Jia, Z. J. (2002). *Planta Med.* **68**, 626–630.

supporting information

Acta Cryst. (2012). E68, o1208 [https://doi.org/10.1107/S1600536812012470]

5 α -Hydroxyeudesm-4(15),11(13)-dien-8 β ,12-olide**Xue Gao and Gang Chen****S1. Comment**

The title compound, 5 α -hydroxyeudesm-4(15),11(13)-dien-8 β ,12-olide (Fig.1), was isolated from the medicinal plant *Carpesium minus* (Compositae). This plant has been used to reduce swelling, relieve pain and as a detoxifying agent. The compound was identified by NMR spectra, which were compared with the previous reports (Lee *et al.*, 2002; Yang *et al.*, 2002; Li *et al.*, 2011). Herewith, we present its crystal structure.

The molecule of the title compound has three fused rings consisting of two six- and one five-membered rings (A/B/C). The A/B ring junction is *trans*-fused and B/C is *cis*-fused. The two cyclohexane rings have chair conformations with puckering parameters (Cremer & Pople,1975) $Q = 0.571$ (2) Å, $\theta = 175.7$ (2) $^\circ$ and $\varphi = 134$ (4) $^\circ$ for the A ring and $Q = 0.512$ (2) Å, $\theta = 156.4$ (2) $^\circ$ and $\varphi = 344.9$ (6) $^\circ$ for the B ring; the cyclopentane ring adopts a twist conformation with puckering parameters $Q = 0.258$ (2) Å and $\varphi = 237.1$ (4) $^\circ$. In the crystal, the molecules are linked into chains by intermolecular O—H \cdots O hydrogen bonds.

S2. Experimental

The air-dried whole plants of *Carpesium minus* (3.1 g) were pulverized and extracted with 95% EtOH and yielded 439 g of crude extract, which was then suspended in 2 L water. The suspension was partitioned with EtOAc (3 \times 800 ml) to give a EtOAc-soluble portion, and a water-soluble fraction. After removal of the EtOAc under reduced pressure, 356 g of dark residue was obtained, and this was subjected to silica-gel chromatography, eluted with a stepwise gradient solvent system of petroleum/acetone 50: 1 to 0: 1 (v/v), to yield six major fractions (monitored by TLC). The third fraction (68 g) was rechromatographed on silica gel using a chloroform/MeOH (1: 0 to 30: 1) system and three fractions (Fr.A—Fr.C) were collected. Fr.B was further fractionated on a silica gel column using petroleum/EtOAc (3: 1) to give pure the title compound as colorless crystals.

S3. Refinement

All H atoms were placed in geometrically calculated positions, and allowed to ride on their parent atoms with O—H = 0.82 Å and C—H = 0.93–0.98 Å, and with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H atoms and hydroxyl group H atoms, and $x = 1.2$ for all other H atoms. In the absence of significant anomalous scattering, Friedel pairs were merged and the absolute configuration is arbitrary.




Figure 1

The molecular structure of the compound, with atom labels and 50% probability displacement ellipsoids.

5 α -Hydroxyeudesm-4(15),11 (13)-dien-8 β ,12-olide

Crystal data

C₁₅H₂₀O₃

$M_r = 248.31$

Monoclinic, $P2_1$

$a = 7.893$ (2) Å

$b = 7.034$ (2) Å

$c = 12.166$ (4) Å

$\beta = 101.154$ (3)°

$V = 662.7$ (3) Å³

$Z = 2$

$F(000) = 268$

$D_x = 1.244$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1583 reflections

$\theta = 2.9$ – 23.8 °

$\mu = 0.09$ mm⁻¹

$T = 296$ K

Block, colorless

$0.23 \times 0.20 \times 0.19$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2006)

$T_{\min} = 0.981$, $T_{\max} = 0.984$

3673 measured reflections

1323 independent reflections

1159 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 25.5$ °, $\theta_{\min} = 2.9$ °

$h = -9 \rightarrow 9$

$k = -8 \rightarrow 7$

$l = -13 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.084$

$S = 1.08$

1323 reflections

165 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.046P)^2 + 0.026P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.12 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|------------|-------------|--------------|----------------------------------|
| C1 | 0.5831 (3) | 0.5040 (4) | 0.2805 (2) | 0.0451 (6) |
| H1 | 0.5843 | 0.6432 | 0.2844 | 0.054* |
| C2 | 0.6238 (3) | 0.4471 (4) | 0.1686 (2) | 0.0462 (6) |
| H2A | 0.7088 | 0.5353 | 0.1504 | 0.055* |
| H2B | 0.5197 | 0.4618 | 0.1120 | 0.055* |
| C3 | 0.6920 (3) | 0.2446 (4) | 0.16056 (19) | 0.0404 (6) |
| C4 | 0.7625 (4) | 0.2265 (4) | 0.0514 (2) | 0.0534 (7) |
| H4A | 0.6677 | 0.2421 | -0.0118 | 0.064* |
| H4B | 0.8442 | 0.3285 | 0.0486 | 0.064* |
| C5 | 0.8506 (4) | 0.0380 (5) | 0.0400 (2) | 0.0663 (9) |
| H5A | 0.8998 | 0.0396 | -0.0272 | 0.080* |
| H5B | 0.7660 | -0.0635 | 0.0325 | 0.080* |
| C6 | 0.9931 (4) | -0.0002 (5) | 0.1417 (2) | 0.0619 (8) |
| H6A | 1.0384 | -0.1275 | 0.1367 | 0.074* |
| H6B | 1.0868 | 0.0893 | 0.1425 | 0.074* |
| C7 | 0.9243 (3) | 0.0189 (4) | 0.2486 (2) | 0.0428 (6) |
| C8 | 0.8428 (3) | 0.2111 (3) | 0.26178 (18) | 0.0370 (5) |
| C9 | 0.7793 (3) | 0.2326 (3) | 0.37155 (18) | 0.0364 (5) |
| H9A | 0.8753 | 0.2109 | 0.4332 | 0.044* |
| H9B | 0.6931 | 0.1357 | 0.3752 | 0.044* |
| C10 | 0.7006 (3) | 0.4286 (3) | 0.38653 (18) | 0.0391 (5) |
| H10 | 0.7921 | 0.5208 | 0.4142 | 0.047* |
| C11 | 0.5786 (3) | 0.4143 (3) | 0.46605 (19) | 0.0397 (5) |
| C12 | 0.4025 (3) | 0.4029 (3) | 0.3975 (2) | 0.0439 (6) |
| C13 | 0.5451 (3) | 0.0994 (4) | 0.1587 (2) | 0.0491 (6) |
| H13A | 0.4607 | 0.1147 | 0.0910 | 0.074* |
| H13B | 0.4917 | 0.1202 | 0.2222 | 0.074* |
| H13C | 0.5915 | -0.0271 | 0.1618 | 0.074* |
| C14 | 0.9291 (3) | -0.1238 (4) | 0.3199 (2) | 0.0514 (6) |
| H14A | 0.9759 | -0.2401 | 0.3049 | 0.062* |
| H14B | 0.8857 | -0.1079 | 0.3852 | 0.062* |
| C15 | 0.6070 (3) | 0.4053 (4) | 0.5758 (2) | 0.0516 (7) |

| | | | | |
|------|--------------|------------|--------------|------------|
| H15A | 0.5149 | 0.3913 | 0.6127 | 0.062* |
| H15B | 0.7192 | 0.4128 | 0.6168 | 0.062* |
| O1 | 0.9648 (2) | 0.3595 (3) | 0.25349 (15) | 0.0503 (5) |
| H1A | 1.0515 | 0.3445 | 0.3021 | 0.075* |
| O2 | 0.40859 (19) | 0.4391 (3) | 0.28997 (13) | 0.0530 (5) |
| O3 | 0.2681 (2) | 0.3667 (3) | 0.42676 (14) | 0.0585 (5) |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|-------------|--------------|
| C1 | 0.0446 (14) | 0.0397 (13) | 0.0508 (14) | 0.0042 (11) | 0.0082 (11) | 0.0056 (12) |
| C2 | 0.0434 (13) | 0.0513 (15) | 0.0420 (13) | 0.0031 (12) | 0.0038 (10) | 0.0127 (12) |
| C3 | 0.0365 (12) | 0.0485 (15) | 0.0358 (12) | 0.0004 (11) | 0.0058 (10) | 0.0046 (11) |
| C4 | 0.0549 (15) | 0.0681 (19) | 0.0381 (13) | 0.0003 (15) | 0.0112 (11) | 0.0055 (13) |
| C5 | 0.0695 (19) | 0.083 (2) | 0.0496 (16) | 0.0125 (18) | 0.0210 (14) | -0.0051 (15) |
| C6 | 0.0571 (17) | 0.072 (2) | 0.0608 (17) | 0.0137 (15) | 0.0213 (14) | -0.0028 (15) |
| C7 | 0.0332 (12) | 0.0468 (14) | 0.0473 (14) | 0.0009 (11) | 0.0050 (10) | -0.0027 (12) |
| C8 | 0.0299 (11) | 0.0393 (12) | 0.0405 (12) | -0.0064 (10) | 0.0037 (9) | 0.0032 (10) |
| C9 | 0.0331 (12) | 0.0388 (13) | 0.0346 (12) | -0.0009 (10) | -0.0001 (9) | 0.0001 (10) |
| C10 | 0.0360 (12) | 0.0373 (13) | 0.0421 (12) | -0.0046 (11) | 0.0030 (9) | -0.0010 (11) |
| C11 | 0.0380 (12) | 0.0347 (12) | 0.0463 (13) | -0.0018 (11) | 0.0077 (10) | -0.0058 (10) |
| C12 | 0.0395 (13) | 0.0425 (14) | 0.0491 (14) | 0.0017 (11) | 0.0076 (10) | -0.0060 (11) |
| C13 | 0.0422 (14) | 0.0570 (16) | 0.0447 (14) | -0.0085 (12) | 0.0003 (11) | -0.0044 (12) |
| C14 | 0.0465 (14) | 0.0438 (15) | 0.0634 (16) | 0.0054 (12) | 0.0096 (12) | -0.0006 (13) |
| C15 | 0.0537 (15) | 0.0541 (16) | 0.0471 (15) | 0.0042 (13) | 0.0099 (11) | -0.0043 (12) |
| O1 | 0.0357 (9) | 0.0532 (11) | 0.0608 (11) | -0.0118 (8) | 0.0062 (7) | 0.0067 (9) |
| O2 | 0.0357 (9) | 0.0746 (13) | 0.0470 (10) | 0.0067 (9) | 0.0036 (7) | 0.0005 (9) |
| O3 | 0.0364 (9) | 0.0772 (13) | 0.0632 (11) | -0.0042 (9) | 0.0130 (8) | -0.0094 (10) |

Geometric parameters (Å, °)

| | | | |
|--------|-----------|----------|-----------|
| C1—O2 | 1.476 (3) | C7—C8 | 1.519 (3) |
| C1—C2 | 1.512 (3) | C8—O1 | 1.437 (3) |
| C1—C10 | 1.531 (3) | C8—C9 | 1.522 (3) |
| C1—H1 | 0.9800 | C9—C10 | 1.538 (3) |
| C2—C3 | 1.533 (4) | C9—H9A | 0.9700 |
| C2—H2A | 0.9700 | C9—H9B | 0.9700 |
| C2—H2B | 0.9700 | C10—C11 | 1.495 (3) |
| C3—C4 | 1.541 (3) | C10—H10 | 0.9800 |
| C3—C13 | 1.542 (3) | C11—C15 | 1.312 (3) |
| C3—C8 | 1.556 (3) | C11—C12 | 1.478 (3) |
| C4—C5 | 1.516 (4) | C12—O3 | 1.210 (3) |
| C4—H4A | 0.9700 | C12—O2 | 1.342 (3) |
| C4—H4B | 0.9700 | C13—H13A | 0.9600 |
| C5—C6 | 1.526 (4) | C13—H13B | 0.9600 |
| C5—H5A | 0.9700 | C13—H13C | 0.9600 |
| C5—H5B | 0.9700 | C14—H14A | 0.9300 |
| C6—C7 | 1.509 (3) | C14—H14B | 0.9300 |

| | | | |
|--------------|-------------|---------------|-------------|
| C6—H6A | 0.9700 | C15—H15A | 0.9300 |
| C6—H6B | 0.9700 | C15—H15B | 0.9300 |
| C7—C14 | 1.323 (4) | O1—H1A | 0.8200 |
| O2—C1—C2 | 110.7 (2) | O1—C8—C7 | 109.61 (18) |
| O2—C1—C10 | 104.44 (17) | O1—C8—C9 | 109.18 (19) |
| C2—C1—C10 | 117.9 (2) | C7—C8—C9 | 113.54 (19) |
| O2—C1—H1 | 107.8 | O1—C8—C3 | 104.78 (17) |
| C2—C1—H1 | 107.8 | C7—C8—C3 | 109.00 (19) |
| C10—C1—H1 | 107.8 | C9—C8—C3 | 110.37 (17) |
| C1—C2—C3 | 116.2 (2) | C8—C9—C10 | 113.71 (18) |
| C1—C2—H2A | 108.2 | C8—C9—H9A | 108.8 |
| C3—C2—H2A | 108.2 | C10—C9—H9A | 108.8 |
| C1—C2—H2B | 108.2 | C8—C9—H9B | 108.8 |
| C3—C2—H2B | 108.2 | C10—C9—H9B | 108.8 |
| H2A—C2—H2B | 107.4 | H9A—C9—H9B | 107.7 |
| C2—C3—C4 | 108.73 (19) | C11—C10—C1 | 101.97 (17) |
| C2—C3—C13 | 110.1 (2) | C11—C10—C9 | 109.97 (19) |
| C4—C3—C13 | 109.2 (2) | C1—C10—C9 | 113.80 (19) |
| C2—C3—C8 | 108.32 (19) | C11—C10—H10 | 110.3 |
| C4—C3—C8 | 108.73 (18) | C1—C10—H10 | 110.3 |
| C13—C3—C8 | 111.67 (18) | C9—C10—H10 | 110.3 |
| C5—C4—C3 | 113.5 (2) | C15—C11—C12 | 121.9 (2) |
| C5—C4—H4A | 108.9 | C15—C11—C10 | 131.1 (2) |
| C3—C4—H4A | 108.9 | C12—C11—C10 | 106.96 (19) |
| C5—C4—H4B | 108.9 | O3—C12—O2 | 121.7 (2) |
| C3—C4—H4B | 108.9 | O3—C12—C11 | 128.9 (2) |
| H4A—C4—H4B | 107.7 | O2—C12—C11 | 109.43 (19) |
| C4—C5—C6 | 111.1 (3) | C3—C13—H13A | 109.5 |
| C4—C5—H5A | 109.4 | C3—C13—H13B | 109.5 |
| C6—C5—H5A | 109.4 | H13A—C13—H13B | 109.5 |
| C4—C5—H5B | 109.4 | C3—C13—H13C | 109.5 |
| C6—C5—H5B | 109.4 | H13A—C13—H13C | 109.5 |
| H5A—C5—H5B | 108.0 | H13B—C13—H13C | 109.5 |
| C7—C6—C5 | 110.5 (2) | C7—C14—H14A | 120.0 |
| C7—C6—H6A | 109.5 | C7—C14—H14B | 120.0 |
| C5—C6—H6A | 109.5 | H14A—C14—H14B | 120.0 |
| C7—C6—H6B | 109.5 | C11—C15—H15A | 120.0 |
| C5—C6—H6B | 109.5 | C11—C15—H15B | 120.0 |
| H6A—C6—H6B | 108.1 | H15A—C15—H15B | 120.0 |
| C14—C7—C6 | 121.9 (2) | C8—O1—H1A | 109.5 |
| C14—C7—C8 | 124.4 (2) | C12—O2—C1 | 110.15 (17) |
| C6—C7—C8 | 113.6 (2) | | |
| O2—C1—C2—C3 | -83.6 (3) | C2—C3—C8—C9 | 60.6 (2) |
| C10—C1—C2—C3 | 36.5 (3) | C4—C3—C8—C9 | 178.6 (2) |
| C1—C2—C3—C4 | -168.0 (2) | C13—C3—C8—C9 | -60.9 (2) |
| C1—C2—C3—C13 | 72.4 (3) | O1—C8—C9—C10 | 56.0 (2) |

| | | | |
|--------------|--------------|----------------|-------------|
| C1—C2—C3—C8 | -49.9 (3) | C7—C8—C9—C10 | 178.61 (18) |
| C2—C3—C4—C5 | 174.0 (2) | C3—C8—C9—C10 | -58.7 (2) |
| C13—C3—C4—C5 | -65.8 (3) | O2—C1—C10—C11 | -26.1 (2) |
| C8—C3—C4—C5 | 56.2 (3) | C2—C1—C10—C11 | -149.3 (2) |
| C3—C4—C5—C6 | -54.6 (3) | O2—C1—C10—C9 | 92.3 (2) |
| C4—C5—C6—C7 | 52.6 (3) | C2—C1—C10—C9 | -31.0 (3) |
| C5—C6—C7—C14 | 120.5 (3) | C8—C9—C10—C11 | 156.08 (18) |
| C5—C6—C7—C8 | -56.6 (3) | C8—C9—C10—C1 | 42.4 (2) |
| C14—C7—C8—O1 | 127.5 (3) | C1—C10—C11—C15 | -160.4 (3) |
| C6—C7—C8—O1 | -55.5 (3) | C9—C10—C11—C15 | 78.5 (3) |
| C14—C7—C8—C9 | 5.1 (3) | C1—C10—C11—C12 | 22.0 (2) |
| C6—C7—C8—C9 | -177.9 (2) | C9—C10—C11—C12 | -99.1 (2) |
| C14—C7—C8—C3 | -118.4 (3) | C15—C11—C12—O3 | -7.9 (4) |
| C6—C7—C8—C3 | 58.6 (2) | C10—C11—C12—O3 | 170.0 (3) |
| C2—C3—C8—O1 | -56.8 (2) | C15—C11—C12—O2 | 172.5 (2) |
| C4—C3—C8—O1 | 61.2 (2) | C10—C11—C12—O2 | -9.7 (3) |
| C13—C3—C8—O1 | -178.3 (2) | O3—C12—O2—C1 | 172.3 (2) |
| C2—C3—C8—C7 | -174.07 (19) | C11—C12—O2—C1 | -8.1 (3) |
| C4—C3—C8—C7 | -56.1 (2) | C2—C1—O2—C12 | 149.8 (2) |
| C13—C3—C8—C7 | 64.5 (2) | C10—C1—O2—C12 | 22.0 (3) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|---------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O1—H1A \cdots O3 ⁱ | 0.82 | 2.06 | 2.868 (2) | 168 |

Symmetry code: (i) $x+1, y, z$.