

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetraaquabis[3-(pyridin-4-yl)benzoato- κ N]manganese(II)

Ru-Qin Gao* and Guo-Ting Li

Department of Environmental and Municipal Engineering, North China University of Water Conservancy and Electric Power, Zhengzhou 450011, People's Republic of China

Correspondence e-mail: gaorquin@ncwu.edu.cn


Received 31 March 2012; accepted 4 April 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.034; wR factor = 0.079; data-to-parameter ratio = 11.9.

In the title compound, $[\text{Mn}(\text{C}_{12}\text{H}_8\text{NO}_2)_2(\text{H}_2\text{O})_4]$, the Mn^{2+} ion lies on a twofold rotation axis and has a distorted N_2O_4 octahedral coordination geometry formed by four water O atoms in the equatorial plane and two apical pyridyl N atoms. A three-dimensional network is formed in the crystal structure by multiple $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds between the coordinating water molecules and the free carboxylate groups.

Related literature

For pyridyl-multicarboxylate-metal frameworks, see: Huang *et al.* (2007). For 3-pyridin-4-ylbenzoate compounds, see: Wu *et al.* (2011). For the isotopic Co complex, see: Wang & Li (2011).



Experimental

Crystal data

$[\text{Mn}(\text{C}_{12}\text{H}_8\text{NO}_2)_2(\text{H}_2\text{O})_4]$
 $M_r = 523.39$
 Monoclinic, $C2/c$

$a = 24.935$ (3) Å
 $b = 7.1911$ (6) Å
 $c = 13.9283$ (16) Å

$\beta = 112.199$ (13)°
 $V = 2312.4$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.63$ mm⁻¹
 $T = 293$ K
 $0.24 \times 0.20 \times 0.16$ mm

Data collection

Siemens SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.875$, $T_{\max} = 0.913$

4456 measured reflections
 2035 independent reflections
 1673 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.079$
 $S = 1.05$
 2035 reflections
 171 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3A}\cdots\text{O2}^{\text{i}}$	0.86 (2)	1.91 (2)	2.732 (2)	160 (2)
$\text{O3}-\text{H3B}\cdots\text{O1}^{\text{ii}}$	0.80 (2)	1.92 (3)	2.715 (2)	175 (2)
$\text{O4}-\text{H4A}\cdots\text{O1}^{\text{iii}}$	0.86 (3)	1.86 (3)	2.728 (2)	176 (2)
$\text{O4}-\text{H4B}\cdots\text{O2}^{\text{iv}}$	0.84 (2)	1.92 (2)	2.726 (2)	161 (2)

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 2$; (ii) $x - \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1994); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

This work was supported by the Natural Science Foundation of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5868).

References

- Huang, Y., Wu, B., Yuan, D., Xu, Y., Jiang, F. & Hong, M. (2007). *Inorg. Chem.* **46**, 1171–1176.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1994). *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Siemens (1996). *SMART*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Wang, H.-R. & Li, G.-T. (2011). *Acta Cryst.* **E67**, m1743.
 Wu, B. L., Wang, R. Y., Zhang, H. Y. & Hou, H. W. (2011). *Inorg. Chim. Acta*, **375**, 2–10.

supporting information

Acta Cryst. (2012). E68, m562 [doi:10.1107/S1600536812014602]

Tetraaquabis[3-(pyridin-4-yl)benzoato- κ N]manganese(II)

Ru-Qin Gao and Guo-Ting Li

S1. Comment

Pyridyl-containing multi-carboxylic acids have been extensively investigated on the construction of various metal-organic frameworks (Huang *et al.*, 2007). Pyridylbenzoate ligands which possess a pyridyl group and a benzoic acid group are typical unsymmetrical spacers. Very recently, a series of coordination polymers of 3-pyridin-4-ylbenzoic acid (PBC) was synthesized and characterized (Wu, *et al.*, 2011). Herein we report a new Mn(II) complex with (PBC), namely, [Mn(PBC)₂(H₂O)₄] (1) which is isostructural with the complex [Co(PBC)₂(H₂O)₄] (Wang & Li, 2011).

As showed in Fig. 1, (1) is a mononuclear complex with a twofold axis passing through the Mn(II) center along *b* axis and equally splitting the whole molecule. In (1) the Mn(II) center is ligated by four O of coordinated water molecules in the equatorial plane, and two PBC acting as monodentate ligands occupy the axial positions through their pyridyl nitrogen atoms coordinating to Mn(II). Thus the Mn(II) ion is in a six-coordinated octahedral geometry. The bond distances of Mn—O and Mn—N range from 2.1867 (16) to 2.2661 (16) Å, while the in-plane and axis-transition angles are 173.04 (6) and 175.06 (8) °, respectively, indicating a slight distortion of the octahedral coordination sphere around the Mn(II) center.

Further aggregation of the monomers (1) is formed by the multiple hydrogen-bonding between the coordinated water molecules (as donors) and the uncoordinated carboxylate groups (as acceptors) (Table 1). Hydrogen-bonding system among monomers (1) is rather complicated: each water molecule forms two O—H...O hydrogen bonds with carboxylate groups of neighbouring complex molecules, while every carboxylate group of PBC forms three hydrogen bonds. Consequently, every monomer acts as a novel six-connected supramolecular synthon to connect with six adjacent monomers. Notably, the hydrogen-bonding models of the carboxyl group of PBC play an important role in the formation of crystal structure of (1). For example, as shown in Fig. 2, the O1 atom of the carboxylate group of PBC in a hydrogen-bonding bridging mode ligates to two water molecules from two neighboring monomers, and as a result, monomers (1) are regularly arrayed in *ab* plane and linked into two-dimensional layers by strong hydrogen bonding (O3...O1, 2.715 (2) Å; O4...O1, 2.728 (2) Å). The layer structure is stabilized by forceful face-to-face π ... π stacking interactions between adjacent benzoate groups and pyridyl groups of PBC with a centroid to centroid distance of 3.62 (1) Å. Intriguingly, the benzoate group and pyridyl group of PBC distort to 27.6 (0) ° to meet the formation of hydrogen bonding. The layers are further bound together to create the three-dimensional supramolecular architecture by hydrogen bonds between the O2 atom of the carboxylate group of PBC and two water molecules in the adjacent complex molecule.

S2. Experimental

The title compound, (1), was prepared according to the following process. A mixture of MnCO₃ (0.012 g, 0.1 mmol), PBC (0.040 g, 0.2 mmol) and deionized water (10 ml) was sealed into a 25 ml Teflon-lined stainless autoclave. The autoclave was heated at 160 °C for four days. As cooled to room temperature gradually, pale yellow needle crystals of (1) suitable for X-ray analysis were obtained in 64% yield (based on Mn).

S3. Refinement

All H atoms were located in a difference map. The coordinates of the water H atoms were refined with $U(H)$ set to $1.2U_{eq}(O)$. H atoms bonded to C were refined as riding with $C-H = 0.95\text{\AA}$ and $1.2U_{eq}(C)$.




Figure 1

ORTEP diagram of (1) with atom numbering scheme (30% probability ellipsoids for all non-hydrogen atoms). Symmetry code: (i) $-x, y, -z + 3/2$.




Figure 2

View of regular arrangement of monomers (1) directed by strong hydrogen bonding to form two-dimensional layers and face-to-face $\pi\cdots\pi$ stacking interactions between adjacent benzoate and pyridyl groups of PBC. Symmetry codes: (i) $-x, y, -z + 3/2$; (ii) $x - 1/2, y + 1/2, z$; (iii) $-x + 1/2, y + 1/2, -z + 3/2$; (iv) $-x + 1/2, y - 1/2, -z + 3/2$; (v) $x - 1/2, y - 1/2, z$.

Tetraaquabis[3-(pyridin-4-yl)benzoato- κN]manganese(II)

Crystal data

$[\text{Mn}(\text{C}_{12}\text{H}_8\text{NO}_2)_2(\text{H}_2\text{O})_4]$

$M_r = 523.39$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 24.935 (3) \text{\AA}$

$b = 7.1911 (6) \text{\AA}$

$c = 13.9283 (16) \text{\AA}$

$\beta = 112.199 (13)^\circ$

$V = 2312.4 (4) \text{\AA}^3$

$Z = 4$

$F(000) = 1084$

$D_x = 1.503 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{\AA}$

Cell parameters from 1769 reflections

$\theta = 3.0\text{--}27.6^\circ$

$\mu = 0.63 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Needle, yellow

$0.24 \times 0.20 \times 0.16 \text{ mm}$

Data collection

Siemens SMART CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scan
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.875$, $T_{\max} = 0.913$

4456 measured reflections
2035 independent reflections
1673 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 3.0^\circ$
 $h = -29 \rightarrow 29$
 $k = -5 \rightarrow 8$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.079$
 $S = 1.05$
2035 reflections
171 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0389P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.20 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.0000	0.30640 (6)	0.7500	0.03055 (16)
O1	0.46120 (6)	0.2912 (2)	0.87784 (13)	0.0494 (4)
O2	0.40494 (7)	0.2263 (2)	0.96408 (12)	0.0557 (5)
O3	0.03234 (7)	0.5054 (2)	0.87792 (12)	0.0434 (4)
O4	-0.02832 (7)	0.0811 (2)	0.63641 (12)	0.0414 (4)
N1	0.08789 (7)	0.3200 (2)	0.73735 (13)	0.0343 (4)
C1	0.41251 (9)	0.2746 (3)	0.88379 (18)	0.0373 (5)
C2	0.35888 (8)	0.3161 (3)	0.78874 (16)	0.0307 (5)
C3	0.30485 (8)	0.3071 (3)	0.79533 (16)	0.0304 (4)
H3	0.3024	0.2747	0.8597	0.036*
C4	0.25382 (8)	0.3442 (3)	0.71007 (15)	0.0296 (5)
C5	0.25909 (9)	0.3921 (3)	0.61674 (16)	0.0372 (5)
H5	0.2253	0.4182	0.5574	0.045*
C6	0.31246 (9)	0.4019 (3)	0.60960 (16)	0.0422 (6)
H6	0.3151	0.4351	0.5455	0.051*

C7	0.36272 (9)	0.3639 (3)	0.69523 (17)	0.0372 (5)
H7	0.3995	0.3707	0.6895	0.045*
C8	0.19674 (8)	0.3344 (3)	0.71925 (15)	0.0294 (5)
C9	0.19034 (9)	0.3677 (3)	0.81302 (16)	0.0368 (5)
H9	0.2233	0.3967	0.8732	0.044*
C10	0.13683 (9)	0.3587 (3)	0.81876 (17)	0.0384 (5)
H10	0.1341	0.3812	0.8840	0.046*
C11	0.09364 (9)	0.2880 (3)	0.64725 (17)	0.0362 (5)
H11	0.0598	0.2597	0.5884	0.043*
C12	0.14589 (9)	0.2939 (3)	0.63492 (16)	0.0364 (5)
H12	0.1473	0.2703	0.5688	0.044*
H3A	0.0499 (10)	0.451 (3)	0.9365 (18)	0.055*
H4A	-0.0072 (10)	-0.013 (3)	0.6343 (18)	0.055*
H3B	0.0120 (11)	0.588 (3)	0.8818 (18)	0.055*
H4B	-0.0448 (10)	0.126 (3)	0.5770 (19)	0.055*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0213 (3)	0.0359 (3)	0.0354 (3)	0.000	0.0119 (2)	0.000
O1	0.0229 (8)	0.0438 (9)	0.0791 (12)	-0.0003 (7)	0.0165 (8)	0.0015 (8)
O2	0.0358 (9)	0.0824 (12)	0.0427 (9)	-0.0011 (8)	0.0076 (8)	0.0058 (9)
O3	0.0351 (10)	0.0453 (10)	0.0458 (10)	0.0083 (7)	0.0107 (8)	-0.0068 (8)
O4	0.0409 (10)	0.0388 (9)	0.0435 (9)	0.0052 (7)	0.0147 (8)	-0.0010 (7)
N1	0.0263 (9)	0.0370 (10)	0.0409 (10)	0.0025 (8)	0.0139 (8)	0.0045 (8)
C1	0.0272 (12)	0.0313 (11)	0.0493 (13)	-0.0015 (10)	0.0098 (10)	-0.0065 (10)
C2	0.0238 (11)	0.0273 (10)	0.0420 (12)	-0.0025 (9)	0.0136 (9)	-0.0063 (9)
C3	0.0271 (11)	0.0300 (10)	0.0365 (11)	-0.0015 (9)	0.0149 (9)	-0.0028 (9)
C4	0.0254 (11)	0.0277 (11)	0.0388 (11)	-0.0013 (9)	0.0157 (9)	-0.0013 (9)
C5	0.0293 (12)	0.0440 (12)	0.0368 (12)	-0.0019 (10)	0.0110 (10)	0.0020 (10)
C6	0.0404 (14)	0.0547 (14)	0.0383 (12)	-0.0023 (12)	0.0227 (11)	0.0028 (11)
C7	0.0294 (12)	0.0384 (12)	0.0508 (13)	-0.0050 (10)	0.0229 (11)	-0.0056 (10)
C8	0.0248 (11)	0.0276 (11)	0.0373 (11)	0.0021 (9)	0.0134 (9)	0.0050 (9)
C9	0.0255 (11)	0.0453 (12)	0.0387 (12)	-0.0006 (10)	0.0110 (10)	-0.0005 (10)
C10	0.0287 (12)	0.0524 (13)	0.0366 (11)	0.0023 (10)	0.0152 (10)	0.0000 (10)
C11	0.0241 (11)	0.0423 (12)	0.0398 (12)	0.0009 (10)	0.0094 (9)	0.0006 (10)
C12	0.0295 (12)	0.0443 (12)	0.0377 (11)	0.0001 (10)	0.0154 (10)	-0.0011 (10)

Geometric parameters (Å, °)

Mn1—O4	2.1867 (16)	C3—C4	1.400 (3)
Mn1—O4 ⁱ	2.1867 (16)	C3—H3	0.9500
Mn1—O3	2.1878 (16)	C4—C5	1.398 (3)
Mn1—O3 ⁱ	2.1878 (16)	C4—C8	1.478 (3)
Mn1—N1	2.2661 (16)	C5—C6	1.373 (3)
Mn1—N1 ⁱ	2.2661 (16)	C5—H5	0.9500
O1—C1	1.253 (2)	C6—C7	1.392 (3)
O2—C1	1.252 (3)	C6—H6	0.9500

O3—H3A	0.86 (2)	C7—H7	0.9500
O3—H3B	0.80 (2)	C8—C9	1.395 (3)
O4—H4A	0.86 (3)	C8—C12	1.395 (3)
O4—H4B	0.84 (2)	C9—C10	1.368 (3)
N1—C11	1.336 (3)	C9—H9	0.9500
N1—C10	1.344 (3)	C10—H10	0.9500
C1—C2	1.514 (3)	C11—C12	1.378 (3)
C2—C7	1.385 (3)	C11—H11	0.9500
C2—C3	1.386 (3)	C12—H12	0.9500
O4—Mn1—O4 ⁱ	84.40 (9)	C2—C3—C4	121.98 (18)
O4—Mn1—O3	173.04 (6)	C2—C3—H3	119.0
O4 ⁱ —Mn1—O3	88.64 (6)	C4—C3—H3	119.0
O4—Mn1—O3 ⁱ	88.64 (6)	C5—C4—C3	117.47 (18)
O4 ⁱ —Mn1—O3 ⁱ	173.04 (6)	C5—C4—C8	121.60 (18)
O3—Mn1—O3 ⁱ	98.31 (10)	C3—C4—C8	120.92 (18)
O4—Mn1—N1	91.89 (6)	C6—C5—C4	120.93 (19)
O4 ⁱ —Mn1—N1	91.76 (6)	C6—C5—H5	119.5
O3—Mn1—N1	88.02 (6)	C4—C5—H5	119.5
O3 ⁱ —Mn1—N1	88.75 (6)	C5—C6—C7	120.74 (19)
O4—Mn1—N1 ⁱ	91.76 (6)	C5—C6—H6	119.6
O4 ⁱ —Mn1—N1 ⁱ	91.89 (6)	C7—C6—H6	119.6
O3—Mn1—N1 ⁱ	88.75 (6)	C2—C7—C6	119.66 (19)
O3 ⁱ —Mn1—N1 ⁱ	88.02 (6)	C2—C7—H7	120.2
N1—Mn1—N1 ⁱ	175.06 (8)	C6—C7—H7	120.2
Mn1—O3—H3A	112.0 (16)	C9—C8—C12	115.78 (18)
Mn1—O3—H3B	119.7 (18)	C9—C8—C4	121.83 (18)
H3A—O3—H3B	113 (2)	C12—C8—C4	122.39 (18)
Mn1—O4—H4A	124.7 (16)	C10—C9—C8	120.39 (19)
Mn1—O4—H4B	109.7 (17)	C10—C9—H9	119.8
H4A—O4—H4B	110 (2)	C8—C9—H9	119.8
C11—N1—C10	116.28 (18)	N1—C10—C9	123.71 (19)
C11—N1—Mn1	121.06 (14)	N1—C10—H10	118.1
C10—N1—Mn1	122.66 (14)	C9—C10—H10	118.1
O2—C1—O1	124.2 (2)	N1—C11—C12	123.65 (19)
O2—C1—C2	117.03 (19)	N1—C11—H11	118.2
O1—C1—C2	118.8 (2)	C12—C11—H11	118.2
C7—C2—C3	119.22 (19)	C11—C12—C8	120.20 (19)
C7—C2—C1	121.28 (19)	C11—C12—H12	119.9
C3—C2—C1	119.50 (18)	C8—C12—H12	119.9

Symmetry code: (i) $-x, y, -z+3/2$.*Hydrogen-bond geometry* (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3A \cdots O2 ⁱⁱ	0.86 (2)	1.91 (2)	2.732 (2)	160 (2)
O3—H3B \cdots O1 ⁱⁱⁱ	0.80 (2)	1.92 (3)	2.715 (2)	175 (2)

O4—H4A···O1 ^{iv}	0.86 (3)	1.86 (3)	2.728 (2)	176 (2)
O4—H4B···O2 ^v	0.84 (2)	1.92 (2)	2.726 (2)	161 (2)

Symmetry codes: (ii) $-x+1/2, -y+1/2, -z+2$; (iii) $x-1/2, y+1/2, z$; (iv) $-x+1/2, y-1/2, -z+3/2$; (v) $x-1/2, -y+1/2, z-1/2$.