

5-(Thiophen-2-ylmethyl)-1,3,4-thiadiazol-2-amine

Yavuz Köysal,^a Sadık Deniz,^b Ray J. Butcher,^{c*} Sema Öztürk Yildirim,^{c,d} Jerry P. Jasinski^e and Amanda C. Keeley^e

^aYeşilyurt Demir Çelik Vocational School, Ondokuz Mayıs University, Samsun, Turkey, ^bDepartment of Chemistry, Karadeniz Technical University, 61080 Trabzon, Turkey, ^cDepartment of Chemistry, Howard University, 525 College Street, NW, Washington, DC 2059, USA, ^dDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, and ^eDepartment of Chemistry, Keene State College, 220 Main Street, Keene, NH 03435-2001, USA
Correspondence e-mail: rbutcher99@yahoo.com


Received 28 March 2012; accepted 29 March 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.045; wR factor = 0.120; data-to-parameter ratio = 14.0.

In the title molecule, $\text{C}_7\text{H}_7\text{N}_3\text{S}_2$, the dihedral angle between the thiophene and thiadiazole rings is 72.99 (5)°; the two rings are oriented so that the S atoms in each ring are on the same side. In the crystal, the three-dimensional network involves strong $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, as well as $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ stacking interactions [centroid-centroid distances = 3.654 (1) and 3.495 (1) Å].

Related literature

For the antitumor activity of 2-amino-1,3,4-thiadiazole, 2-ethylamino-1,3,4-thiadiazole and 2,2'-(methylenediamino)bis-1,3,4-thiadiazole, see: Olesan *et al.* (1955); Mishra *et al.* (1995). For their anti-HIV, antiproliferative, germicidal and D2 dopaminergic activity, see: Mohareb *et al.* (2004). For the synthesis of the title compound, see: Sancak *et al.*, (2007). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_7\text{H}_7\text{N}_3\text{S}_2$

$M_r = 197.28$

Monoclinic, $P2_1/c$
 $a = 11.2970$ (6) Å
 $b = 6.6094$ (3) Å
 $c = 11.2480$ (6) Å
 $\beta = 97.243$ (5)°
 $V = 833.15$ (7) Å³

$Z = 4$
Cu $K\alpha$ radiation
 $\mu = 5.33$ mm⁻¹
 $T = 173$ K
 $0.46 \times 0.28 \times 0.15$ mm

Data collection

Agilent Xcalibur Eos Gemini diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.209$, $T_{\max} = 0.450$

4375 measured reflections
1539 independent reflections
1497 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.120$
 $S = 1.09$
1539 reflections

110 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.64$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.38$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg is the centroid of the S1/C1–C4 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3B}\cdots\text{N1}^{\text{i}}$	0.86	2.13	2.991 (2)	175
$\text{N3}-\text{H3A}\cdots\text{N2}^{\text{ii}}$	0.86	2.17	3.013 (2)	167
$\text{C1}-\text{H1}\cdots\text{Cg}^{\text{iii}}$	0.93	2.83	3.549 (2)	135

Symmetry codes: (i) $x, -y + \frac{5}{2}, z - \frac{1}{2}$; (ii) $-x + 2, -y + 3, -z + 1$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

RJB acknowledges the NSF-MRI program (grant No. CHE-0619278) for funds to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG5202).

References

- Agilent (2010). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies, Yarnton, England.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Mishra, L., Said, M. K., Itokawa, H. & Takeva, K. (1995). *Bioorg. Med. Chem.* **3**, 1241–1245.
- Mohareb, M., Sherif, M., Gaber, M., Ghabrial, S. & Aziz, I. (2004). *Heteroat. Chem.* **15**, 15–20.
- Olesan, J. J., Sloboda, A., Troy, W. P., Halliday, S. L., Landes, M. J., Angier, R. B., Semb, J., Cvr, K. & Williams, J. H. (1955). *J. Am. Chem. Soc.* **77**, 6713–6714.
- Sancak, K., Ünver, Y. & Er, M. (2007). *Turk. J. Chem.* **31**, 125–134.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o1279 [https://doi.org/10.1107/S1600536812013633]

5-(Thiophen-2-ylmethyl)-1,3,4-thiadiazol-2-amine

Yavuz Köysal, Sadık Deniz, Ray J. Butcher, Sema Öztürk Yildirim, Jerry P. Jasinski and Amanda C. Keeley

S1. Comment

The antitumor activities of 2-amino-1,3,4-thiadiazole (ATDA, NSC-4728) and the related compounds: 2-ethyl-amino-1,3,4-thiadiazole (EATDA), 2,2'-(methylene-diamino) bis-1,3,4-thiadiazole (NSC-143019) were found in several experimental tumor systems about 50 years ago (Olesan *et al.*, 1955). 2-Amino-1,3,4-thiadiazole (ATDA), as the most promising compound, was used in phase II clinical trials in patients with different tumors: renal, colon, ovarian, and others. Recently new derivatives with the 1,3,4-thiadiazole nucleus as well as Fe(II) / Fe (III) complexes of 2-amino-1,3,4-thiadiazoles have been synthesized and evaluated for their antiproliferative activity against a panel of human cancer cell lines (Mishra *et al.*, 1995). Over recent years, there has been an increasing interest in the chemistry of thiophenes because of their biological significance. Many of them have been widely investigated for therapeutic uses, especially as antifungal, antibacterial, anti-inflammatory, anticonvulsant, antiasthmatic, and analgesic agents. They also were known to show anti-HIV, antiproliferative, germicidal, and D2 dopaminergic activities (Mohareb *et al.*, 2004). In view of these facts, the aim of this present study is to obtain a structure of 1,3,4-oxadiazole incorporating the thiophene ring.

In the molecule of the title compound (Fig 1), the bond lengths are within normal ranges (Allen *et al.*, 1987). In the molecule of (I) atom S1 is oriented towards the thiadiazol ring, Fig. 1. The dihedral angle between the planar thiophene (r.m.s. deviation = 0.007 Å) and planar thiadiazol (r.m.s. deviation = 0.004 Å) rings of 72.99 (5)° indicates a twist between planes as seen in the S1–C4–C5–C6 torsion angle of 94.86 (17) °. The amine group is effectively co-planar with the thiadiazol ring to which it is attached as seen in the N3–C7–S2–C6 torsion angle of 178.53 (16) °.

In the crystal structure, there are strong intermolecular N–H···N hydrogen bonds which lead to the formation of centrosymmetric dimers in the crystal. In addition there are C–H··· π and π - π stacking interactions [$Cg1 \cdots Cg1(1-x, -y, -z) = 3.654(1)$ Å and $Cg2 \cdots Cg2(-x, 1-y, -z) = 3.495(1)$ Å, $Cg1(S1/C1-C4)$ and $Cg2(S2/N1/N2/C6/C7)$ are the centroids of the thiophene and thiadiazol rings]. This pattern is the primary supramolecular structure for this compound (Fig. 2).

S2. Experimental

The title compound was synthesized using the published method (Sancak *et al.*, 2007).

S3. Refinement

The amine H atoms were seen in a difference Fourier map and then idealized with $U_{iso}(H) = 1.2U_{eq}(N)$ with N–H bond length of 0.86 Å. The C-bound H-atoms were positioned geometrically with C–H = 0.93 and 0.97 Å, for aromatic and CH₂ H-atoms, respectively, and constrained to ride on their parent atoms, with $U_{iso}(H) = 1.2U_{eq}(C)$. During the refinement it was noticed that for the strongest reflections ($F_o/F_c(\max)$ close to 1.00) the observed value (F_o) was much smaller than the calculated value (F_c) indicating detector saturation problems. These reflections were omitted from the refinement.




Figure 1

View of the molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level for non-hydrogen atoms.




Figure 2

The packing view showing the hydrogen bonds network. Dashed lines indicate intermolecular N—H...N hydrogen bonds (see Table 1 for details).

5-(Thiophen-2-ylmethyl)-1,3,4-thiadiazol-2-amine

Crystal data

$C_7H_7N_3S_2$

$M_r = 197.28$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1\ ybc$

$a = 11.2970\ (6)\ \text{\AA}$

$b = 6.6094\ (3)\ \text{\AA}$

$c = 11.2480\ (6)\ \text{\AA}$

$\beta = 97.243\ (5)^\circ$

$V = 833.15\ (7)\ \text{\AA}^3$

$Z = 4$

$F(000) = 408$

$D_x = 1.573\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184\ \text{\AA}$

Cell parameters from 2744 reflections

$\theta = 3.9\text{--}70.0^\circ$

$\mu = 5.33\ \text{mm}^{-1}$

$T = 173\text{ K}$
 Chunk, colorless

$0.46 \times 0.28 \times 0.15\text{ mm}$

Data collection

Agilent Xcalibur Eos Gemini
 diffractometer
 Radiation source: Enhance (Cu) X-ray Source
 Graphite monochromator
 Detector resolution: $16.1500\text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.209$, $T_{\max} = 0.450$

4375 measured reflections
 1539 independent reflections
 1497 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\max} = 70.1^\circ$, $\theta_{\min} = 3.9^\circ$
 $h = -13 \rightarrow 13$
 $k = -5 \rightarrow 8$
 $l = -12 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.120$
 $S = 1.09$
 1539 reflections
 110 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0928P)^2 + 0.0849P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.64\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.38\text{ e \AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.035 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.64166 (4)	0.57641 (7)	0.32224 (4)	0.0210 (2)
S2	0.82410 (4)	1.05521 (6)	0.34150 (3)	0.0182 (2)
N1	0.87197 (12)	1.0981 (2)	0.56747 (13)	0.0175 (4)
N2	0.91662 (12)	1.2714 (2)	0.51908 (13)	0.0179 (4)
N3	0.93579 (14)	1.4168 (2)	0.33358 (14)	0.0251 (4)
H3A	0.9732	1.5198	0.3665	0.030*
H3B	0.9220	1.4085	0.2568	0.030*
C1	0.48911 (16)	0.5889 (3)	0.30609 (17)	0.0216 (4)
H1	0.4395	0.5306	0.2430	0.026*
C2	0.44958 (15)	0.6943 (3)	0.39653 (16)	0.0226 (4)
H2	0.3693	0.7160	0.4030	0.027*
C3	0.54422 (16)	0.7680 (3)	0.48045 (16)	0.0206 (4)

H3	0.5323	0.8420	0.5483	0.025*
C4	0.65455 (15)	0.7192 (2)	0.45147 (15)	0.0171 (4)
C5	0.77380 (16)	0.7709 (3)	0.51957 (16)	0.0212 (4)
H5A	0.8306	0.6668	0.5044	0.025*
H5B	0.7667	0.7689	0.6046	0.025*
C6	0.82264 (14)	0.9732 (3)	0.48849 (14)	0.0159 (4)
C7	0.89906 (14)	1.2698 (3)	0.40174 (15)	0.0165 (4)

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0161 (3)	0.0215 (4)	0.0250 (4)	−0.00005 (14)	0.0010 (2)	−0.00692 (15)
S2	0.0184 (3)	0.0192 (3)	0.0158 (3)	−0.00404 (13)	−0.0027 (2)	−0.00264 (13)
N1	0.0172 (7)	0.0156 (7)	0.0191 (7)	−0.0010 (5)	0.0000 (6)	0.0004 (5)
N2	0.0170 (7)	0.0175 (8)	0.0184 (7)	−0.0032 (5)	−0.0012 (5)	−0.0010 (5)
N3	0.0272 (9)	0.0290 (9)	0.0172 (8)	−0.0138 (6)	−0.0039 (6)	0.0017 (6)
C1	0.0172 (8)	0.0152 (8)	0.0311 (9)	−0.0020 (6)	−0.0017 (7)	0.0002 (7)
C2	0.0176 (8)	0.0169 (9)	0.0337 (10)	0.0024 (6)	0.0052 (7)	0.0078 (7)
C3	0.0248 (9)	0.0135 (9)	0.0239 (9)	0.0041 (6)	0.0054 (7)	0.0012 (6)
C4	0.0211 (8)	0.0084 (8)	0.0213 (9)	−0.0011 (6)	0.0007 (7)	0.0000 (6)
C5	0.0236 (9)	0.0130 (9)	0.0251 (9)	−0.0007 (6)	−0.0039 (7)	0.0016 (7)
C6	0.0139 (7)	0.0152 (8)	0.0179 (8)	0.0022 (6)	−0.0004 (6)	−0.0003 (6)
C7	0.0100 (7)	0.0184 (9)	0.0199 (8)	−0.0002 (6)	−0.0025 (6)	−0.0031 (6)

Geometric parameters ($\text{\AA}$, $^\circ$)

S1—C1	1.7119 (18)	C1—C2	1.354 (3)
S1—C4	1.7237 (17)	C1—H1	0.9300
S2—C6	1.7419 (17)	C2—C3	1.420 (3)
S2—C7	1.7445 (17)	C2—H2	0.9300
N1—C6	1.287 (2)	C3—C4	1.366 (2)
N1—N2	1.390 (2)	C3—H3	0.9300
N2—C7	1.310 (2)	C4—C5	1.503 (2)
N3—C7	1.336 (2)	C5—C6	1.505 (2)
N3—H3A	0.8600	C5—H5A	0.9700
N3—H3B	0.8600	C5—H5B	0.9700
C1—S1—C4	92.29 (9)	C3—C4—C5	127.66 (16)
C6—S2—C7	86.93 (8)	C3—C4—S1	110.36 (13)
C6—N1—N2	113.90 (14)	C5—C4—S1	121.96 (13)
C7—N2—N1	111.83 (14)	C4—C5—C6	114.53 (14)
C7—N3—H3A	120.0	C4—C5—H5A	108.6
C7—N3—H3B	120.0	C6—C5—H5A	108.6
H3A—N3—H3B	120.0	C4—C5—H5B	108.6
C2—C1—S1	111.59 (14)	C6—C5—H5B	108.6
C2—C1—H1	124.2	H5A—C5—H5B	107.6
S1—C1—H1	124.2	N1—C6—C5	123.32 (15)
C1—C2—C3	112.57 (16)	N1—C6—S2	113.61 (13)

C1—C2—H2	123.7	C5—C6—S2	123.00 (12)
C3—C2—H2	123.7	N2—C7—N3	123.65 (16)
C4—C3—C2	113.16 (16)	N2—C7—S2	113.71 (13)
C4—C3—H3	123.4	N3—C7—S2	122.63 (13)
C2—C3—H3	123.4		
C6—N1—N2—C7	−0.27 (19)	N2—N1—C6—C5	176.73 (15)
C4—S1—C1—C2	−1.08 (14)	N2—N1—C6—S2	−0.36 (18)
S1—C1—C2—C3	0.4 (2)	C4—C5—C6—N1	136.56 (17)
C1—C2—C3—C4	0.7 (2)	C4—C5—C6—S2	−46.6 (2)
C2—C3—C4—C5	−179.68 (15)	C7—S2—C6—N1	0.64 (13)
C2—C3—C4—S1	−1.50 (19)	C7—S2—C6—C5	−176.45 (15)
C1—S1—C4—C3	1.47 (14)	N1—N2—C7—N3	−178.55 (16)
C1—S1—C4—C5	179.78 (14)	N1—N2—C7—S2	0.78 (18)
C3—C4—C5—C6	−87.1 (2)	C6—S2—C7—N2	−0.80 (13)
S1—C4—C5—C6	94.86 (17)	C6—S2—C7—N3	178.53 (16)

Hydrogen-bond geometry ($\text{\AA}$, $^\circ$)

Cg is the centroid of the S1/C1—C4 ring.

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N3—H3B $\cdots$ N1 ⁱ	0.86	2.13	2.991 (2)	175
N3—H3A $\cdots$ N2 ⁱⁱ	0.86	2.17	3.013 (2)	167
C1—H1 $\cdots$ Cg ⁱⁱⁱ	0.93	2.83	3.549 (2)	135

Symmetry codes: (i) $x, -y+5/2, z-1/2$; (ii) $-x+2, -y+3, -z+1$; (iii) $-x+1, y-1/2, -z+1/2$.