

Bis(μ -4,4';6,6'-bis(biphenyl-2,2'-diyl-dioxy)-2,2-bis{2-[5-(pyridin-4-yl)-1,3,4-oxadiazol-2-yl]phenoxy}cyclo-triphenylphosphazene)di- μ -chlorido-bis[chloridocopper(II)]

Xiang-Wen Wu, Xiao-Yan Wang, Jian-Ping Ma and Yu-Bin Dong*

College of Chemistry, Chemical Engineering and Materials Science, Key Laboratory of Molecular and Nano Probes, Engineering Research Center of Pesticide and Medicine Intermediate Clean Production, Ministry of Education, Shandong Provincial Key Laboratory of Clean Production of Fine Chemicals, Shandong Normal University, Jinan 250014, People's Republic of China
Correspondence e-mail: yubindong@sdnu.edu.cn


Received 28 March 2012; accepted 14 April 2012

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.056; wR factor = 0.139; data-to-parameter ratio = 13.1.

In the crystal of the title compound, $[\text{Cu}_2\text{Cl}_4(\text{C}_{50}\text{H}_{32}\text{N}_9\text{O}_8\text{P}_3)_2]$, the binuclear molecule is located across an inversion center. Each Cu^{2+} cation is coordinated by two pyridine N atoms from symmetry-related 4,4';6,6'-bis(biphenyl-2,2'-diyl-dioxy)-2,2-bis{2-[5-(pyridin-4-yl)-1,3,4-oxadiazol-2-yl]phenoxy}cyclo-triphenylphosphazene (*L*) ligands, a pair of bridging Cl^- anions and a terminal Cl^- anion, forming a distorted CuCl_3N_2 square-pyramidal geometry. Weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ and intermolecular $\text{C}-\text{H}\cdots\text{N}$ interactions occur in the crystal.

Related literature

For our interest in the coordination chemistry of bent organic ligands bridged by five-membered heterocycles such as oxadiazole and triazole, see: Dong *et al.* (2005, 2007). For the impact of different types of linkages as well as distinct coordinating orientations on the structures of various coordination-driven supramolecular compounds, see: Zhao *et al.* (2007). For bond lengths and angles in related structures, see: Ainscough *et al.* (2008); Du *et al.* (2010). Zhao *et al.* (2009).



Experimental

Crystal data

$[\text{Cu}_2\text{Cl}_4(\text{C}_{50}\text{H}_{32}\text{N}_9\text{O}_8\text{P}_3)_2]$
 $M_r = 2228.39$
 Triclinic, $P\bar{1}$
 $a = 10.554(5)\text{ \AA}$
 $b = 14.486(7)\text{ \AA}$
 $c = 15.551(8)\text{ \AA}$
 $\alpha = 85.401(7)^\circ$
 $\beta = 81.984(7)^\circ$
 $\gamma = 89.844(7)^\circ$
 $V = 2347(2)\text{ \AA}^3$
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.75\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.35 \times 0.30 \times 0.08\text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.779$, $T_{\max} = 0.942$
 12440 measured reflections
 8593 independent reflections
 6217 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.139$
 $S = 1.02$
 8593 reflections
 658 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.51\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.48\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry ($\text{\AA}$, $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C26}-\text{H26}\cdots\text{O3}$	0.93	2.38	3.271 (5)	160
$\text{C36}-\text{H36}\cdots\text{N5}^i$	0.93	2.51	3.396 (6)	159

Symmetry code: (i) $x, y + 1, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the NSFC (Nos. 91027003 and 21072118), the 973 Program (No. 2012CB821705), the PCSIRT, the Shangdong Natural Science Foundation (No. JQ200803) and the Taishan Scholars' Construction Project Special Fund in China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5503).

References

- Ainscough, E. W., Brodie, A. M., Davidson, R. J., Moubaraki, B., Murray, K. M., Otter, C. A. & Waterland, M. R. (2008). *Inorg. Chem.* **47**, 9182–9192.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dong, Y.-B., Wang, H.-Y., Ma, J.-P., Shen, D.-Z. & Huang, R.-Q. (2005). *Inorg. Chem.* **44**, 4679–4692.
- Dong, Y.-B., Wang, P., Ma, J.-P., Wang, H.-Y., Tang, B. & Huang, R.-Q. (2007). *J. Am. Chem. Soc.* **129**, 4872–4873.
- Du, M., Wang, Q., Li, C.-P., Zhao, X.-J. & Ribas, J. (2010). *Cryst. Growth Des.* **10**, 3285–3296.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhao, X.-X., Ma, J.-P., Dong, Y.-B., Huang, R.-Q. & Lai, T.-S. (2007). *Cryst. Growth Des.* **7**, 1058–1068.
- Zhao, X.-X., Ma, J.-P., Shen, D.-Z., Dong, Y.-B. & Huang, R.-Q. (2009). *CrystEngComm*, **11**, 1281–1290.

supporting information

Acta Cryst. (2012). E68, m648–m649 [https://doi.org/10.1107/S1600536812016285]

Bis(μ -4,4';6,6-bis(biphenyl-2,2'-diyldioxy)-2,2-bis{2-[5-(pyridin-4-yl)-1,3,4-oxadiazol-2-yl]phenoxy}cyclotriphosphazene)di- μ -chlorido-bis[chloridocopper(II)]

Xiang-Wen Wu, Xiao-Yan Wang, Jian-Ping Ma and Yu-Bin Dong

S1. Comment

In our research, we have a longstanding interest in the coordination chemistry of bent organic ligands bridged by the five-membered heterocycles such as oxadiazole (Dong *et al.*, 2005) and triazole (Dong *et al.*, 2007), especially in how different types of linkages as well as distinct coordinating orientations impact the structures of various coordination-driven supramolecular compounds (Zhao *et al.*, 2007). In this contribution, we present a new ligand containing 1,3,4-oxadiazole and cyclotriphosphazene units, namely, bis(2,2'-dioxybiphenyl)bis(2-phenoxy-5-(4-pyridyl)-1,3,4-oxadiazole) cyclotriphosphazene (**L**) and the title compound, $[\text{CuCl}_2(\text{C}_{50}\text{H}_{32}\text{O}_8\text{N}_9\text{P}_3)]_2$ based on it (Scheme 1).

In the crystal of the title compound, $[\text{CuCl}_2(\text{C}_{50}\text{H}_{32}\text{O}_8\text{N}_9\text{P}_3)]_2$, the binuclear molecule is located across an inversion center (Figure 1). The Cu^{II} cation is N,N'-chelated by a bis(2,2'-dioxybiphenyl)bis(2-phenoxy-5-(4-pyridyl)-1,3,4-oxadiazole) cyclotriphosphazene (**L**) ligand, and a pair of bridging Cl^- anions and a terminal Cl^- anion further coordinate to the Cu^{II} cation to complete the distorted CuCl_3N_2 square-pyramidal geometry (Ainscough *et al.*, 2008). The $\text{N}8^{\text{i}}\text{-Cu1-N9}$ angle is $174.68(1)^\circ$ and the dihedral angle between C1-C5/N9 and $\text{C21}^{\text{i}}\text{-C25}^{\text{i}}/\text{N8}^{\text{i}}$ is 21.272° (Zhao *et al.*, 2009) [symmetry code: $i = -x, -y, 1-z$]. The corresponding dihedral angles between the planes of the three rings in one 1,3,4-oxadiazole arm are 9.565° (between C1-C5/N9 and C6-C7-O8/N6-N7) and 5.322° (between C6-C7-O8/N6-N7 and C8-C13), and the corresponding dihedral angles of three rings in the other 1,3,4-oxadiazole arm are 30.941° (between C22-C26/N8 and C20-C21-O8/N4-N5) and 12.694° (between C20-C21-O8/N4-N5 and C14-C19). The Cu-N distances range from $2.021(3) \text{ \AA}$ to $2.045 \text{ \AA}$ which are close to those found in $\{[\text{Cu}(\text{4-bpo})(\text{dca})(\text{H}_2\text{O})_2](\text{NO}_3)(\text{H}_2\text{O})\}_n$ ($\text{4-bp} = 2,5\text{-bis}(4\text{-pyridyl})\text{-1,3,4-oxadiazole}$, $\text{dca} = \text{N}(\text{CN})_2$) (Du *et al.*, 2010). Moreover, the angles of Cl1-Cu1-Cl2 and $\text{Cl1-Cu1-Cl2}^{\text{i}}$ are $167.54(5)^\circ$ and $101.39(5)^\circ$, respectively [symmetry code: $i = -x, -y, 1-z$]. The bridged Cu-Cl bond length ($2.700(6) \text{ \AA}$) is longer than other Cu-Cl bonds ($2.261(9) \text{ \AA}$ and $2.300(8) \text{ \AA}$) (Ainscough *et al.*, 2008). Additionally, neighbouring Cu^{II} cations are connected by two bridged chlorine ions into Cu_2Cl_2 loops with a $\text{Cu}\cdots\text{Cu}$ separation $3.522(1) \text{ \AA}$.

In the solid state, binuclear molecules are linked together by weak intramolecular $\text{C}\cdots\text{H}\cdots\text{O}$ and intermolecular $\text{C}\cdots\text{H}\cdots\text{N}$ interactions into one-dimensional double-chains (Figure 2, Table 1).

S2. Experimental

Preparation of **L**:

The solution of 2-phenoxy-5-(4-pyridyl)-1,3,4-oxadiazole (0.50 g, 2.1 mmol) in 30 ml THF was added NaH (50.4 mg, 2.1 mmol), the mixture was stirred for 1 h at room temperature, then $[\text{N}_3\text{P}_3(2,2'\text{-dioxybiphenyl})_2\text{Cl}_2]$ (0.574 g, 1 mmol) in 20 ml THF was added. The mixture was heated at reflux over 3 h, allowed to cool, the precipitation was separated by filtration and washed several times with water, then the residue was purified on a silica gel column using $\text{CH}_2\text{Cl}_2\text{:MeOH} = 20\text{:1}$ as the eluent to afford **L** as a white crystalline solid. IR(KBr pellet, cm^{-1}): 1608(*m*), 1540(*w*), 1477(*m*), 1435(*m*), 1410(*w*), 1276(*m*), 1233(*m*), 1175(*s*), 1090(*m*), 1033(*m*), 932(*w*), 890(*s*), 818(*w*), 785(*m*), 750(*m*), 716(*m*), 701(*m*),

630(w), 608(m), 532(m), 409(w). $^1\text{H-NMR}$ (300 MHz, 25°C , DMSO-d_6 , TMS, p.p.m.): 8.41–8.40(d, 2H, $-\text{C}_5\text{H}_4\text{N}$), 8.26–8.24(d, 1H, $-\text{C}_6\text{H}_4\text{O}$), 8.02–7.84(d, 2H, $-\text{C}_5\text{H}_4\text{N}$), 7.76(d, 1H, $-\text{C}_6\text{H}_4\text{O}$), 7.54(t, 1H, $p\text{-C}_6\text{H}_4\text{O}$), 7.51–7.39(m, 3H, $-\text{C}_6\text{H}_4\text{O}$, $\text{C}_{12}\text{H}_8\text{O}_2$), 7.28–7.26(m, 4H, $-\text{C}_{12}\text{H}_8\text{O}_2$), 6.98–6.26(d, 3H, $-\text{C}_6\text{H}_4\text{O}$, $-\text{C}_{12}\text{H}_8\text{O}_2$). Anal. Calcd. for $\text{C}_{50}\text{H}_{32}\text{N}_9\text{O}_8\text{P}_3$: C, 61.29; H, 3.30; N, 12.86. Found: C, 61.33; H, 3.24; N, 12.89.

Preparation of Cu^{II} compound:

L (49 mg, 0.05 mmol) in CH_2Cl_2 (7 ml) and $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (9 mg, 0.05 mmol) in CH_3OH (7 ml) were mixed. After about one week, green block-shaped crystals were obtained in 75% yield based on *L*.

S3. Refinement

H atoms attached to carbon were placed in geometrically idealized positions with $\text{C-H} = 0.93 \text{ \AA}$ and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

A view of Cu^{II} compound, showing the coordination around Cu^{II} . Displacement ellipsoids are drawn at the 30% probability level and H atoms have been omitted for clarity. [symmetry code: (i) $-x, -y, -z + 1$].




Figure 2

A diagram showing weak intermolecular C-H...N interaction.

Bis(μ -4,4';6,6-bis(biphenyl-2,2'-diyl)dioxy)-2,2-bis{2-[5-(pyridin-4-yl)-1,3,4-oxadiazol-2-yl]phenoxy}cyclotriphosphazene)di- μ -chlorido-bis[chloridocopper(II)]

Crystal data

$[\text{Cu}_2\text{Cl}_4(\text{C}_{50}\text{H}_{32}\text{N}_9\text{O}_8\text{P}_3)_2]$

$M_r = 2228.39$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 10.554 (5) \text{ \AA}$

$b = 14.486 (7) \text{ \AA}$

$c = 15.551 (8) \text{ \AA}$

$\alpha = 85.401 (7)^\circ$

$\beta = 81.984 (7)^\circ$

$\gamma = 89.844 (7)^\circ$

$V = 2347 (2) \text{ \AA}^3$
 $Z = 1$
 $F(000) = 1134$
 $D_x = 1.577 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 2838 reflections

$\theta = 2.2\text{--}24.7^\circ$
 $\mu = 0.75 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
 Block, green
 $0.35 \times 0.30 \times 0.08 \text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.779$, $T_{\max} = 0.942$

12440 measured reflections
 8593 independent reflections
 6217 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$
 $\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -12 \rightarrow 12$
 $k = -17 \rightarrow 13$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.139$
 $S = 1.02$
 8593 reflections
 658 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0629P)^2 + 0.7991P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.48 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0712 (4)	−0.0313 (3)	0.2710 (3)	0.0457 (11)
H1	0.0268	−0.0868	0.2868	0.055*
C2	0.1569 (4)	−0.0249 (3)	0.1965 (3)	0.0453 (10)
H2	0.1706	−0.0753	0.1629	0.054*
C3	0.2228 (4)	0.0568 (3)	0.1715 (3)	0.0374 (9)
C4	0.2035 (4)	0.1283 (3)	0.2261 (3)	0.0465 (11)
H4	0.2500	0.1833	0.2130	0.056*
C5	0.1149 (4)	0.1168 (3)	0.2994 (3)	0.0478 (11)
H5	0.1009	0.1658	0.3347	0.057*
C6	0.3034 (4)	0.0679 (3)	0.0879 (3)	0.0374 (9)
C7	0.4205 (4)	0.1374 (3)	−0.0201 (2)	0.0362 (9)

C8	0.4932 (4)	0.2111 (3)	−0.0736 (2)	0.0341 (9)
C9	0.5458 (4)	0.1951 (3)	−0.1588 (3)	0.0462 (11)
H9	0.5371	0.1369	−0.1787	0.055*
C10	0.6096 (4)	0.2634 (3)	−0.2131 (3)	0.0518 (12)
H10	0.6462	0.2505	−0.2687	0.062*
C11	0.6206 (4)	0.3511 (3)	−0.1870 (3)	0.0467 (11)
H11	0.6610	0.3979	−0.2254	0.056*
C12	0.5710 (4)	0.3691 (3)	−0.1029 (3)	0.0379 (9)
H12	0.5781	0.4281	−0.0845	0.045*
C13	0.5108 (3)	0.2990 (3)	−0.0465 (2)	0.0317 (9)
C14	0.7375 (3)	0.1807 (2)	0.0721 (2)	0.0295 (8)
C15	0.8049 (4)	0.2253 (3)	−0.0015 (3)	0.0372 (9)
H15	0.7672	0.2738	−0.0314	0.045*
C16	0.9278 (4)	0.1988 (3)	−0.0314 (3)	0.0448 (10)
H16	0.9738	0.2301	−0.0804	0.054*
C17	0.9821 (4)	0.1253 (3)	0.0120 (3)	0.0453 (11)
H17	1.0648	0.1068	−0.0077	0.054*
C18	0.9139 (4)	0.0801 (3)	0.0839 (3)	0.0421 (10)
H18	0.9515	0.0308	0.1128	0.051*
C19	0.7891 (4)	0.1055 (3)	0.1157 (2)	0.0340 (9)
C20	0.7238 (4)	0.0482 (3)	0.1890 (3)	0.0357 (9)
C21	0.5882 (4)	−0.0021 (3)	0.2956 (3)	0.0413 (10)
C22	0.4747 (4)	−0.0044 (3)	0.3607 (3)	0.0408 (10)
C23	0.4200 (5)	−0.0890 (3)	0.3937 (3)	0.0535 (12)
H23	0.4551	−0.1438	0.3742	0.064*
C24	0.3138 (4)	−0.0910 (3)	0.4553 (3)	0.0530 (12)
H24	0.2772	−0.1483	0.4759	0.064*
C25	0.3143 (4)	0.0655 (3)	0.4566 (3)	0.0544 (12)
H25	0.2792	0.1191	0.4788	0.065*
C26	0.4198 (4)	0.0741 (3)	0.3937 (3)	0.0536 (12)
H26	0.4539	0.1323	0.3737	0.064*
C27	0.9432 (3)	0.4069 (2)	0.1586 (2)	0.0302 (8)
C28	1.0052 (4)	0.3287 (3)	0.1306 (3)	0.0423 (10)
H28	0.9665	0.2707	0.1433	0.051*
C29	1.1253 (4)	0.3382 (3)	0.0836 (3)	0.0521 (12)
H29	1.1684	0.2860	0.0641	0.062*
C30	1.1819 (4)	0.4234 (3)	0.0652 (3)	0.0525 (12)
H30	1.2630	0.4289	0.0329	0.063*
C31	1.1200 (4)	0.5013 (3)	0.0940 (3)	0.0440 (10)
H31	1.1596	0.5590	0.0809	0.053*
C32	0.9980 (3)	0.4944 (3)	0.1429 (2)	0.0330 (9)
C33	0.9357 (3)	0.5754 (2)	0.1815 (2)	0.0317 (8)
C34	1.0032 (4)	0.6362 (3)	0.2232 (3)	0.0449 (10)
H34	1.0895	0.6258	0.2264	0.054*
C35	0.9449 (5)	0.7115 (3)	0.2601 (3)	0.0538 (12)
H35	0.9917	0.7507	0.2886	0.065*
C36	0.8176 (5)	0.7291 (3)	0.2551 (3)	0.0557 (13)
H36	0.7788	0.7804	0.2796	0.067*

C37	0.7478 (4)	0.6702 (3)	0.2135 (3)	0.0436 (10)
H37	0.6624	0.6821	0.2086	0.052*
C38	0.8072 (3)	0.5931 (2)	0.1793 (2)	0.0314 (8)
C39	0.4887 (4)	0.3345 (2)	0.4399 (2)	0.0332 (9)
C40	0.6046 (4)	0.3320 (3)	0.4705 (3)	0.0419 (10)
H40	0.6766	0.3081	0.4386	0.050*
C41	0.6109 (4)	0.3662 (3)	0.5505 (3)	0.0494 (11)
H41	0.6879	0.3650	0.5732	0.059*
C42	0.5037 (4)	0.4021 (3)	0.5964 (3)	0.0478 (11)
H42	0.5083	0.4244	0.6504	0.057*
C43	0.3902 (4)	0.4050 (3)	0.5632 (3)	0.0440 (10)
H43	0.3188	0.4302	0.5948	0.053*
C44	0.3791 (4)	0.3712 (3)	0.4831 (2)	0.0356 (9)
C45	0.2570 (4)	0.3736 (3)	0.4468 (3)	0.0378 (9)
C46	0.1428 (4)	0.3526 (4)	0.4998 (3)	0.0613 (13)
H46	0.1432	0.3374	0.5590	0.074*
C47	0.0272 (5)	0.3538 (4)	0.4663 (4)	0.0782 (17)
H47	−0.0488	0.3409	0.5033	0.094*
C48	0.0249 (4)	0.3740 (4)	0.3794 (4)	0.0704 (15)
H48	−0.0522	0.3729	0.3570	0.084*
C49	0.1367 (4)	0.3958 (3)	0.3248 (3)	0.0502 (11)
H49	0.1359	0.4100	0.2655	0.060*
C50	0.2493 (3)	0.3961 (3)	0.3593 (3)	0.0341 (9)
Cl1	−0.18724 (11)	0.15324 (8)	0.35268 (7)	0.0535 (3)
Cl2	−0.02705 (10)	−0.11280 (7)	0.46697 (7)	0.0485 (3)
Cu1	−0.10057 (5)	0.02987 (3)	0.42097 (3)	0.04299 (16)
N1	0.4580 (3)	0.3014 (2)	0.2006 (2)	0.0334 (7)
N2	0.6647 (3)	0.3800 (2)	0.09859 (19)	0.0323 (7)
N3	0.5960 (3)	0.4339 (2)	0.2594 (2)	0.0349 (8)
N4	0.7652 (3)	−0.0316 (2)	0.2173 (3)	0.0536 (10)
N5	0.6752 (4)	−0.0637 (2)	0.2876 (3)	0.0573 (10)
N6	0.3977 (4)	0.0552 (3)	−0.0424 (3)	0.0599 (11)
N7	0.3211 (4)	0.0102 (3)	0.0285 (3)	0.0599 (11)
N8	0.2597 (3)	−0.0152 (2)	0.4874 (2)	0.0429 (8)
N9	0.0475 (3)	0.0384 (2)	0.3230 (2)	0.0422 (8)
O1	0.8252 (2)	0.39650 (16)	0.21153 (16)	0.0315 (6)
O2	0.7351 (2)	0.53487 (16)	0.13627 (16)	0.0332 (6)
O3	0.4819 (2)	0.29617 (16)	0.35980 (16)	0.0348 (6)
O4	0.3585 (2)	0.42701 (16)	0.30209 (16)	0.0328 (6)
O5	0.6110 (2)	0.20458 (16)	0.10054 (16)	0.0322 (6)
O6	0.4665 (2)	0.31792 (16)	0.03865 (16)	0.0306 (6)
O7	0.6120 (2)	0.07148 (17)	0.23493 (17)	0.0354 (6)
O8	0.3621 (2)	0.15034 (17)	0.06086 (16)	0.0357 (6)
P1	0.55400 (9)	0.30498 (6)	0.11381 (6)	0.0272 (2)
P2	0.69841 (9)	0.43337 (6)	0.17686 (6)	0.0290 (2)
P3	0.47897 (9)	0.36458 (6)	0.27569 (6)	0.0295 (2)

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.051 (3)	0.031 (2)	0.052 (3)	−0.0080 (19)	0.000 (2)	0.0047 (19)
C2	0.051 (3)	0.037 (2)	0.047 (3)	−0.005 (2)	0.001 (2)	−0.0062 (19)
C3	0.032 (2)	0.039 (2)	0.040 (2)	−0.0045 (17)	−0.0041 (19)	0.0002 (18)
C4	0.051 (3)	0.041 (2)	0.045 (3)	−0.018 (2)	0.001 (2)	−0.002 (2)
C5	0.052 (3)	0.038 (2)	0.051 (3)	−0.008 (2)	0.002 (2)	−0.009 (2)
C6	0.036 (2)	0.036 (2)	0.041 (2)	−0.0054 (18)	−0.0062 (19)	−0.0015 (18)
C7	0.035 (2)	0.043 (2)	0.031 (2)	−0.0004 (18)	−0.0019 (18)	−0.0093 (18)
C8	0.034 (2)	0.039 (2)	0.031 (2)	0.0005 (17)	−0.0085 (18)	−0.0037 (17)
C9	0.051 (3)	0.054 (3)	0.036 (3)	0.000 (2)	−0.007 (2)	−0.012 (2)
C10	0.049 (3)	0.072 (3)	0.031 (3)	−0.003 (2)	0.002 (2)	−0.003 (2)
C11	0.039 (2)	0.059 (3)	0.039 (3)	−0.009 (2)	−0.008 (2)	0.013 (2)
C12	0.037 (2)	0.041 (2)	0.037 (2)	−0.0014 (18)	−0.0115 (19)	0.0049 (18)
C13	0.0245 (19)	0.044 (2)	0.027 (2)	0.0016 (17)	−0.0081 (16)	0.0002 (17)
C14	0.029 (2)	0.030 (2)	0.031 (2)	0.0004 (16)	−0.0055 (17)	−0.0085 (16)
C15	0.039 (2)	0.038 (2)	0.035 (2)	0.0017 (18)	−0.0056 (19)	−0.0047 (18)
C16	0.047 (3)	0.046 (3)	0.039 (2)	−0.002 (2)	0.009 (2)	−0.010 (2)
C17	0.035 (2)	0.048 (3)	0.050 (3)	0.007 (2)	0.006 (2)	−0.009 (2)
C18	0.040 (2)	0.035 (2)	0.050 (3)	0.0127 (19)	−0.004 (2)	−0.0047 (19)
C19	0.035 (2)	0.033 (2)	0.034 (2)	0.0050 (17)	−0.0053 (18)	−0.0079 (17)
C20	0.031 (2)	0.034 (2)	0.043 (2)	0.0031 (17)	−0.0073 (19)	−0.0045 (18)
C21	0.043 (2)	0.035 (2)	0.045 (3)	−0.0003 (19)	−0.003 (2)	0.0003 (19)
C22	0.043 (2)	0.036 (2)	0.042 (3)	−0.0013 (19)	−0.004 (2)	0.0021 (18)
C23	0.065 (3)	0.034 (2)	0.057 (3)	0.001 (2)	0.005 (3)	0.000 (2)
C24	0.062 (3)	0.035 (2)	0.057 (3)	−0.005 (2)	0.009 (2)	0.002 (2)
C25	0.053 (3)	0.030 (2)	0.075 (4)	0.000 (2)	0.010 (3)	−0.004 (2)
C26	0.050 (3)	0.029 (2)	0.075 (3)	−0.006 (2)	0.011 (2)	0.005 (2)
C27	0.028 (2)	0.033 (2)	0.032 (2)	0.0014 (16)	−0.0115 (17)	−0.0037 (16)
C28	0.042 (2)	0.035 (2)	0.053 (3)	0.0091 (19)	−0.015 (2)	−0.013 (2)
C29	0.048 (3)	0.057 (3)	0.055 (3)	0.022 (2)	−0.011 (2)	−0.020 (2)
C30	0.032 (2)	0.075 (3)	0.048 (3)	0.015 (2)	0.005 (2)	−0.002 (2)
C31	0.033 (2)	0.050 (3)	0.047 (3)	0.0009 (19)	−0.004 (2)	0.004 (2)
C32	0.027 (2)	0.035 (2)	0.038 (2)	0.0000 (16)	−0.0070 (17)	−0.0032 (17)
C33	0.030 (2)	0.029 (2)	0.035 (2)	−0.0053 (16)	−0.0026 (17)	0.0024 (16)
C34	0.041 (2)	0.044 (3)	0.050 (3)	−0.013 (2)	−0.010 (2)	0.001 (2)
C35	0.073 (3)	0.035 (2)	0.055 (3)	−0.014 (2)	−0.011 (3)	−0.010 (2)
C36	0.083 (4)	0.029 (2)	0.052 (3)	0.006 (2)	0.002 (3)	−0.007 (2)
C37	0.047 (3)	0.032 (2)	0.050 (3)	0.0112 (19)	−0.004 (2)	−0.0016 (19)
C38	0.037 (2)	0.0268 (19)	0.030 (2)	−0.0067 (16)	−0.0049 (17)	0.0031 (16)
C39	0.042 (2)	0.026 (2)	0.032 (2)	0.0009 (17)	−0.0075 (18)	−0.0056 (16)
C40	0.043 (2)	0.041 (2)	0.042 (3)	0.0095 (19)	−0.010 (2)	−0.0032 (19)
C41	0.055 (3)	0.050 (3)	0.048 (3)	−0.003 (2)	−0.025 (2)	−0.001 (2)
C42	0.069 (3)	0.047 (3)	0.032 (2)	0.000 (2)	−0.019 (2)	−0.0078 (19)
C43	0.056 (3)	0.044 (2)	0.032 (2)	0.005 (2)	−0.003 (2)	−0.0062 (18)
C44	0.043 (2)	0.033 (2)	0.029 (2)	0.0008 (18)	−0.0039 (18)	−0.0014 (16)
C45	0.034 (2)	0.045 (2)	0.033 (2)	−0.0004 (18)	0.0013 (18)	−0.0040 (18)

C46	0.048 (3)	0.090 (4)	0.039 (3)	−0.002 (3)	0.010 (2)	0.010 (2)
C47	0.032 (3)	0.120 (5)	0.074 (4)	−0.003 (3)	0.008 (3)	0.020 (3)
C48	0.033 (3)	0.105 (4)	0.071 (4)	−0.008 (3)	−0.011 (3)	0.009 (3)
C49	0.039 (3)	0.067 (3)	0.045 (3)	−0.002 (2)	−0.010 (2)	0.001 (2)
C50	0.030 (2)	0.035 (2)	0.036 (2)	−0.0014 (17)	−0.0028 (18)	−0.0008 (17)
Cl1	0.0586 (7)	0.0497 (7)	0.0504 (7)	0.0054 (5)	−0.0059 (6)	0.0038 (5)
Cl2	0.0525 (6)	0.0362 (6)	0.0536 (7)	−0.0018 (5)	−0.0003 (5)	0.0037 (5)
Cu1	0.0438 (3)	0.0355 (3)	0.0460 (3)	−0.0034 (2)	0.0028 (2)	0.0034 (2)
N1	0.0297 (17)	0.0375 (18)	0.0327 (18)	−0.0091 (14)	0.0003 (14)	−0.0093 (14)
N2	0.0288 (17)	0.0365 (18)	0.0306 (18)	−0.0071 (14)	0.0017 (14)	−0.0060 (14)
N3	0.0347 (18)	0.0342 (18)	0.0373 (19)	−0.0070 (14)	−0.0036 (15)	−0.0132 (14)
N4	0.051 (2)	0.038 (2)	0.065 (3)	0.0100 (18)	0.008 (2)	0.0075 (18)
N5	0.059 (2)	0.040 (2)	0.064 (3)	0.0108 (19)	0.013 (2)	0.0109 (18)
N6	0.078 (3)	0.049 (2)	0.050 (2)	−0.018 (2)	0.012 (2)	−0.0161 (19)
N7	0.077 (3)	0.048 (2)	0.051 (3)	−0.020 (2)	0.011 (2)	−0.0135 (19)
N8	0.044 (2)	0.0320 (19)	0.049 (2)	−0.0043 (16)	0.0027 (17)	0.0017 (16)
N9	0.043 (2)	0.0317 (19)	0.049 (2)	−0.0015 (15)	0.0023 (17)	−0.0002 (16)
O1	0.0271 (13)	0.0271 (13)	0.0404 (16)	−0.0027 (10)	−0.0078 (12)	0.0023 (11)
O2	0.0288 (14)	0.0320 (14)	0.0402 (16)	−0.0006 (11)	−0.0099 (12)	−0.0021 (11)
O3	0.0455 (16)	0.0297 (14)	0.0301 (15)	0.0030 (12)	−0.0061 (12)	−0.0068 (11)
O4	0.0320 (14)	0.0348 (14)	0.0299 (15)	0.0003 (11)	0.0006 (12)	−0.0006 (11)
O5	0.0244 (13)	0.0305 (14)	0.0415 (16)	0.0024 (11)	−0.0046 (12)	−0.0022 (11)
O6	0.0260 (13)	0.0369 (14)	0.0301 (15)	0.0021 (11)	−0.0057 (11)	−0.0066 (11)
O7	0.0359 (15)	0.0322 (14)	0.0367 (16)	0.0044 (12)	−0.0026 (12)	0.0021 (12)
O8	0.0388 (15)	0.0338 (15)	0.0336 (16)	−0.0073 (12)	−0.0004 (12)	−0.0051 (12)
P1	0.0260 (5)	0.0304 (5)	0.0260 (5)	−0.0018 (4)	−0.0041 (4)	−0.0056 (4)
P2	0.0260 (5)	0.0290 (5)	0.0322 (6)	−0.0027 (4)	−0.0044 (4)	−0.0039 (4)
P3	0.0298 (5)	0.0305 (5)	0.0278 (5)	−0.0019 (4)	−0.0002 (4)	−0.0062 (4)

Geometric parameters (Å, °)

C1—N9	1.344 (5)	C30—C31	1.377 (6)
C1—C2	1.364 (6)	C30—H30	0.9300
C1—H1	0.9300	C31—C32	1.400 (5)
C2—C3	1.376 (5)	C31—H31	0.9300
C2—H2	0.9300	C32—C33	1.472 (5)
C3—C4	1.387 (5)	C33—C38	1.384 (5)
C3—C6	1.449 (5)	C33—C34	1.389 (5)
C4—C5	1.370 (6)	C34—C35	1.377 (6)
C4—H4	0.9300	C34—H34	0.9300
C5—N9	1.339 (5)	C35—C36	1.378 (6)
C5—H5	0.9300	C35—H35	0.9300
C6—N7	1.289 (5)	C36—C37	1.384 (6)
C6—O8	1.356 (4)	C36—H36	0.9300
C7—N6	1.299 (5)	C37—C38	1.387 (5)
C7—O8	1.349 (4)	C37—H37	0.9300
C7—C8	1.450 (5)	C38—O2	1.405 (4)
C8—C13	1.394 (5)	C39—C40	1.372 (5)

C8—C9	1.401 (5)	C39—C44	1.382 (5)
C9—C10	1.363 (6)	C39—O3	1.414 (4)
C9—H9	0.9300	C40—C41	1.386 (5)
C10—C11	1.375 (6)	C40—H40	0.9300
C10—H10	0.9300	C41—C42	1.375 (6)
C11—C12	1.383 (6)	C41—H41	0.9300
C11—H11	0.9300	C42—C43	1.368 (6)
C12—C13	1.383 (5)	C42—H42	0.9300
C12—H12	0.9300	C43—C44	1.394 (5)
C13—O6	1.390 (4)	C43—H43	0.9300
C14—C15	1.374 (5)	C44—C45	1.477 (5)
C14—C19	1.387 (5)	C45—C46	1.382 (6)
C14—O5	1.399 (4)	C45—C50	1.387 (5)
C15—C16	1.381 (5)	C46—C47	1.391 (7)
C15—H15	0.9300	C46—H46	0.9300
C16—C17	1.380 (6)	C47—C48	1.363 (7)
C16—H16	0.9300	C47—H47	0.9300
C17—C18	1.363 (6)	C48—C49	1.376 (6)
C17—H17	0.9300	C48—H48	0.9300
C18—C19	1.400 (5)	C49—C50	1.371 (5)
C18—H18	0.9300	C49—H49	0.9300
C19—C20	1.448 (5)	C50—O4	1.405 (4)
C20—N4	1.300 (5)	Cl1—Cu1	2.2619 (14)
C20—O7	1.348 (4)	Cl2—Cu1	2.3008 (15)
C21—N5	1.279 (5)	Cl2—Cu1 ⁱ	2.7006 (15)
C21—O7	1.366 (5)	Cu1—N9	2.021 (3)
C21—C22	1.455 (6)	Cu1—N8 ⁱ	2.045 (4)
C22—C26	1.378 (5)	Cu1—Cl2 ⁱ	2.7006 (15)
C22—C23	1.385 (6)	N1—P1	1.568 (3)
C23—C24	1.368 (6)	N1—P3	1.579 (3)
C23—H23	0.9300	N2—P2	1.575 (3)
C24—N8	1.337 (5)	N2—P1	1.578 (3)
C24—H24	0.9300	N3—P2	1.560 (3)
C25—N8	1.329 (5)	N3—P3	1.573 (3)
C25—C26	1.374 (6)	N4—N5	1.395 (5)
C25—H25	0.9300	N6—N7	1.391 (5)
C26—H26	0.9300	N8—Cu1 ⁱ	2.045 (4)
C27—C28	1.377 (5)	O1—P2	1.587 (2)
C27—C32	1.384 (5)	O2—P2	1.580 (3)
C27—O1	1.395 (4)	O3—P3	1.580 (3)
C28—C29	1.373 (6)	O4—P3	1.584 (3)
C28—H28	0.9300	O5—P1	1.590 (3)
C29—C30	1.365 (6)	O6—P1	1.587 (2)
C29—H29	0.9300		
N9—C1—C2	123.5 (4)	C35—C34—H34	119.4
N9—C1—H1	118.2	C33—C34—H34	119.4
C2—C1—H1	118.2	C34—C35—C36	120.4 (4)

C1—C2—C3	119.3 (4)	C34—C35—H35	119.8
C1—C2—H2	120.4	C36—C35—H35	119.8
C3—C2—H2	120.4	C35—C36—C37	119.7 (4)
C2—C3—C4	118.0 (4)	C35—C36—H36	120.2
C2—C3—C6	119.7 (4)	C37—C36—H36	120.2
C4—C3—C6	122.1 (4)	C36—C37—C38	119.0 (4)
C5—C4—C3	119.1 (4)	C36—C37—H37	120.5
C5—C4—H4	120.5	C38—C37—H37	120.5
C3—C4—H4	120.5	C33—C38—C37	122.2 (3)
N9—C5—C4	123.2 (4)	C33—C38—O2	119.6 (3)
N9—C5—H5	118.4	C37—C38—O2	118.0 (3)
C4—C5—H5	118.4	C40—C39—C44	123.8 (4)
N7—C6—O8	111.7 (4)	C40—C39—O3	117.5 (3)
N7—C6—C3	128.3 (4)	C44—C39—O3	118.6 (3)
O8—C6—C3	119.8 (3)	C39—C40—C41	117.9 (4)
N6—C7—O8	111.7 (4)	C39—C40—H40	121.1
N6—C7—C8	126.8 (4)	C41—C40—H40	121.1
O8—C7—C8	121.4 (3)	C42—C41—C40	120.2 (4)
C13—C8—C9	117.2 (4)	C42—C41—H41	119.9
C13—C8—C7	124.0 (4)	C40—C41—H41	119.9
C9—C8—C7	118.8 (4)	C43—C42—C41	120.3 (4)
C10—C9—C8	121.1 (4)	C43—C42—H42	119.8
C10—C9—H9	119.5	C41—C42—H42	119.8
C8—C9—H9	119.5	C42—C43—C44	121.5 (4)
C9—C10—C11	121.0 (4)	C42—C43—H43	119.3
C9—C10—H10	119.5	C44—C43—H43	119.3
C11—C10—H10	119.5	C39—C44—C43	116.2 (4)
C10—C11—C12	119.4 (4)	C39—C44—C45	121.8 (3)
C10—C11—H11	120.3	C43—C44—C45	122.0 (4)
C12—C11—H11	120.3	C46—C45—C50	116.4 (4)
C13—C12—C11	119.7 (4)	C46—C45—C44	120.6 (4)
C13—C12—H12	120.1	C50—C45—C44	123.0 (4)
C11—C12—H12	120.1	C45—C46—C47	121.2 (5)
C12—C13—O6	118.6 (3)	C45—C46—H46	119.4
C12—C13—C8	121.4 (4)	C47—C46—H46	119.4
O6—C13—C8	120.0 (3)	C48—C47—C46	120.3 (5)
C15—C14—C19	120.9 (3)	C48—C47—H47	119.9
C15—C14—O5	120.8 (3)	C46—C47—H47	119.9
C19—C14—O5	118.0 (3)	C47—C48—C49	120.0 (4)
C14—C15—C16	120.6 (4)	C47—C48—H48	120.0
C14—C15—H15	119.7	C49—C48—H48	120.0
C16—C15—H15	119.7	C50—C49—C48	118.8 (4)
C17—C16—C15	119.5 (4)	C50—C49—H49	120.6
C17—C16—H16	120.3	C48—C49—H49	120.6
C15—C16—H16	120.3	C49—C50—C45	123.2 (4)
C18—C17—C16	119.7 (4)	C49—C50—O4	116.3 (4)
C18—C17—H17	120.2	C45—C50—O4	120.3 (3)
C16—C17—H17	120.2	Cu1—Cl2—Cu1 ⁱ	89.14 (5)

C17—C18—C19	122.1 (4)	N9—Cu1—N8 ⁱ	174.68 (13)
C17—C18—H18	119.0	N9—Cu1—Cl1	88.10 (10)
C19—C18—H18	119.0	N8 ⁱ —Cu1—Cl1	90.93 (10)
C14—C19—C18	117.2 (4)	N9—Cu1—Cl2	88.67 (10)
C14—C19—C20	126.0 (3)	N8 ⁱ —Cu1—Cl2	91.18 (10)
C18—C19—C20	116.8 (3)	Cl1—Cu1—Cl2	167.54 (5)
N4—C20—O7	112.5 (4)	N9—Cu1—Cl2 ⁱ	94.56 (11)
N4—C20—C19	124.0 (4)	N8 ⁱ —Cu1—Cl2 ⁱ	90.77 (10)
O7—C20—C19	123.5 (3)	Cl1—Cu1—Cl2 ⁱ	101.39 (5)
N5—C21—O7	112.5 (4)	Cl2—Cu1—Cl2 ⁱ	90.86 (5)
N5—C21—C22	126.9 (4)	P1—N1—P3	120.96 (19)
O7—C21—C22	120.6 (3)	P2—N2—P1	120.2 (2)
C26—C22—C23	117.5 (4)	P2—N3—P3	121.78 (18)
C26—C22—C21	123.2 (4)	C20—N4—N5	105.9 (3)
C23—C22—C21	119.3 (4)	C21—N5—N4	106.6 (3)
C24—C23—C22	119.2 (4)	C7—N6—N7	106.3 (3)
C24—C23—H23	120.4	C6—N7—N6	106.7 (3)
C22—C23—H23	120.4	C25—N8—C24	116.7 (4)
N8—C24—C23	123.6 (4)	C25—N8—Cu1 ⁱ	124.4 (3)
N8—C24—H24	118.2	C24—N8—Cu1 ⁱ	118.8 (3)
C23—C24—H24	118.2	C5—N9—C1	116.7 (4)
N8—C25—C26	123.5 (4)	C5—N9—Cu1	122.2 (3)
N8—C25—H25	118.2	C1—N9—Cu1	120.5 (3)
C26—C25—H25	118.2	C27—O1—P2	120.5 (2)
C25—C26—C22	119.4 (4)	C38—O2—P2	120.3 (2)
C25—C26—H26	120.3	C39—O3—P3	118.2 (2)
C22—C26—H26	120.3	C50—O4—P3	124.2 (2)
C28—C27—C32	122.7 (4)	C14—O5—P1	128.2 (2)
C28—C27—O1	118.5 (3)	C13—O6—P1	121.9 (2)
C32—C27—O1	118.5 (3)	C20—O7—C21	102.5 (3)
C29—C28—C27	118.7 (4)	C7—O8—C6	103.6 (3)
C29—C28—H28	120.7	N1—P1—N2	118.73 (16)
C27—C28—H28	120.7	N1—P1—O6	104.80 (15)
C30—C29—C28	120.5 (4)	N2—P1—O6	110.24 (15)
C30—C29—H29	119.7	N1—P1—O5	109.29 (16)
C28—C29—H29	119.7	N2—P1—O5	110.41 (15)
C29—C30—C31	120.5 (4)	O6—P1—O5	101.93 (13)
C29—C30—H30	119.7	N3—P2—N2	117.71 (16)
C31—C30—H30	119.7	N3—P2—O2	111.60 (15)
C30—C31—C32	120.6 (4)	N2—P2—O2	105.09 (16)
C30—C31—H31	119.7	N3—P2—O1	105.30 (16)
C32—C31—H31	119.7	N2—P2—O1	112.52 (15)
C27—C32—C31	116.9 (3)	O2—P2—O1	103.87 (13)
C27—C32—C33	121.6 (3)	N3—P3—N1	118.07 (17)
C31—C32—C33	121.3 (3)	N3—P3—O3	112.23 (16)
C38—C33—C34	117.3 (3)	N1—P3—O3	105.70 (16)
C38—C33—C32	121.5 (3)	N3—P3—O4	104.77 (15)
C34—C33—C32	121.2 (3)	N1—P3—O4	112.16 (15)

C35—C34—C33	121.3 (4)	O3—P3—O4	103.00 (14)
N9—C1—C2—C3	−0.6 (7)	C46—C47—C48—C49	−1.9 (9)
C1—C2—C3—C4	3.1 (6)	C47—C48—C49—C50	0.5 (8)
C1—C2—C3—C6	−173.3 (4)	C48—C49—C50—C45	1.4 (7)
C2—C3—C4—C5	−3.7 (6)	C48—C49—C50—O4	−174.4 (4)
C6—C3—C4—C5	172.6 (4)	C46—C45—C50—C49	−1.8 (6)
C3—C4—C5—N9	1.9 (7)	C44—C45—C50—C49	178.0 (4)
C2—C3—C6—N7	2.3 (7)	C46—C45—C50—O4	173.8 (4)
C4—C3—C6—N7	−173.9 (4)	C44—C45—C50—O4	−6.3 (6)
C2—C3—C6—O8	175.9 (4)	Cu1 ⁱ —Cl2—Cu1—N9	94.54 (11)
C4—C3—C6—O8	−0.3 (6)	Cu1 ⁱ —Cl2—Cu1—N8 ⁱ	−90.78 (11)
N6—C7—C8—C13	177.9 (4)	Cu1 ⁱ —Cl2—Cu1—Cl1	169.5 (2)
O8—C7—C8—C13	1.2 (6)	Cu1 ⁱ —Cl2—Cu1—Cl2 ⁱ	0.0
N6—C7—C8—C9	0.2 (6)	O7—C20—N4—N5	−0.4 (5)
O8—C7—C8—C9	−176.5 (3)	C19—C20—N4—N5	−179.7 (4)
C13—C8—C9—C10	−1.0 (6)	O7—C21—N5—N4	0.0 (5)
C7—C8—C9—C10	176.8 (4)	C22—C21—N5—N4	−179.3 (4)
C8—C9—C10—C11	−2.2 (7)	C20—N4—N5—C21	0.2 (5)
C9—C10—C11—C12	2.9 (6)	O8—C7—N6—N7	−1.0 (5)
C10—C11—C12—C13	−0.2 (6)	C8—C7—N6—N7	−178.0 (4)
C11—C12—C13—O6	177.4 (3)	O8—C6—N7—N6	1.0 (5)
C11—C12—C13—C8	−3.2 (6)	C3—C6—N7—N6	175.0 (4)
C9—C8—C13—C12	3.7 (5)	C7—N6—N7—C6	0.0 (5)
C7—C8—C13—C12	−174.0 (3)	C26—C25—N8—C24	1.1 (7)
C9—C8—C13—O6	−176.8 (3)	C26—C25—N8—Cu1 ⁱ	−178.7 (4)
C7—C8—C13—O6	5.5 (5)	C23—C24—N8—C25	−0.1 (7)
C19—C14—C15—C16	−2.9 (6)	C23—C24—N8—Cu1 ⁱ	179.8 (4)
O5—C14—C15—C16	−177.2 (3)	C4—C5—N9—C1	0.7 (6)
C14—C15—C16—C17	1.5 (6)	C4—C5—N9—Cu1	−170.6 (3)
C15—C16—C17—C18	−0.2 (6)	C2—C1—N9—C5	−1.4 (6)
C16—C17—C18—C19	0.2 (6)	C2—C1—N9—Cu1	170.1 (3)
C15—C14—C19—C18	2.8 (5)	N8 ⁱ —Cu1—N9—C5	133.5 (13)
O5—C14—C19—C18	177.3 (3)	Cl1—Cu1—N9—C5	54.0 (3)
C15—C14—C19—C20	−174.4 (3)	Cl2—Cu1—N9—C5	−138.0 (3)
O5—C14—C19—C20	0.1 (5)	Cl2 ⁱ —Cu1—N9—C5	−47.2 (3)
C17—C18—C19—C14	−1.4 (6)	N8 ⁱ —Cu1—N9—C1	−37.5 (15)
C17—C18—C19—C20	176.0 (4)	Cl1—Cu1—N9—C1	−117.0 (3)
C14—C19—C20—N4	165.9 (4)	Cl2—Cu1—N9—C1	51.0 (3)
C18—C19—C20—N4	−11.3 (6)	Cl2 ⁱ —Cu1—N9—C1	141.8 (3)
C14—C19—C20—O7	−13.3 (6)	C28—C27—O1—P2	−113.1 (3)
C18—C19—C20—O7	169.6 (3)	C32—C27—O1—P2	72.5 (4)
N5—C21—C22—C26	147.9 (5)	C33—C38—O2—P2	71.7 (4)
O7—C21—C22—C26	−31.5 (6)	C37—C38—O2—P2	−112.3 (3)
N5—C21—C22—C23	−30.1 (7)	C40—C39—O3—P3	−103.8 (4)
O7—C21—C22—C23	150.6 (4)	C44—C39—O3—P3	77.0 (4)
C26—C22—C23—C24	1.5 (7)	C49—C50—O4—P3	−116.9 (3)
C21—C22—C23—C24	179.5 (4)	C45—C50—O4—P3	67.1 (4)

C22—C23—C24—N8	−1.3 (7)	C15—C14—O5—P1	−49.2 (4)
N8—C25—C26—C22	−0.9 (8)	C19—C14—O5—P1	136.3 (3)
C23—C22—C26—C25	−0.5 (7)	C12—C13—O6—P1	−89.1 (4)
C21—C22—C26—C25	−178.5 (4)	C8—C13—O6—P1	91.4 (4)
C32—C27—C28—C29	−1.4 (6)	N4—C20—O7—C21	0.5 (4)
O1—C27—C28—C29	−175.6 (3)	C19—C20—O7—C21	179.7 (3)
C27—C28—C29—C30	0.1 (6)	N5—C21—O7—C20	−0.3 (5)
C28—C29—C30—C31	0.5 (7)	C22—C21—O7—C20	179.1 (3)
C29—C30—C31—C32	0.2 (7)	N6—C7—O8—C6	1.5 (4)
C28—C27—C32—C31	1.9 (5)	C8—C7—O8—C6	178.7 (3)
O1—C27—C32—C31	176.1 (3)	N7—C6—O8—C7	−1.6 (4)
C28—C27—C32—C33	−173.3 (3)	C3—C6—O8—C7	−176.1 (3)
O1—C27—C32—C33	0.9 (5)	P3—N1—P1—N2	−6.0 (3)
C30—C31—C32—C27	−1.3 (6)	P3—N1—P1—O6	−129.6 (2)
C30—C31—C32—C33	173.9 (4)	P3—N1—P1—O5	121.8 (2)
C27—C32—C33—C38	−47.2 (5)	P2—N2—P1—N1	14.5 (3)
C31—C32—C33—C38	137.8 (4)	P2—N2—P1—O6	135.4 (2)
C27—C32—C33—C34	131.5 (4)	P2—N2—P1—O5	−112.8 (2)
C31—C32—C33—C34	−43.5 (6)	C13—O6—P1—N1	−162.0 (3)
C38—C33—C34—C35	−0.7 (6)	C13—O6—P1—N2	69.2 (3)
C32—C33—C34—C35	−179.4 (4)	C13—O6—P1—O5	−48.1 (3)
C33—C34—C35—C36	−1.1 (7)	C14—O5—P1—N1	−137.8 (3)
C34—C35—C36—C37	0.7 (7)	C14—O5—P1—N2	−5.5 (3)
C35—C36—C37—C38	1.4 (7)	C14—O5—P1—O6	111.7 (3)
C34—C33—C38—C37	2.8 (6)	P3—N3—P2—N2	15.3 (3)
C32—C33—C38—C37	−178.4 (4)	P3—N3—P2—O2	136.9 (2)
C34—C33—C38—O2	178.6 (3)	P3—N3—P2—O1	−111.0 (2)
C32—C33—C38—O2	−2.6 (5)	P1—N2—P2—N3	−18.9 (3)
C36—C37—C38—C33	−3.3 (6)	P1—N2—P2—O2	−143.86 (19)
C36—C37—C38—O2	−179.1 (4)	P1—N2—P2—O1	103.8 (2)
C44—C39—C40—C41	1.7 (6)	C38—O2—P2—N3	71.1 (3)
O3—C39—C40—C41	−177.6 (3)	C38—O2—P2—N2	−160.3 (3)
C39—C40—C41—C42	−0.6 (6)	C38—O2—P2—O1	−41.9 (3)
C40—C41—C42—C43	−0.6 (7)	C27—O1—P2—N3	−165.3 (2)
C41—C42—C43—C44	0.8 (7)	C27—O1—P2—N2	65.3 (3)
C40—C39—C44—C43	−1.5 (6)	C27—O1—P2—O2	−47.8 (3)
O3—C39—C44—C43	177.8 (3)	P2—N3—P3—N1	−7.1 (3)
C40—C39—C44—C45	179.1 (4)	P2—N3—P3—O3	116.2 (2)
O3—C39—C44—C45	−1.7 (6)	P2—N3—P3—O4	−132.7 (2)
C42—C43—C44—C39	0.2 (6)	P1—N1—P3—N3	2.3 (3)
C42—C43—C44—C45	179.7 (4)	P1—N1—P3—O3	−124.2 (2)
C39—C44—C45—C46	138.6 (4)	P1—N1—P3—O4	124.2 (2)
C43—C44—C45—C46	−40.9 (6)	C39—O3—P3—N3	55.3 (3)
C39—C44—C45—C50	−41.3 (6)	C39—O3—P3—N1	−174.7 (3)
C43—C44—C45—C50	139.3 (4)	C39—O3—P3—O4	−56.9 (3)
C50—C45—C46—C47	0.4 (7)	C50—O4—P3—N3	−148.7 (3)

C44—C45—C46—C47	−179.5 (5)	C50—O4—P3—N1	82.0 (3)
C45—C46—C47—C48	1.5 (9)	C50—O4—P3—O3	−31.2 (3)

Symmetry code: (i) $-x, -y, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C26—H26 $\cdots$ O3	0.93	2.38	3.271 (5)	160
C36—H36 $\cdots$ N5 ⁱⁱ	0.93	2.51	3.396 (6)	159

Symmetry code: (ii) $x, y+1, z$.