

Trichloridotris[*N*-[phenyl(pyridin-2-yl)-methylidene]hydroxylamine- κ^2 N,N']-neodymium(III)

Hua Yang

College of Chemistry and Chemical Engineering, Yan'an University, Yan'an, Shaanxi 716000, People's Republic of China

Correspondence e-mail: yanghua7687@163.com


Received 16 March 2012; accepted 31 March 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.022; wR factor = 0.059; data-to-parameter ratio = 17.5.

In the title compound, $[\text{NdCl}_3(\text{C}_{12}\text{H}_{10}\text{N}_2\text{O})_3]$, the central Nd^{III} ion is nine-coordinated by six N atoms from three bidentate chelate *N*-[phenyl(pyridin-2-yl)methylidene]hydroxylamine ligands and three Cl^- ions, and displays a distorted tricapped trigonal prismatic geometry. The complex molecules are stabilized by intramolecular $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonds.

Related literature

For complexes of oximes, see: Kukushkin & Pombeiro (1999); Milios *et al.* (2007); Fritsky *et al.* (2004); Xu *et al.* (2007); Papatriantafyllopoulou *et al.* (2009). For 3d-metal complexes of *N*-[phenyl(pyridine-2-yl)methylidene]hydroxylamine, see: Milios *et al.* (2003); Milios *et al.* (2004). For an Sm complex with this ligand, see: Lei *et al.* (2012).



Experimental

Crystal data

$[\text{NdCl}_3(\text{C}_{12}\text{H}_{10}\text{N}_2\text{O})_3]$
 $M_r = 845.25$
 Triclinic, $P\bar{1}$
 $a = 8.6367$ (17) Å
 $b = 10.460$ (2) Å
 $c = 19.847$ (4) Å
 $\alpha = 91.87$ (3)°
 $\beta = 94.38$ (3)°

$\gamma = 92.80$ (3)°
 $V = 1784.4$ (6) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.72$ mm⁻¹
 $T = 293$ K
 $0.31 \times 0.18 \times 0.13$ mm

Data collection

Bruker SMART CCD-detector
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2000)
 $T_{\min} = 0.617$, $T_{\max} = 0.807$

30524 measured reflections
 7775 independent reflections
 7155 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.022$
 $wR(F^2) = 0.059$
 $S = 1.04$
 7775 reflections

445 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.50$ e Å⁻³
 $\Delta\rho_{\min} = -0.44$ e Å⁻³

Table 1

Selected bond lengths (Å).

Nd1—N2	2.604 (2)	Nd1—N3	2.742 (2)
Nd1—N1	2.661 (2)	Nd1—Cl3	2.7686 (8)
Nd1—N5	2.680 (2)	Nd1—Cl2	2.7903 (9)
Nd1—N4	2.6953 (19)	Nd1—Cl1	2.8296 (10)
Nd1—N6	2.7018 (19)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A $\cdots$ Cl3	0.82	2.22	2.966 (2)	152
O2—H2A $\cdots$ Cl1	0.82	2.19	2.9290 (19)	151
O3—H3A $\cdots$ Cl2	0.82	2.19	2.930 (2)	150

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The author appreciates financial support from Yanan University (grant No. YD2011–20) and the Science and Technology Bureau of Yanan City (grant No. kn2009–16).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2190).

References

- Bruker (2000). *SADABS*, *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fritsky, O., Swiatek-Kozłowska, J., Dobosz, A., Sliva, T. Y. & Dudarenko, N. M. (2004). *Inorg. Chim. Acta*, **357**, 3746–3752.
- Kukushkin, Y. & Pombeiro, A. J. L. (1999). *Coord. Chem. Rev.* **181**, 147–175.
- Lei, T., Chen, W., Chen, Y., Hu, B. & Li, Y. (2012). *Acta Cryst. E* **68**, m344–m345.
- Milios, C. J., Inglis, R., Vinslava, A., Bagai, R., Wernsdorfer, W., Parsons, S., Perlepes, S. P., Christou, G. & Brechin, E. K. (2007). *J. Am. Chem. Soc.* **129**, 12505–12511.
- Milios, C. J., Kefalloniti, E., Raptopoulou, C. P., Terzis, A., Vicente, R., Lalioti, N., Escuer, A. & Perlepes, S. P. (2003). *Chem. Commun.* pp. 819–821.
- Milios, C. J., Stamatatos, T. C., Kyritsis, P., Terzis, A., Raptopoulou, C. P., Vicente, R., Escuer, A. & Perlepes, S. P. (2004). *Eur. J. Inorg. Chem.* pp. 2885–2901.
- Papatriantafyllopoulou, C., Estrader, M., Efthymiou, C. G., Dermitzaki, D., Gkotsis, K., Terzis, A., Diaz, C. & Perlepes, S. P. (2009). *Polyhedron*, **28**, 1652–1655.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xu, H. B., Wang, B. W., Pan, F., Wang, Z. M. & Gao, S. (2007). *Angew. Chem. Int. Ed.* **46**, 7388–7392.

supporting information

Acta Cryst. (2012). E68, m578–m579 [https://doi.org/10.1107/S1600536812014055]

Trichloridotris[*N*-[phenyl(pyridin-2-yl)methylidene]hydroxylamine- $\kappa^2 N, N'$]neodymium(III)

Hua Yang

S1. Comment

The coordination chemistry of oximes (Kukushkin & Pombeiro, 1999; Milios *et al.*, 2007) continues to attract considerable attention, with the efforts of several research groups driven by a number of considerations. These include the use of metal oxime complexes in supramolecular chemistry (Fritsky *et al.*, 2004) and the employment of oximate ligands in the synthesis of complexes with interesting magnetic properties (Xu *et al.*, 2007; Papatriantafyllopoulou *et al.*, 2009; Milios *et al.*, 2007). *N*-[phenyl(pyridine-2-yl)methylidene]hydroxylamine [(py)C(ph)NOH], is one of the oximes that is currently a popular ligand for synthesis of the 3*d*-metal complexes (Milios *et al.*, 2003; Milios *et al.*, 2004). However, the structures of rare earth metal complexes with this ligand are uncommon in the crystallographic literature. Here we report the structure of the neodymium complex with [(py)C(ph)NOH], the title compound [NdCl₃(C₁₂H₁₀N₂O)₃], which was synthesized by the reaction of NdCl₃ · 6H₂O with the ligand under autogenous pressure. The title compound is isomorphous with the Sm^{III} analogue (Lei *et al.*, 2012).

In the title complex, the central Nd^{III} ion is nine-coordinated by six nitrogen atoms from three bidentate chelate ligands and three Cl[−] ions [Nd—N range, 2.604 (2)–2.742 (2) Å; Nd—Cl, 2.7686 (8)–2.8296 (10) Å (Table 1)] and displays a distorted tricapped trigonal prismatic geometry (Fig. 1). The discrete complex molecules are stabilized by intramolecular O—H···Cl hydrogen bonds (Table 2, Fig. 2).

S2. Experimental

A mixture of phenyl-2-pyridyl ketone oxime (0.0198 g, 0.10 mmol), NdCl₃ · 6H₂O (0.0179 g, 0.05 mmol), and ethanol (2 mL) was sealed in a 6 mL Pyrex tube. The tube was heated at 80 °C for 4 days under autogenous pressure. Cooling of the resultant solution to room temperature gave colourless crystals of the product. The crystals were collected by filtration, washed with ethanol (2 mL) and dried in air. Yield: 54%. Anal. Calcd. for C₃₆H₃₀Cl₃N₆NdO₃: C, 51.15; H, 3.58; N, 9.94%. Found: C, 50.93; H, 3.43; N, 9.76%.

S3. Refinement

H atoms were placed in calculated positions and included in the refinement using a riding-model approximation, with C—H = 0.93 Å and O—H = 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.




Figure 1

The molecular structure of the title complex, showing atom labels and 30% probability displacement ellipsoids.




Figure 2

Intramolecular hydrogen-bonding interactions in the title complex, with hydrogen bonds shown as dashed lines.

Trichloridotris{*N*-[phenyl(pyridin-2-yl)methylidene]hydroxylamine- κ^2N,N' }neodymium(III)

Crystal data

[NdCl₃(C₁₂H₁₀N₂O)₃]

$M_r = 845.25$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 8.6367(17) \text{ \AA}$

$b = 10.460(2) \text{ \AA}$

$c = 19.847(4) \text{ \AA}$
 $\alpha = 91.87(3)^\circ$
 $\beta = 94.38(3)^\circ$
 $\gamma = 92.80(3)^\circ$
 $V = 1784.4(6) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 846$
 char

$D_x = 1.573 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 7737 reflections
 $\theta = 2.2\text{--}27.0^\circ$
 $\mu = 1.72 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
 Block, colourless
 $0.31 \times 0.18 \times 0.13 \text{ mm}$

Data collection

Bruker SMART CCD-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2000)
 $T_{\min} = 0.617$, $T_{\max} = 0.807$

30524 measured reflections
 7775 independent reflections
 7155 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$
 $\theta_{\max} = 27.0^\circ$, $\theta_{\min} = 1.0^\circ$
 $h = -11 \rightarrow 11$
 $k = -13 \rightarrow 13$
 $l = -25 \rightarrow 25$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.022$
 $wR(F^2) = 0.059$
 $S = 1.04$
 7775 reflections
 445 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0316P)^2 + 0.3895P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.50 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.44 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Nd1	0.170030 (12)	0.594157 (10)	0.248609 (5)	0.03370 (4)
Cl2	0.10720 (8)	0.35456 (5)	0.30113 (3)	0.05379 (15)
Cl1	0.27975 (7)	0.70288 (6)	0.13140 (3)	0.05048 (14)
Cl3	0.48680 (7)	0.57109 (6)	0.27455 (3)	0.05349 (15)
N2	0.2436 (2)	0.41331 (18)	0.16542 (10)	0.0429 (4)
N6	0.2375 (2)	0.61037 (18)	0.38390 (9)	0.0437 (4)
N1	-0.0455 (2)	0.48814 (17)	0.15808 (9)	0.0397 (4)
N3	-0.0819 (2)	0.65372 (18)	0.31740 (9)	0.0415 (4)
N5	0.2564 (2)	0.81945 (17)	0.31196 (10)	0.0434 (4)

C27	0.3334 (5)	1.0579 (3)	0.37346 (19)	0.0884 (11)
H27	0.3572	1.1381	0.3943	0.106*
N4	−0.0185 (2)	0.77901 (18)	0.20749 (9)	0.0418 (4)
C7	0.2007 (3)	0.2515 (2)	0.07372 (11)	0.0416 (5)
C18	−0.1209 (2)	0.8276 (2)	0.24228 (11)	0.0378 (4)
O1	0.3882 (2)	0.36271 (19)	0.17316 (10)	0.0627 (5)
H1A	0.4419	0.4034	0.2031	0.094*
O2	−0.0065 (2)	0.83670 (19)	0.14624 (9)	0.0592 (5)
H2A	0.0687	0.8104	0.1282	0.089*
C19	−0.2129 (3)	0.9365 (2)	0.21929 (11)	0.0397 (5)
C5	−0.0047 (3)	0.4056 (2)	0.10935 (11)	0.0394 (5)
O3	0.2290 (2)	0.50316 (16)	0.42307 (8)	0.0576 (4)
H3A	0.2028	0.4398	0.3987	0.086*
C17	−0.1480 (2)	0.7670 (2)	0.30694 (11)	0.0384 (5)
C13	−0.1152 (3)	0.5941 (2)	0.37345 (13)	0.0502 (6)
H13	−0.0759	0.5139	0.3803	0.060*
C31	0.2881 (3)	0.7204 (3)	0.49380 (12)	0.0508 (6)
C6	0.1547 (3)	0.3581 (2)	0.11766 (11)	0.0399 (5)
C21	−0.2296 (4)	1.1591 (3)	0.19732 (14)	0.0624 (7)
H21	−0.1853	1.2421	0.1996	0.075*
C1	−0.1928 (3)	0.5235 (2)	0.15376 (12)	0.0459 (5)
H1	−0.2241	0.5772	0.1879	0.055*
C4	−0.1058 (3)	0.3652 (3)	0.05469 (12)	0.0521 (6)
H4	−0.0734	0.3112	0.0210	0.063*
C30	0.2678 (3)	0.7144 (2)	0.41872 (11)	0.0440 (5)
C24	−0.3642 (3)	0.9139 (3)	0.19283 (13)	0.0512 (6)
H24	−0.4110	0.8319	0.1923	0.061*
C14	−0.2040 (3)	0.6440 (3)	0.42155 (13)	0.0569 (6)
H14	−0.2222	0.5994	0.4602	0.068*
C25	0.2728 (3)	0.9250 (2)	0.27624 (14)	0.0537 (6)
H25	0.2582	0.9167	0.2294	0.064*
C20	−0.1452 (3)	1.0601 (2)	0.22234 (13)	0.0516 (6)
H20	−0.0437	1.0759	0.2411	0.062*
C29	0.2820 (3)	0.8323 (2)	0.37946 (12)	0.0475 (5)
C23	−0.4461 (3)	1.0143 (3)	0.16705 (14)	0.0630 (7)
H23	−0.5476	0.9994	0.1482	0.076*
C32	0.4105 (3)	0.6620 (3)	0.52790 (13)	0.0559 (6)
H32	0.4837	0.6224	0.5036	0.067*
C16	−0.2384 (3)	0.8229 (2)	0.35265 (13)	0.0516 (6)
H16	−0.2810	0.9013	0.3441	0.062*
C33	0.4240 (4)	0.6624 (3)	0.59736 (15)	0.0744 (9)
H33	0.5055	0.6222	0.6199	0.089*
C11	0.1861 (4)	0.0299 (3)	0.03858 (14)	0.0635 (7)
H11	0.1448	−0.0530	0.0427	0.076*
C12	0.1387 (3)	0.1293 (2)	0.07905 (13)	0.0549 (6)
H12	0.0648	0.1129	0.1098	0.066*
C2	−0.3004 (3)	0.4845 (3)	0.10144 (14)	0.0554 (6)
H2	−0.4018	0.5110	0.1006	0.067*

C22	−0.3780 (4)	1.1357 (3)	0.16916 (14)	0.0676 (8)
H22	−0.4332	1.2026	0.1513	0.081*
C8	0.3121 (3)	0.2733 (3)	0.02820 (14)	0.0605 (7)
H8	0.3573	0.3552	0.0250	0.073*
C9	0.3564 (4)	0.1743 (3)	−0.01235 (15)	0.0738 (9)
H9	0.4298	0.1897	−0.0434	0.089*
C3	−0.2561 (3)	0.4062 (3)	0.05076 (14)	0.0612 (7)
H3	−0.3258	0.3808	0.0142	0.073*
C15	−0.2652 (3)	0.7611 (3)	0.41137 (14)	0.0597 (7)
H15	−0.3238	0.7982	0.4435	0.072*
C35	0.1970 (5)	0.7814 (4)	0.60097 (17)	0.0867 (11)
H35	0.1257	0.8220	0.6260	0.104*
C26	0.3100 (4)	1.0446 (3)	0.30484 (18)	0.0729 (8)
H26	0.3191	1.1153	0.2780	0.087*
C10	0.2927 (4)	0.0537 (3)	−0.00695 (14)	0.0663 (8)
H10	0.3223	−0.0128	−0.0346	0.080*
C36	0.1811 (4)	0.7812 (3)	0.53085 (15)	0.0708 (8)
H36	0.0992	0.8216	0.5088	0.085*
C34	0.3174 (5)	0.7219 (4)	0.63336 (16)	0.0888 (12)
H34	0.3270	0.7220	0.6803	0.107*
C28	0.3211 (4)	0.9497 (3)	0.41152 (16)	0.0735 (9)
H28	0.3391	0.9562	0.4583	0.088*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Nd1	0.03692 (7)	0.02906 (7)	0.03506 (7)	0.00424 (4)	0.00134 (4)	−0.00019 (4)
Cl2	0.0757 (4)	0.0322 (3)	0.0531 (3)	−0.0011 (3)	0.0043 (3)	0.0023 (2)
Cl1	0.0546 (3)	0.0522 (3)	0.0471 (3)	0.0083 (3)	0.0152 (3)	0.0057 (3)
Cl3	0.0391 (3)	0.0638 (4)	0.0559 (3)	0.0057 (3)	−0.0057 (2)	−0.0067 (3)
N2	0.0381 (9)	0.0402 (10)	0.0505 (11)	0.0122 (8)	0.0013 (8)	−0.0055 (8)
N6	0.0537 (11)	0.0375 (10)	0.0399 (10)	0.0020 (8)	0.0022 (8)	0.0021 (8)
N1	0.0399 (9)	0.0357 (10)	0.0434 (10)	0.0052 (8)	0.0020 (8)	−0.0007 (8)
N3	0.0434 (10)	0.0372 (10)	0.0449 (10)	0.0043 (8)	0.0065 (8)	0.0070 (8)
N5	0.0473 (10)	0.0341 (10)	0.0486 (11)	−0.0012 (8)	0.0056 (8)	−0.0010 (8)
C27	0.132 (3)	0.0428 (16)	0.087 (2)	−0.0277 (18)	0.021 (2)	−0.0182 (16)
N4	0.0471 (10)	0.0395 (10)	0.0406 (10)	0.0087 (8)	0.0080 (8)	0.0085 (8)
C7	0.0463 (12)	0.0391 (12)	0.0395 (11)	0.0083 (10)	0.0026 (9)	−0.0033 (9)
C18	0.0390 (11)	0.0327 (11)	0.0421 (11)	0.0045 (9)	0.0043 (9)	0.0014 (9)
O1	0.0439 (9)	0.0694 (12)	0.0732 (13)	0.0254 (9)	−0.0082 (8)	−0.0251 (10)
O2	0.0709 (12)	0.0669 (12)	0.0464 (9)	0.0307 (10)	0.0216 (8)	0.0219 (9)
C19	0.0465 (12)	0.0367 (11)	0.0378 (11)	0.0112 (9)	0.0092 (9)	0.0038 (9)
C5	0.0432 (11)	0.0347 (11)	0.0400 (11)	0.0033 (9)	0.0012 (9)	0.0004 (9)
O3	0.0852 (13)	0.0419 (9)	0.0445 (9)	0.0005 (9)	−0.0032 (9)	0.0069 (7)
C17	0.0398 (11)	0.0352 (11)	0.0402 (11)	0.0009 (9)	0.0047 (9)	0.0012 (9)
C13	0.0487 (13)	0.0478 (14)	0.0557 (14)	0.0033 (11)	0.0073 (11)	0.0153 (11)
C31	0.0542 (14)	0.0535 (15)	0.0435 (13)	−0.0110 (12)	0.0077 (11)	−0.0074 (11)
C6	0.0438 (12)	0.0349 (11)	0.0412 (11)	0.0060 (9)	0.0044 (9)	−0.0029 (9)

C21	0.087 (2)	0.0373 (13)	0.0674 (17)	0.0171 (13)	0.0259 (16)	0.0089 (12)
C1	0.0426 (12)	0.0432 (13)	0.0523 (13)	0.0081 (10)	0.0033 (10)	−0.0012 (10)
C4	0.0551 (14)	0.0561 (15)	0.0441 (13)	0.0093 (12)	−0.0024 (11)	−0.0088 (11)
C30	0.0434 (12)	0.0454 (13)	0.0424 (12)	−0.0003 (10)	0.0028 (9)	−0.0039 (10)
C24	0.0435 (13)	0.0515 (14)	0.0597 (15)	0.0071 (11)	0.0062 (11)	0.0058 (12)
C14	0.0586 (15)	0.0644 (17)	0.0491 (14)	−0.0020 (13)	0.0124 (12)	0.0132 (12)
C25	0.0626 (15)	0.0399 (13)	0.0585 (15)	−0.0035 (11)	0.0066 (12)	0.0045 (11)
C20	0.0555 (14)	0.0422 (13)	0.0585 (15)	0.0070 (11)	0.0094 (12)	0.0036 (11)
C29	0.0498 (13)	0.0422 (13)	0.0497 (13)	−0.0042 (10)	0.0073 (11)	−0.0068 (10)
C23	0.0500 (14)	0.084 (2)	0.0590 (16)	0.0265 (14)	0.0082 (12)	0.0115 (15)
C32	0.0616 (16)	0.0540 (15)	0.0503 (14)	−0.0100 (12)	0.0015 (12)	−0.0005 (12)
C16	0.0565 (14)	0.0481 (14)	0.0521 (14)	0.0096 (11)	0.0131 (11)	0.0009 (11)
C33	0.086 (2)	0.077 (2)	0.0556 (17)	−0.0239 (18)	−0.0106 (16)	0.0101 (15)
C11	0.089 (2)	0.0368 (13)	0.0653 (17)	0.0058 (13)	0.0113 (15)	−0.0059 (12)
C12	0.0658 (16)	0.0435 (14)	0.0570 (15)	0.0007 (12)	0.0186 (12)	−0.0038 (11)
C2	0.0427 (13)	0.0603 (16)	0.0625 (16)	0.0095 (12)	−0.0041 (11)	0.0003 (13)
C22	0.085 (2)	0.0652 (19)	0.0611 (17)	0.0458 (17)	0.0277 (15)	0.0205 (14)
C8	0.0739 (18)	0.0513 (15)	0.0576 (16)	−0.0027 (13)	0.0204 (14)	−0.0051 (12)
C9	0.084 (2)	0.078 (2)	0.0618 (17)	0.0028 (17)	0.0327 (16)	−0.0137 (15)
C3	0.0541 (15)	0.0714 (18)	0.0550 (15)	0.0049 (13)	−0.0141 (12)	−0.0037 (13)
C15	0.0648 (16)	0.0647 (17)	0.0521 (15)	0.0055 (14)	0.0203 (13)	0.0008 (13)
C35	0.100 (3)	0.098 (3)	0.063 (2)	−0.018 (2)	0.0377 (19)	−0.0240 (18)
C26	0.094 (2)	0.0370 (14)	0.088 (2)	−0.0098 (14)	0.0204 (18)	0.0034 (14)
C10	0.080 (2)	0.0618 (18)	0.0586 (16)	0.0186 (15)	0.0136 (15)	−0.0186 (14)
C36	0.0677 (18)	0.082 (2)	0.0618 (17)	−0.0014 (16)	0.0121 (14)	−0.0123 (15)
C34	0.115 (3)	0.103 (3)	0.0442 (16)	−0.040 (2)	0.0093 (19)	−0.0032 (17)
C28	0.102 (2)	0.0538 (17)	0.0621 (17)	−0.0224 (16)	0.0136 (16)	−0.0154 (14)

Geometric parameters (Å, °)

Nd1—N2	2.604 (2)	C21—H21	0.9300
Nd1—N1	2.661 (2)	C1—C2	1.377 (3)
Nd1—N5	2.680 (2)	C1—H1	0.9300
Nd1—N4	2.6953 (19)	C4—C3	1.384 (4)
Nd1—N6	2.7018 (19)	C4—H4	0.9300
Nd1—N3	2.742 (2)	C30—C29	1.485 (3)
Nd1—C13	2.7686 (8)	C24—C23	1.384 (4)
Nd1—C12	2.7903 (9)	C24—H24	0.9300
Nd1—C11	2.8296 (10)	C14—C15	1.372 (4)
N2—C6	1.277 (3)	C14—H14	0.9300
N2—O1	1.380 (2)	C25—C26	1.370 (4)
N6—C30	1.275 (3)	C25—H25	0.9300
N6—O3	1.387 (2)	C20—H20	0.9300
N1—C1	1.339 (3)	C29—C28	1.380 (4)
N1—C5	1.354 (3)	C23—C22	1.371 (4)
N3—C13	1.336 (3)	C23—H23	0.9300
N3—C17	1.355 (3)	C32—C33	1.374 (4)
N5—C25	1.339 (3)	C32—H32	0.9300

N5—C29	1.342 (3)	C16—C15	1.380 (4)
C27—C26	1.363 (5)	C16—H16	0.9300
C27—C28	1.385 (4)	C33—C34	1.369 (5)
C27—H27	0.9300	C33—H33	0.9300
N4—C18	1.278 (3)	C11—C10	1.358 (4)
N4—O2	1.383 (2)	C11—C12	1.390 (4)
C7—C12	1.372 (3)	C11—H11	0.9300
C7—C8	1.386 (3)	C12—H12	0.9300
C7—C6	1.481 (3)	C2—C3	1.365 (4)
C18—C17	1.479 (3)	C2—H2	0.9300
C18—C19	1.485 (3)	C22—H22	0.9300
O1—H1A	0.8200	C8—C9	1.378 (4)
O2—H2A	0.8200	C8—H8	0.9300
C19—C24	1.377 (3)	C9—C10	1.362 (4)
C19—C20	1.390 (3)	C9—H9	0.9300
C5—C4	1.381 (3)	C3—H3	0.9300
C5—C6	1.485 (3)	C15—H15	0.9300
O3—H3A	0.8200	C35—C34	1.367 (5)
C17—C16	1.376 (3)	C35—C36	1.388 (4)
C13—C14	1.375 (4)	C35—H35	0.9300
C13—H13	0.9300	C26—H26	0.9300
C31—C32	1.389 (4)	C10—H10	0.9300
C31—C36	1.389 (4)	C36—H36	0.9300
C31—C30	1.486 (3)	C34—H34	0.9300
C21—C22	1.366 (4)	C28—H28	0.9300
C21—C20	1.379 (4)		
N2—Nd1—N1	60.38 (6)	C7—C6—C5	120.62 (19)
N2—Nd1—N5	146.60 (6)	C22—C21—C20	120.3 (3)
N1—Nd1—N5	140.98 (6)	C22—C21—H21	119.9
N2—Nd1—N4	121.50 (6)	C20—C21—H21	119.9
N1—Nd1—N4	72.25 (6)	N1—C1—C2	123.5 (2)
N5—Nd1—N4	68.74 (6)	N1—C1—H1	118.3
N2—Nd1—N6	126.82 (6)	C2—C1—H1	118.3
N1—Nd1—N6	139.68 (6)	C5—C4—C3	119.1 (2)
N5—Nd1—N6	59.14 (6)	C5—C4—H4	120.4
N4—Nd1—N6	111.33 (6)	C3—C4—H4	120.4
N2—Nd1—N3	137.44 (6)	N6—C30—C29	115.7 (2)
N1—Nd1—N3	83.49 (6)	N6—C30—C31	123.3 (2)
N5—Nd1—N3	75.89 (6)	C29—C30—C31	121.0 (2)
N4—Nd1—N3	58.57 (6)	C19—C24—C23	119.5 (3)
N6—Nd1—N3	67.56 (6)	C19—C24—H24	120.3
N2—Nd1—Cl3	74.36 (5)	C23—C24—H24	120.3
N1—Nd1—Cl3	134.58 (4)	C15—C14—C13	118.6 (2)
N5—Nd1—Cl3	78.41 (5)	C15—C14—H14	120.7
N4—Nd1—Cl3	136.62 (5)	C13—C14—H14	120.7
N6—Nd1—Cl3	71.65 (5)	N5—C25—C26	123.7 (3)
N3—Nd1—Cl3	138.76 (4)	N5—C25—H25	118.1

N2—Nd1—Cl2	69.77 (5)	C26—C25—H25	118.1
N1—Nd1—Cl2	77.37 (5)	C21—C20—C19	119.4 (3)
N5—Nd1—Cl2	130.24 (4)	C21—C20—H20	120.3
N4—Nd1—Cl2	131.64 (4)	C19—C20—H20	120.3
N6—Nd1—Cl2	71.34 (5)	N5—C29—C28	121.7 (2)
N3—Nd1—Cl2	81.64 (5)	N5—C29—C30	117.4 (2)
Cl3—Nd1—Cl2	91.19 (4)	C28—C29—C30	120.9 (2)
N2—Nd1—Cl1	70.22 (5)	C22—C23—C24	120.2 (3)
N1—Nd1—Cl1	81.67 (5)	C22—C23—H23	119.9
N5—Nd1—Cl1	86.33 (5)	C24—C23—H23	119.9
N4—Nd1—Cl1	70.85 (5)	C33—C32—C31	120.5 (3)
N6—Nd1—Cl1	138.42 (5)	C33—C32—H32	119.8
N3—Nd1—Cl1	129.42 (4)	C31—C32—H32	119.8
Cl3—Nd1—Cl1	79.77 (3)	C17—C16—C15	119.1 (2)
Cl2—Nd1—Cl1	139.95 (3)	C17—C16—H16	120.5
C6—N2—O1	113.33 (18)	C15—C16—H16	120.5
C6—N2—Nd1	126.43 (14)	C34—C33—C32	119.9 (3)
O1—N2—Nd1	120.13 (13)	C34—C33—H33	120.0
C30—N6—O3	113.23 (18)	C32—C33—H33	120.0
C30—N6—Nd1	125.00 (15)	C10—C11—C12	120.1 (3)
O3—N6—Nd1	121.56 (13)	C10—C11—H11	120.0
C1—N1—C5	117.13 (19)	C12—C11—H11	120.0
C1—N1—Nd1	122.39 (15)	C7—C12—C11	120.2 (2)
C5—N1—Nd1	120.07 (14)	C7—C12—H12	119.9
C13—N3—C17	116.6 (2)	C11—C12—H12	119.9
C13—N3—Nd1	121.58 (15)	C3—C2—C1	119.0 (2)
C17—N3—Nd1	119.27 (14)	C3—C2—H2	120.5
C25—N5—C29	117.5 (2)	C1—C2—H2	120.5
C25—N5—Nd1	120.04 (16)	C21—C22—C23	120.4 (2)
C29—N5—Nd1	122.42 (15)	C21—C22—H22	119.8
C26—C27—C28	118.7 (3)	C23—C22—H22	119.8
C26—C27—H27	120.6	C9—C8—C7	120.3 (3)
C28—C27—H27	120.6	C9—C8—H8	119.8
C18—N4—O2	112.71 (17)	C7—C8—H8	119.8
C18—N4—Nd1	125.20 (14)	C10—C9—C8	120.1 (3)
O2—N4—Nd1	122.00 (12)	C10—C9—H9	120.0
C12—C7—C8	118.9 (2)	C8—C9—H9	120.0
C12—C7—C6	120.9 (2)	C2—C3—C4	119.0 (2)
C8—C7—C6	120.1 (2)	C2—C3—H3	120.5
N4—C18—C17	116.40 (18)	C4—C3—H3	120.5
N4—C18—C19	122.86 (19)	C14—C15—C16	118.9 (2)
C17—C18—C19	120.71 (18)	C14—C15—H15	120.6
N2—O1—H1A	109.5	C16—C15—H15	120.6
N4—O2—H2A	109.5	C34—C35—C36	120.1 (3)
C24—C19—C20	120.1 (2)	C34—C35—H35	120.0
C24—C19—C18	119.6 (2)	C36—C35—H35	120.0
C20—C19—C18	120.2 (2)	C27—C26—C25	118.7 (3)
N1—C5—C4	122.2 (2)	C27—C26—H26	120.6

N1—C5—C6	116.83 (19)	C25—C26—H26	120.6
C4—C5—C6	120.9 (2)	C11—C10—C9	120.5 (3)
N6—O3—H3A	109.5	C11—C10—H10	119.8
N3—C17—C16	122.7 (2)	C9—C10—H10	119.8
N3—C17—C18	116.38 (19)	C35—C36—C31	119.7 (3)
C16—C17—C18	120.9 (2)	C35—C36—H36	120.2
N3—C13—C14	124.0 (2)	C31—C36—H36	120.2
N3—C13—H13	118.0	C35—C34—C33	120.7 (3)
C14—C13—H13	118.0	C35—C34—H34	119.7
C32—C31—C36	119.1 (2)	C33—C34—H34	119.7
C32—C31—C30	120.8 (2)	C29—C28—C27	119.5 (3)
C36—C31—C30	120.1 (3)	C29—C28—H28	120.2
N2—C6—C7	124.1 (2)	C27—C28—H28	120.2
N2—C6—C5	115.26 (19)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O1—H1A $\cdots$ Cl3	0.82	2.22	2.966 (2)	152
O2—H2A $\cdots$ Cl1	0.82	2.19	2.9290 (19)	151
O3—H3A $\cdots$ Cl2	0.82	2.19	2.930 (2)	150