

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

# catena-Poly[[*(benzil bis[(pyridin-2-yl)-methylidene]hydrazone)-κ<sup>4</sup>N,N',N'',N'''*)-mercury(II)]-μ-chlorido-[dichlorido-mercury(II)]-μ-chlorido]

Mehmet Akkurt,<sup>a\*</sup> Ali Akbar Khandar,<sup>b</sup> Muhammad Nawaz Tahir,<sup>c</sup> Seyed Abolfazl Hosseini-Yazdi<sup>b</sup> and Ghodrat Mahmoudi<sup>b</sup>

<sup>a</sup>Department of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, <sup>b</sup>Department of Inorganic Chemistry, Faculty of Chemistry, University of Tabriz, P.O. Box 51666, Tabriz, Iran, and <sup>c</sup>Department of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: akkurt@erciyes.edu.tr


Received 10 May 2012; accepted 6 June 2012

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.049; data-to-parameter ratio = 20.6.

In the title coordination polymer,  $[\text{Hg}_2\text{Cl}_4(\text{C}_{26}\text{H}_{20}\text{N}_6)]_n$ , one  $\text{Hg}^{\text{II}}$  ion is coordinated by four N atoms from the benzylbis((pyridin-2-yl)methylidene)hydrazone ligand and two  $\text{Cl}^-$  ions in a very distorted *cis*- $\text{HgCl}_2\text{N}_4$  octahedral geometry. The other  $\text{Hg}^{\text{II}}$  ion is coordinated in a distorted tetrahedral geometry by four  $\text{Cl}^-$  ions. Bridging chloride ions link the  $\text{Hg}^{\text{II}}$  ions into a chain propagating in [010]: the  $\text{Hg}-\text{Cl}$  bridging bonds are significantly longer than the terminal bonds. The dihedral angle between the central benzene rings of the ligand is  $83.3(2)^\circ$ . The packing is consolidated by weak  $\text{C}-\text{H}\cdots\text{Cl}$  hydrogen bonds and  $\text{C}-\text{H}\cdots\pi$  interactions.

## Related literature

For background to polyimine ligands, see: Bai *et al.* (2005); Chowdhury *et al.* (2003); Drew *et al.* (2006); Pal *et al.* (2000); Sun *et al.* (2006).



## Experimental

### Crystal data

$[\text{HgCl}_4(\text{C}_{26}\text{H}_{20}\text{N}_6)]$   
 $M_r = 959.46$   
 Monoclinic,  $P2_1/c$   
 $a = 8.8560(2)$  Å  
 $b = 13.8093(4)$  Å  
 $c = 23.6960(7)$  Å  
 $\beta = 96.702(1)^\circ$

$V = 2878.10(14)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 11.06$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.36 \times 0.18 \times 0.16$  mm

### Data collection

Bruker Kappa APEXII CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 2005)  
 $T_{\text{min}} = 0.106$ ,  $T_{\text{max}} = 0.171$

26994 measured reflections  
 7066 independent reflections  
 5295 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.049$   
 $S = 1.02$   
 7066 reflections

343 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.85$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.85$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Hg1—Cl1	2.5560 (10)	Hg1—N6	2.435 (3)
Hg1—Cl2	2.5488 (11)	Hg2—Cl1	2.7777 (10)
Hg1—N2	2.471 (3)	Hg2—Cl3	2.3328 (12)
Hg1—N3	2.499 (3)	Hg2—Cl4	2.3308 (12)
Hg1—N5	2.558 (3)	Hg2—Cl2 <sup>i</sup>	2.7227 (11)

Symmetry code: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

*Cg*5 is the centroid of the C1–C6 phenyl ring.

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C14—H14 $\cdots$ Cl3 <sup>iii</sup>	0.93	2.82	3.647 (4)	149
C25—H25 $\cdots$ <i>Cg</i> 5 <sup>iii</sup>	0.93	2.96	3.814 (5)	153

Symmetry codes: (ii)  $-x + 1, -y + 2, -z + 1$ ; (iii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of a diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6787).

## References

- Bai, Y., Duan, C. Y., Cai, P., Dang, D. B. & Meng, Q. J. (2005). *Dalton Trans.* pp. 2678–2680.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chowdhury, S., Iveson, P. B., Drew, M. G. B., Tocher, D. A. & Datta, D. (2003). *New J. Chem.* **27**, 193–196.
- Drew, M. G. B., Parui, D., Naskar, J. P. & Datta, D. (2006). *Eur. J. Inorg. Chem.* pp. 4026–4028.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Pal, P. K., Chowdhury, S., Drew, M. G. B. & Datta, D. (2000). *New J. Chem.* **24**, 931–933.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sun, Q. Z., Bai, Y., He, G. J., Duan, C. Y., Lin, Z. H. & Meng, Q. J. (2006). *Chem. Commun.* pp. 2777–2779.

## supporting information

*Acta Cryst.* (2012). E68, m903–m904 [https://doi.org/10.1107/S1600536812025718]

***catena*-Poly[[[(benzil bis{[(pyridin-2-yl)methylidene]hydrazone}- $\kappa^4N,N',N'',N'''$ )mercury(II)]- $\mu$ -chlorido-[dichloridomercury(II)]- $\mu$ -chlorido]**

**Mehmet Akkurt, Ali Akbar Khandar, Muhammad Nawaz Tahir, Seyed Abolfazl Hosseini-Yazdi and Ghodrat Mahmoudi**

### S1. Comment

Research of polyimine compound is well established currently of great interest because of their potential applications as useful organic ligands, in which the amine nitrogen atoms have strong coordination ability to transition metal ions and recognition function (Bai *et al.*, 2005; Pal *et al.*, 2000; Chowdhury *et al.*, 2003; Drew *et al.*, 2006; Sun *et al.*, 2006). In this paper, we report the synthesis and crystal structure of the title compound.


The molecular structure of the title bimetallic coordination polymer  $\text{Cl}_2\text{Hg}-\text{Cl}-(\text{C}_{26}\text{H}_{20}\text{N}_6)\text{Hg}-\text{Cl}-\text{HgCl}_2$  (I) is shown in Fig. 1. Atom Hg2 is four-coordinated in a distorted tetrahedral coordination geometry by two bridging Cl atoms and two terminal Cl atoms. The bond distances of Hg—N are in the range of 2.435 (3) – 2.5558 (3) Å, the bond distances of Hg—Cl are in the range of 2.3308 (12) – 2.7777 (10) Å. *N',N'*-bis[1-(pyridin-2-yl)methylidene]benzil dihydrazone acts as cleating ligand here. The packing and hydrogen bonding (Table 1) is shown in Fig. 2. In addition, a C—H $\cdots$  $\pi$  interaction contributes to the stabilization of the crystal packing.

### S2. Experimental

Benzilbis((pyridin-2-yl)methylidenehydrazone) (*L*) was readily prepared by the reaction of benzil dihydrazone with 2-pyridinecarboxaldehyde in a 1:2 ratio. The ligand *L* (0.5 mmol, 0.208 g) and  $\text{HgCl}_2$  (0.5 mmol, 0.135 g) were mixed in methanol (40 ml). The solution was left for 10 d at room temperature to afford green prisms (yield 75%). Analysis calculated for  $\text{C}_{26}\text{H}_{20}\text{Cl}_4\text{Hg}_2\text{N}_6$ : C 66.2, H 4.0, N 11.0%; found: C 66.3, H 4.1, N 10.9%.

### S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .



**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.




Figure 2

The packing and hydrogen bonding (dashed lines) of (I) viewing down  $a$  axis. H atoms not involved in hydrogen bonding are omitted.

**catena-Poly[[bis(benzil bis((pyridin-2-yl)methylidene)hydrazone)- $\kappa^4N,N',N'',N'''$ ]mercury(II)]- $\mu$ -chlorido-[dichloridomercury(II)]- $\mu$ -chlorido]**

*Crystal data*

[HgCl<sub>4</sub>(C<sub>26</sub>H<sub>20</sub>N<sub>6</sub>)]

$M_r = 959.46$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.8560$  (2) Å

$b = 13.8093$  (4) Å

$c = 23.6960$  (7) Å

$\beta = 96.702$  (1)°

$V = 2878.10$  (14) Å<sup>3</sup>

$Z = 4$

$F(000) = 1784$

$D_x = 2.214$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 700 reflections

$\theta = 3.5$ – $20.5$ °

$\mu = 11.06$  mm<sup>-1</sup>

$T = 296$  K

Prism, green

$0.36 \times 0.18 \times 0.16$  mm

*Data collection*

Bruker Kappa APEXII CCD diffractometer	26994 measured reflections
Radiation source: fine-focus sealed tube	7066 independent reflections
Graphite monochromator	5295 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.035$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$\theta_{\text{max}} = 28.3^\circ$ , $\theta_{\text{min}} = 2.3^\circ$
$T_{\text{min}} = 0.106$ , $T_{\text{max}} = 0.171$	$h = -11 \rightarrow 11$
	$k = -18 \rightarrow 18$
	$l = -31 \rightarrow 31$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.026$	H-atom parameters constrained
$wR(F^2) = 0.049$	$w = 1/[\sigma^2(F_o^2) + (0.0162P)^2 + 0.1878P]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
7066 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
343 parameters	$\Delta\rho_{\text{max}} = 0.85 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.85 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement on  $F^2$  for ALL reflections except those flagged by the user for potential systematic errors. Weighted  $R$ -factors  $wR$  and all goodnesses of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The observed criterion of  $F^2 > \sigma(F^2)$  is used only for calculating  $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.62651 (2)	0.77072 (1)	0.34093 (1)	0.0365 (1)
Hg2	0.40107 (2)	1.03230 (1)	0.29193 (1)	0.0478 (1)
Cl1	0.68420 (11)	0.94448 (7)	0.31308 (5)	0.0508 (4)
Cl2	0.47008 (11)	0.68234 (8)	0.25964 (5)	0.0501 (3)
Cl3	0.37878 (15)	1.09629 (9)	0.38169 (5)	0.0692 (5)
Cl4	0.29001 (14)	0.94051 (8)	0.21557 (5)	0.0598 (4)
N1	0.7872 (3)	0.8179 (2)	0.47703 (13)	0.0386 (11)
N2	0.6483 (4)	0.8111 (2)	0.44313 (12)	0.0361 (10)
N3	0.3796 (3)	0.8157 (2)	0.37525 (13)	0.0376 (11)
N4	0.7524 (3)	0.5949 (2)	0.44393 (12)	0.0382 (11)
N5	0.7914 (4)	0.6422 (2)	0.39605 (12)	0.0370 (10)
N6	0.8447 (3)	0.7191 (2)	0.29480 (13)	0.0395 (11)
C1	0.9891 (4)	0.7451 (3)	0.53615 (14)	0.0346 (11)
C2	1.0814 (4)	0.6645 (3)	0.54636 (16)	0.0443 (14)
C3	1.2180 (5)	0.6703 (4)	0.58042 (18)	0.0551 (17)
C4	1.2655 (5)	0.7574 (4)	0.60399 (18)	0.0572 (18)

C5	1.1760 (5)	0.8389 (3)	0.59423 (16)	0.0518 (16)
C6	1.0372 (4)	0.8329 (3)	0.56081 (15)	0.0431 (14)
C7	0.8431 (4)	0.7388 (3)	0.49870 (14)	0.0316 (11)
C8	0.7681 (4)	0.6415 (3)	0.49102 (15)	0.0331 (11)
C9	0.7156 (4)	0.5941 (3)	0.54105 (15)	0.0351 (12)
C10	0.6896 (5)	0.4958 (3)	0.54188 (18)	0.0520 (17)
C11	0.6362 (5)	0.4532 (3)	0.5883 (2)	0.0650 (19)
C12	0.6057 (5)	0.5077 (4)	0.6335 (2)	0.0641 (19)
C13	0.6267 (5)	0.6055 (4)	0.63283 (17)	0.0541 (18)
C14	0.6849 (4)	0.6493 (3)	0.58726 (16)	0.0436 (14)
C15	0.5333 (5)	0.8437 (3)	0.46300 (16)	0.0450 (16)
C16	0.3852 (4)	0.8468 (3)	0.42897 (15)	0.0392 (12)
C17	0.2613 (5)	0.8829 (3)	0.45179 (19)	0.0579 (17)
C18	0.1243 (5)	0.8889 (3)	0.4169 (2)	0.0631 (19)
C19	0.1184 (5)	0.8602 (3)	0.3623 (2)	0.0598 (19)
C20	0.2472 (5)	0.8220 (3)	0.34252 (18)	0.0502 (17)
C21	0.8909 (4)	0.6034 (3)	0.36920 (16)	0.0417 (12)
C22	0.9269 (4)	0.6436 (3)	0.31518 (15)	0.0362 (12)
C23	1.0385 (4)	0.6025 (3)	0.28678 (18)	0.0510 (16)
C24	1.0661 (5)	0.6424 (4)	0.23574 (19)	0.0589 (19)
C25	0.9829 (5)	0.7200 (3)	0.21479 (18)	0.0569 (18)
C26	0.8728 (5)	0.7570 (3)	0.24518 (17)	0.0516 (16)
H2	1.05060	0.60550	0.52990	0.0530*
H3	1.27800	0.61530	0.58740	0.0660*
H4	1.35840	0.76160	0.62660	0.0690*
H5	1.20890	0.89790	0.61010	0.0620*
H6	0.97610	0.88760	0.55490	0.0520*
H10	0.70800	0.45790	0.51100	0.0620*
H11	0.62100	0.38660	0.58880	0.0780*
H12	0.57050	0.47830	0.66480	0.0770*
H13	0.60190	0.64300	0.66300	0.0650*
H14	0.70330	0.71560	0.58770	0.0520*
H15	0.54270	0.86640	0.50020	0.0540*
H17	0.26880	0.90280	0.48950	0.0690*
H18	0.03770	0.91240	0.43100	0.0750*
H19	0.02840	0.86610	0.33800	0.0720*
H20	0.24080	0.80000	0.30520	0.0600*
H21	0.94200	0.54850	0.38400	0.0500*
H23	1.09360	0.54930	0.30180	0.0610*
H24	1.14100	0.61660	0.21570	0.0710*
H25	1.00020	0.74760	0.18030	0.0690*
H26	0.81620	0.81000	0.23070	0.0620*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Hg1	0.0406 (1)	0.0377 (1)	0.0319 (1)	0.0028 (1)	0.0073 (1)	-0.0034 (1)
Hg2	0.0651 (1)	0.0414 (1)	0.0374 (1)	0.0024 (1)	0.0082 (1)	-0.0028 (1)

C11	0.0480 (6)	0.0347 (6)	0.0703 (7)	-0.0031 (5)	0.0096 (5)	0.0021 (5)
C12	0.0486 (6)	0.0474 (6)	0.0531 (6)	0.0031 (5)	0.0014 (5)	-0.0196 (5)
C13	0.1034 (10)	0.0603 (8)	0.0489 (7)	-0.0135 (7)	0.0297 (7)	-0.0188 (6)
C14	0.0793 (8)	0.0545 (7)	0.0430 (6)	-0.0037 (6)	-0.0032 (6)	-0.0048 (5)
N1	0.0452 (19)	0.037 (2)	0.0329 (17)	-0.0006 (16)	0.0013 (15)	-0.0003 (15)
N2	0.0432 (19)	0.0331 (18)	0.0310 (17)	0.0027 (15)	0.0001 (15)	-0.0007 (14)
N3	0.0425 (19)	0.0355 (19)	0.0342 (18)	-0.0027 (15)	0.0019 (15)	0.0019 (14)
N4	0.0486 (19)	0.0345 (19)	0.0313 (17)	-0.0030 (15)	0.0041 (15)	-0.0012 (14)
N5	0.0472 (19)	0.0383 (19)	0.0258 (16)	-0.0045 (15)	0.0051 (14)	-0.0060 (14)
N6	0.0398 (18)	0.047 (2)	0.0325 (17)	0.0084 (16)	0.0071 (14)	-0.0015 (15)
C1	0.040 (2)	0.042 (2)	0.0216 (18)	-0.0065 (18)	0.0033 (16)	0.0001 (16)
C2	0.044 (2)	0.047 (3)	0.040 (2)	-0.004 (2)	-0.0032 (19)	-0.0044 (19)
C3	0.044 (3)	0.064 (3)	0.055 (3)	0.001 (2)	-0.004 (2)	-0.002 (2)
C4	0.052 (3)	0.081 (4)	0.037 (2)	-0.015 (3)	-0.002 (2)	0.001 (2)
C5	0.065 (3)	0.057 (3)	0.034 (2)	-0.029 (2)	0.009 (2)	-0.011 (2)
C6	0.048 (2)	0.046 (3)	0.035 (2)	-0.008 (2)	0.0037 (19)	-0.0015 (19)
C7	0.0364 (19)	0.032 (2)	0.0267 (18)	-0.0028 (17)	0.0056 (16)	0.0003 (16)
C8	0.0323 (19)	0.032 (2)	0.034 (2)	0.0006 (16)	0.0002 (16)	0.0004 (17)
C9	0.034 (2)	0.036 (2)	0.035 (2)	-0.0006 (17)	0.0023 (16)	0.0039 (17)
C10	0.069 (3)	0.042 (3)	0.047 (3)	-0.005 (2)	0.015 (2)	0.003 (2)
C11	0.072 (3)	0.047 (3)	0.079 (4)	-0.005 (2)	0.021 (3)	0.022 (3)
C12	0.056 (3)	0.088 (4)	0.050 (3)	-0.001 (3)	0.013 (2)	0.031 (3)
C13	0.048 (3)	0.081 (4)	0.035 (2)	0.001 (2)	0.012 (2)	0.005 (2)
C14	0.045 (2)	0.047 (3)	0.039 (2)	-0.0078 (19)	0.0054 (19)	-0.0036 (19)
C15	0.058 (3)	0.047 (3)	0.031 (2)	0.005 (2)	0.009 (2)	-0.0040 (19)
C16	0.045 (2)	0.045 (2)	0.029 (2)	-0.0008 (19)	0.0098 (18)	0.0050 (18)
C17	0.054 (3)	0.076 (3)	0.048 (3)	0.004 (2)	0.024 (2)	0.003 (2)
C18	0.045 (3)	0.065 (3)	0.083 (4)	-0.002 (2)	0.023 (3)	0.004 (3)
C19	0.041 (3)	0.052 (3)	0.083 (4)	-0.010 (2)	-0.007 (3)	0.004 (3)
C20	0.057 (3)	0.040 (3)	0.052 (3)	-0.011 (2)	0.000 (2)	-0.001 (2)
C21	0.045 (2)	0.039 (2)	0.040 (2)	0.0019 (19)	0.0000 (19)	0.0002 (19)
C22	0.035 (2)	0.038 (2)	0.035 (2)	-0.0014 (17)	0.0018 (17)	-0.0053 (17)
C23	0.042 (2)	0.058 (3)	0.054 (3)	0.016 (2)	0.010 (2)	-0.001 (2)
C24	0.048 (3)	0.078 (4)	0.054 (3)	0.005 (2)	0.020 (2)	-0.012 (3)
C25	0.061 (3)	0.077 (4)	0.035 (2)	0.005 (3)	0.015 (2)	0.001 (2)
C26	0.053 (3)	0.067 (3)	0.035 (2)	0.009 (2)	0.006 (2)	0.005 (2)

*Geometric parameters (Å, °)*

Hg1—C11	2.5560 (10)	C12—C13	1.364 (8)
Hg1—C12	2.5488 (11)	C13—C14	1.388 (6)
Hg1—N2	2.471 (3)	C15—C16	1.458 (6)
Hg1—N3	2.499 (3)	C16—C17	1.372 (6)
Hg1—N5	2.558 (3)	C17—C18	1.389 (6)
Hg1—N6	2.435 (3)	C18—C19	1.348 (7)
Hg2—C11	2.7777 (10)	C19—C20	1.387 (6)
Hg2—C13	2.3328 (12)	C21—C22	1.464 (5)
Hg2—C14	2.3308 (12)	C22—C23	1.381 (5)



Hg2—C12 <sup>i</sup>	2.7227 (11)	C23—C24	1.377 (6)
N1—N2	1.392 (4)	C24—C25	1.361 (7)
N1—C7	1.282 (5)	C25—C26	1.376 (6)
N2—C15	1.254 (5)	C2—H2	0.9300
N3—C16	1.339 (5)	C3—H3	0.9300
N3—C20	1.331 (5)	C4—H4	0.9300
N4—N5	1.387 (4)	C5—H5	0.9300
N4—C8	1.282 (5)	C6—H6	0.9300
N5—C21	1.265 (5)	C10—H10	0.9300
N6—C22	1.330 (5)	C11—H11	0.9300
N6—C26	1.337 (5)	C12—H12	0.9300
C1—C2	1.385 (6)	C13—H13	0.9300
C1—C6	1.391 (6)	C14—H14	0.9300
C1—C7	1.483 (5)	C15—H15	0.9300
C2—C3	1.376 (6)	C17—H17	0.9300
C3—C4	1.371 (7)	C18—H18	0.9300
C4—C5	1.380 (7)	C19—H19	0.9300
C5—C6	1.385 (6)	C20—H20	0.9300
C7—C8	1.501 (6)	C21—H21	0.9300
C8—C9	1.476 (5)	C23—H23	0.9300
C9—C10	1.377 (6)	C24—H24	0.9300
C9—C14	1.387 (5)	C25—H25	0.9300
C10—C11	1.379 (6)	C26—H26	0.9300
C11—C12	1.362 (7)		
C11—Hg1—C12	111.21 (4)	C9—C14—C13	120.0 (4)
C11—Hg1—N2	92.64 (7)	N2—C15—C16	121.4 (3)
C11—Hg1—N3	93.41 (7)	N3—C16—C15	116.6 (3)
C11—Hg1—N5	131.59 (8)	N3—C16—C17	123.3 (3)
C11—Hg1—N6	88.05 (7)	C15—C16—C17	120.1 (4)
C12—Hg1—N2	145.31 (8)	C16—C17—C18	118.1 (4)
C12—Hg1—N3	86.90 (7)	C17—C18—C19	119.2 (4)
C12—Hg1—N5	106.34 (7)	C18—C19—C20	119.7 (4)
C12—Hg1—N6	84.86 (7)	N3—C20—C19	122.1 (4)
N2—Hg1—N3	66.18 (11)	N5—C21—C22	121.0 (4)
N2—Hg1—N5	71.43 (9)	N6—C22—C21	116.7 (3)
N2—Hg1—N6	122.06 (11)	N6—C22—C23	122.5 (3)
N3—Hg1—N5	118.39 (10)	C21—C22—C23	120.8 (4)
N3—Hg1—N6	171.59 (10)	C22—C23—C24	118.3 (4)
N5—Hg1—N6	65.83 (10)	C23—C24—C25	119.4 (4)
C11—Hg2—C13	99.87 (4)	C24—C25—C26	119.2 (4)
C11—Hg2—C14	101.12 (4)	N6—C26—C25	122.1 (4)
C11—Hg2—C12 <sup>i</sup>	89.80 (3)	C1—C2—H2	119.00
C13—Hg2—C14	147.23 (5)	C3—C2—H2	119.00
C12 <sup>i</sup> —Hg2—C13	101.86 (4)	C2—C3—H3	120.00
C12 <sup>i</sup> —Hg2—C14	103.09 (4)	C4—C3—H3	120.00
Hg1—C11—Hg2	104.66 (3)	C3—C4—H4	120.00
Hg1—C12—Hg2 <sup>ii</sup>	118.96 (4)	C5—C4—H4	120.00

N2—N1—C7	116.6 (3)	C4—C5—H5	120.00
Hg1—N2—N1	122.9 (2)	C6—C5—H5	120.00
Hg1—N2—C15	118.4 (3)	C1—C6—H6	120.00
N1—N2—C15	117.4 (3)	C5—C6—H6	120.00
Hg1—N3—C16	116.9 (2)	C9—C10—H10	120.00
Hg1—N3—C20	124.9 (3)	C11—C10—H10	120.00
C16—N3—C20	117.7 (3)	C10—C11—H11	120.00
N5—N4—C8	117.6 (3)	C12—C11—H11	120.00
Hg1—N5—N4	124.2 (2)	C11—C12—H12	120.00
Hg1—N5—C21	115.3 (2)	C13—C12—H12	120.00
N4—N5—C21	117.8 (3)	C12—C13—H13	120.00
Hg1—N6—C22	119.7 (2)	C14—C13—H13	120.00
Hg1—N6—C26	121.3 (2)	C9—C14—H14	120.00
C22—N6—C26	118.4 (3)	C13—C14—H14	120.00
C2—C1—C6	118.8 (3)	N2—C15—H15	119.00
C2—C1—C7	120.9 (4)	C16—C15—H15	119.00
C6—C1—C7	120.4 (4)	C16—C17—H17	121.00
C1—C2—C3	121.1 (4)	C18—C17—H17	121.00
C2—C3—C4	119.8 (5)	C17—C18—H18	120.00
C3—C4—C5	120.3 (4)	C19—C18—H18	120.00
C4—C5—C6	120.1 (4)	C18—C19—H19	120.00
C1—C6—C5	120.0 (4)	C20—C19—H19	120.00
N1—C7—C1	117.3 (3)	N3—C20—H20	119.00
N1—C7—C8	124.7 (3)	C19—C20—H20	119.00
C1—C7—C8	117.9 (3)	N5—C21—H21	120.00
N4—C8—C7	123.8 (3)	C22—C21—H21	120.00
N4—C8—C9	117.7 (3)	C22—C23—H23	121.00
C7—C8—C9	118.4 (3)	C24—C23—H23	121.00
C8—C9—C10	121.2 (4)	C23—C24—H24	120.00
C8—C9—C14	120.0 (4)	C25—C24—H24	120.00
C10—C9—C14	118.8 (4)	C24—C25—H25	120.00
C9—C10—C11	120.4 (4)	C26—C25—H25	120.00
C10—C11—C12	120.6 (4)	N6—C26—H26	119.00
C11—C12—C13	119.9 (4)	C25—C26—H26	119.00
C12—C13—C14	120.3 (4)		
C12—Hg1—C11—Hg2	62.36 (4)	C16—N3—C20—C19	0.7 (6)
N2—Hg1—C11—Hg2	-91.91 (9)	Hg1—N3—C16—C17	173.4 (3)
N3—Hg1—C11—Hg2	-25.64 (8)	C20—N3—C16—C17	1.2 (6)
N5—Hg1—C11—Hg2	-159.25 (9)	C20—N3—C16—C15	-177.0 (4)
N6—Hg1—C11—Hg2	146.07 (8)	Hg1—N3—C20—C19	-170.8 (3)
N3—Hg1—N2—N1	-173.7 (3)	N5—N4—C8—C7	-7.7 (5)
N5—Hg1—N2—N1	52.1 (2)	C8—N4—N5—C21	122.7 (4)
C11—Hg1—N5—N4	125.2 (2)	N5—N4—C8—C9	175.4 (3)
C11—Hg1—C12—Hg2 <sup>ii</sup>	112.83 (5)	C8—N4—N5—Hg1	-76.8 (4)
N2—Hg1—C12—Hg2 <sup>ii</sup>	-116.81 (13)	Hg1—N5—C21—C22	11.5 (5)
N3—Hg1—C12—Hg2 <sup>ii</sup>	-154.71 (8)	N4—N5—C21—C22	173.7 (3)
N5—Hg1—C12—Hg2 <sup>ii</sup>	-36.01 (9)	Hg1—N6—C22—C21	-7.5 (4)

N6—Hg1—C12—Hg2 <sup>ii</sup>	26.94 (8)	C22—N6—C26—C25	0.4 (6)
C12—Hg1—N5—C21	65.9 (3)	C26—N6—C22—C23	-0.7 (5)
N2—Hg1—N5—C21	-150.4 (3)	Hg1—N6—C26—C25	-171.1 (3)
N3—Hg1—N5—C21	161.3 (3)	C26—N6—C22—C21	-179.1 (3)
N6—Hg1—N5—C21	-10.5 (3)	Hg1—N6—C22—C23	170.9 (3)
C11—Hg1—N6—C22	147.2 (3)	C7—C1—C6—C5	-177.9 (3)
C11—Hg1—N2—N1	-81.1 (2)	C7—C1—C2—C3	179.1 (4)
C12—Hg1—N2—N1	144.20 (19)	C2—C1—C7—N1	-158.9 (3)
N5—Hg1—N3—C20	-132.0 (3)	C6—C1—C7—C8	-157.5 (3)
N5—Hg1—N6—C22	9.2 (3)	C6—C1—C7—N1	19.8 (5)
N6—Hg1—N2—N1	8.2 (3)	C6—C1—C2—C3	0.3 (6)
C11—Hg1—N2—C15	85.7 (3)	C2—C1—C7—C8	23.8 (5)
C12—Hg1—N2—C15	-49.0 (3)	C2—C1—C6—C5	0.9 (5)
N3—Hg1—N2—C15	-6.9 (3)	C1—C2—C3—C4	-1.1 (6)
N5—Hg1—N2—C15	-141.1 (3)	C2—C3—C4—C5	0.8 (6)
N6—Hg1—N2—C15	175.0 (3)	C3—C4—C5—C6	0.4 (6)
C11—Hg1—N3—C16	-85.6 (3)	C4—C5—C6—C1	-1.2 (6)
C12—Hg1—N3—C16	163.4 (3)	N1—C7—C8—C9	-113.3 (4)
N2—Hg1—N3—C16	5.8 (2)	C1—C7—C8—C9	63.8 (4)
N5—Hg1—N3—C16	56.5 (3)	C1—C7—C8—N4	-113.1 (4)
C11—Hg1—N3—C20	86.0 (3)	N1—C7—C8—N4	69.8 (5)
C12—Hg1—N3—C20	-25.1 (3)	N4—C8—C9—C14	-159.8 (3)
N2—Hg1—N3—C20	177.4 (3)	C7—C8—C9—C14	23.0 (5)
N2—Hg1—N6—C22	55.3 (3)	N4—C8—C9—C10	17.2 (5)
C11—Hg1—N5—C21	-73.9 (3)	C7—C8—C9—C10	-159.9 (4)
C11—Hg1—N6—C26	-41.4 (3)	C10—C9—C14—C13	-1.0 (6)
C12—Hg1—N6—C26	70.1 (3)	C8—C9—C14—C13	176.1 (4)
C12—Hg1—N5—N4	-95.0 (2)	C8—C9—C10—C11	-178.0 (4)
N2—Hg1—N5—N4	48.7 (2)	C14—C9—C10—C11	-1.0 (6)
N3—Hg1—N5—N4	0.4 (3)	C9—C10—C11—C12	1.2 (7)
N6—Hg1—N5—N4	-171.5 (3)	C10—C11—C12—C13	0.6 (7)
C12—Hg1—N6—C22	-101.3 (3)	C11—C12—C13—C14	-2.6 (7)
N2—Hg1—N6—C26	-133.3 (3)	C12—C13—C14—C9	2.9 (6)
N5—Hg1—N6—C26	-179.5 (3)	N2—C15—C16—C17	-180.0 (4)
C14—Hg2—C11—Hg1	-62.53 (5)	N2—C15—C16—N3	-1.8 (6)
C11 <sup>ii</sup> —Hg2 <sup>ii</sup> —C12—Hg1	148.59 (5)	N3—C16—C17—C18	-1.2 (6)
C13 <sup>ii</sup> —Hg2 <sup>ii</sup> —C12—Hg1	-111.36 (5)	C15—C16—C17—C18	176.8 (4)
C14 <sup>ii</sup> —Hg2 <sup>ii</sup> —C12—Hg1	47.22 (6)	C16—C17—C18—C19	-0.6 (6)
C13—Hg2—C11—Hg1	92.17 (5)	C17—C18—C19—C20	2.4 (6)
C12 <sup>i</sup> —Hg2—C11—Hg1	-165.83 (4)	C18—C19—C20—N3	-2.5 (6)
C7—N1—N2—Hg1	-85.4 (3)	N5—C21—C22—C23	178.3 (4)
N2—N1—C7—C1	-178.5 (3)	N5—C21—C22—N6	-3.3 (5)
C7—N1—N2—C15	107.7 (4)	C21—C22—C23—C24	179.1 (4)
N2—N1—C7—C8	-1.4 (5)	N6—C22—C23—C24	0.8 (6)
N1—N2—C15—C16	175.1 (3)	C22—C23—C24—C25	-0.4 (6)

Hg1—N2—C15—C16	7.5 (5)	C23—C24—C25—C26	0.0 (7)
Hg1—N3—C16—C15	-4.7 (4)	C24—C25—C26—N6	0.0 (7)

Symmetry codes: (i)  $-x+1, y+1/2, -z+1/2$ ; (ii)  $-x+1, y-1/2, -z+1/2$ .

*Hydrogen-bond geometry (Å, °)*

Cg5 is the centroid of the C1–C6 phenyl ring.

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C14—H14 $\cdots$ C13 <sup>iii</sup>	0.93	2.82	3.647 (4)	149
C25—H25 $\cdots$ Cg5 <sup>iv</sup>	0.93	2.96	3.814 (5)	153

Symmetry codes: (iii)  $-x+1, -y+2, -z+1$ ; (iv)  $x, -y+3/2, z-1/2$ .