

Ethyl 2-(pyridine-4-carboxamido)-4,5,6,7-tetrahydro-1-benzothiophene-3-carboxylate

Asma Mukhtar,^a M. Nawaz Tahir,^{b*} Misbahul Ain Khan,^c Abdul Qayyum Ather^d and Naveed Sajid^c

^aInstitute of Chemistry, University of the Punjab, Lahore, Pakistan, ^bUniversity of Sargodha, Department of Physics, Sargodha, Pakistan, ^cDepartment of Chemistry, Islamia University, Bahawalpur, Pakistan, and ^dApplied Chemistry Research Center, PCSIR Laboratories Complex, Lahore 54600, Pakistan
Correspondence e-mail: dmntahir_uos@yahoo.com


Received 3 June 2012; accepted 3 June 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; disorder in main residue; R factor = 0.045; wR factor = 0.116; data-to-parameter ratio = 12.0.

In the title compound, $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$, the dihedral angles between the thiophene ring and the ethyl ester group and the pyridine-4-carboxamide unit are $7.1(2)$ and $9.47(11)^\circ$, respectively. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond generates an $S(6)$ ring. In the crystal, inversion dimers linked by pairs of $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds between the tetrahydro-1-benzothiophene and the pyridine-4-carboxamide residues generate $R_2^2(16)$ loops. There exists positional disorder in three methylene groups of the cyclohexane ring and the terminal C atom of the ethyl ester side chain in a $0.691(14):0.309(14)$ occupancy ratio.

Related literature

For related structures, see: Mukhtar *et al.* (2010*a,b*).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$

$M_r = 330.39$

Triclinic, $P\bar{1}$
 $a = 8.5604(6)\text{ \AA}$
 $b = 9.3481(7)\text{ \AA}$
 $c = 11.7443(10)\text{ \AA}$
 $\alpha = 105.121(3)^\circ$
 $\beta = 99.748(2)^\circ$
 $\gamma = 110.806(3)^\circ$

$V = 811.59(11)\text{ \AA}^3$
 $Z = 2$
 $\text{Mo } K\alpha \text{ radiation}$
 $\mu = 0.22\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.24 \times 0.18 \times 0.15\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.953$, $T_{\max} = 0.958$

12176 measured reflections
2914 independent reflections
1842 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.116$
 $S = 1.00$
2914 reflections
243 parameters

10 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.15\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.18\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O2	0.86	1.99	2.650 (3)	132
C7A—H7A \cdots O3 ⁱ	0.97	2.56	3.323 (12)	136

Symmetry code: (i) $-x + 1, -y, -z + 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the provision of funds for the purchase of a diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6838).

References

- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Mukhtar, A., Tahir, M. N., Khan, M. A. & Khan, M. N. (2010*a*). *Acta Cryst. E66*, o2652.
- Mukhtar, A., Tahir, M. N., Khan, M. A. & Khan, M. N. (2010*b*). *Acta Cryst. E66*, o3171.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supporting information

Acta Cryst. (2012). E68, o2041 [https://doi.org/10.1107/S1600536812025251]

Ethyl 2-(pyridine-4-carboxamido)-4,5,6,7-tetrahydro-1-benzothiophene-3-carboxylate

Asma Mukhtar, M. Nawaz Tahir, Misbahul Ain Khan, Abdul Qayyum Ather and Naveed Sajid

S1. Comment

We reported the crystal structures of ethyl 2-benzamido-4,5,6,7-tetrahydro-1-benzothiophene-3-carboxylate (Mukhtar *et al.*, 2010a) and diethyl 5-acetamido-3-methylthiophene-2,4-dicarboxylate (Mukhtar *et al.*, 2010b) which are related to the title compound (I), (Fig. 1).

In (I), the thiophene ring A (S1/C8/C3/C2/C9), ethyl ester group B (O1/C1/O2/C16/C17A) and pyridine-4-carboxamide moiety C (C10—C15/N1/N2/O3) are planar with r. m. s. deviation of 0.0010, 0.0906 and 0.0520 Å, respectively. The dihedral angle between A/B, A/C and B/C is 7.07 (21), 9.47 (11) and 3.30 (20)°, respectively. In the title compound an S(6) ring motif is formed due to intramolecular H-bonding of N—H···O type (Table 1, Fig. 1). The molecules are linked in the form of dimers with $R_2^2(16)$ ring motif due to C—H···O type of H-bonding (Table 1, Fig. 2). Three methylene groups of cyclohexane ring and terminal C-atom of ethyl ester are disordered over two set of sites with occupancy ratio of 0.691 (14):0.309 (14).


S2. Experimental

A mixture of (0.4 g, 3 mmol) of pyridine-4-carboxylic acid and 0.5 ml of thionyl chloride was heated for 5 minutes. Ethyl 2-pyridyl-4-amido-4,5,6,7-tetrahydro-1-benzothiophene-3-carboxylate (0.7 g, 3 mmol) was dissolved in 30 ml chloroform separately and then added to the former mixture. The whole reaction mixture was refluxed for 45 minutes. The solvent was removed and residue was recrystallized in acetone to give colorless prisms of (I). M.p.: 431 K, yield: 0.92 g, 80%.


S3. Refinement

In the cyclohexane ring three methylene groups and terminal C-atom of ethyl ester are disordered over two set of sites with occupancy ratio of 0.691 (14):0.309 (14).

The H-atoms were positioned geometrically (N—H = 0.86, C—H = 0.93–0.97 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl and $x = 1.2$ for other H-atoms.

**Figure 1**

View of the title compound with displacement ellipsoids drawn at the 50% probability level. The dotted line show intramolecular H-bonding.

**Figure 2**

The partial packing, which shows that molecules form dimers with $R_2^2(16)$ ring.

Ethyl 2-(pyridine-4-carboxamido)-4,5,6,7-tetrahydro-1-benzothiophene- 3-carboxylate

Crystal data

$C_{17}H_{18}N_2O_3S$
 $M_r = 330.39$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 8.5604 (6) \text{ \AA}$
 $b = 9.3481 (7) \text{ \AA}$
 $c = 11.7443 (10) \text{ \AA}$
 $\alpha = 105.121 (3)^\circ$
 $\beta = 99.748 (2)^\circ$
 $\gamma = 110.806 (3)^\circ$
 $V = 811.59 (11) \text{ \AA}^3$

$Z = 2$
 $F(000) = 384$
 $D_x = 1.352 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 1842 reflections
 $\theta = 2.5\text{--}25.3^\circ$
 $\mu = 0.22 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
Prism, colorless
 $0.24 \times 0.18 \times 0.15 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 8.10 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.953$, $T_{\max} = 0.958$

12176 measured reflections
 2914 independent reflections
 1842 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -10 \rightarrow 8$
 $k = -11 \rightarrow 11$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.116$
 $S = 1.00$
 2914 reflections
 243 parameters
 10 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.049P)^2 + 0.1304P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.57947 (9)	0.19759 (8)	0.46283 (7)	0.0767 (3)	
O1	0.6976 (2)	0.68634 (18)	0.35270 (15)	0.0742 (6)	
O2	0.8949 (2)	0.73080 (19)	0.52291 (15)	0.0730 (6)	
O3	0.8381 (3)	0.2881 (3)	0.68270 (19)	0.1003 (8)	
N1	0.8635 (2)	0.4829 (2)	0.60071 (17)	0.0615 (7)	
N2	1.3610 (3)	0.7692 (4)	0.9953 (2)	0.0968 (11)	
C1	0.7620 (3)	0.6392 (3)	0.4390 (2)	0.0580 (8)	
C2	0.6600 (3)	0.4677 (3)	0.42082 (19)	0.0517 (8)	
C3	0.5015 (3)	0.3510 (3)	0.3252 (2)	0.0568 (8)	
C4	0.4040 (3)	0.3808 (3)	0.2210 (2)	0.0684 (9)	
C5A	0.2613 (14)	0.2210 (11)	0.1232 (9)	0.084 (3)	0.691 (14)
C6A	0.1643 (9)	0.1068 (9)	0.1818 (7)	0.090 (2)	0.691 (14)
C7A	0.2885 (14)	0.0544 (12)	0.2537 (10)	0.093 (5)	0.691 (14)
C8	0.4453 (3)	0.2032 (3)	0.3374 (2)	0.0647 (9)	
C9	0.7149 (3)	0.3990 (3)	0.5005 (2)	0.0571 (8)	
C10	0.9163 (3)	0.4265 (3)	0.6877 (2)	0.0681 (10)	
C11	1.0748 (3)	0.5483 (3)	0.7916 (2)	0.0624 (9)	

C12	1.1234 (4)	0.5047 (4)	0.8909 (3)	0.0915 (12)	
C13	1.2663 (5)	0.6197 (5)	0.9893 (3)	0.1068 (16)	
C14	1.3131 (4)	0.8068 (4)	0.8988 (3)	0.0810 (11)	
C15	1.1737 (3)	0.7024 (3)	0.7966 (2)	0.0690 (10)	
C16	0.7948 (4)	0.8546 (3)	0.3616 (3)	0.0935 (11)	
C17A	0.7281 (10)	0.8628 (10)	0.2439 (6)	0.113 (3)	0.691 (14)
C5B	0.221 (2)	0.242 (2)	0.159 (3)	0.083 (7)	0.310 (14)
C6B	0.225 (2)	0.0756 (16)	0.1323 (18)	0.091 (6)	0.310 (14)
C7B	0.275 (3)	0.054 (2)	0.2566 (18)	0.074 (8)	0.310 (14)
C17B	0.687 (2)	0.894 (2)	0.2657 (6)	0.103 (5)	0.310 (14)
H4A	0.48597	0.43620	0.18225	0.0821*	
H4B	0.35058	0.45180	0.25392	0.0821*	
H1	0.92888	0.58106	0.60821	0.0738*	
H7A	0.23001	-0.00598	0.30131	0.1119*	0.691 (14)
H7B	0.32310	-0.01579	0.19682	0.1119*	0.691 (14)
H12	1.06183	0.40089	0.89190	0.1100*	
H13	1.29763	0.58902	1.05578	0.1283*	
H14	1.37777	0.91097	0.89972	0.0972*	
H15	1.14681	0.73660	0.73125	0.0828*	
H16A	0.77654	0.92982	0.42663	0.1120*	
H16B	0.91878	0.88177	0.37895	0.1120*	
H17A	0.60304	0.82038	0.22283	0.1699*	0.691 (14)
H17B	0.76116	0.79910	0.18235	0.1699*	0.691 (14)
H17C	0.77538	0.97428	0.24783	0.1699*	0.691 (14)
H5A	0.17977	0.24607	0.07281	0.1004*	0.691 (14)
H5B	0.31459	0.16809	0.06985	0.1004*	0.691 (14)
H6A	0.06979	0.01117	0.11859	0.1079*	0.691 (14)
H6B	0.11397	0.16052	0.23733	0.1079*	0.691 (14)
H5C	0.14711	0.25201	0.21193	0.0997*	0.310 (14)
H5D	0.17039	0.25150	0.08217	0.0997*	0.310 (14)
H6C	0.31023	0.06938	0.08848	0.1103*	0.310 (14)
H6D	0.11147	-0.00965	0.08180	0.1103*	0.310 (14)
H7C	0.29232	-0.04536	0.24441	0.0886*	0.310 (14)
H7D	0.18307	0.04701	0.29589	0.0886*	0.310 (14)
H17D	0.56995	0.86379	0.27276	0.1534*	0.310 (14)
H17E	0.68253	0.83476	0.18437	0.1534*	0.310 (14)
H17F	0.73952	1.00931	0.28019	0.1534*	0.310 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0865 (5)	0.0500 (4)	0.0840 (5)	0.0152 (3)	0.0208 (4)	0.0297 (3)
O1	0.0827 (12)	0.0456 (10)	0.0725 (11)	0.0125 (9)	-0.0016 (9)	0.0224 (8)
O2	0.0732 (11)	0.0510 (10)	0.0704 (11)	0.0079 (9)	-0.0004 (10)	0.0225 (9)
O3	0.1113 (16)	0.0709 (13)	0.1059 (15)	0.0190 (12)	0.0106 (12)	0.0512 (12)
N1	0.0666 (13)	0.0511 (12)	0.0605 (12)	0.0171 (10)	0.0133 (10)	0.0238 (10)
N2	0.0930 (19)	0.110 (2)	0.0823 (17)	0.0380 (17)	0.0115 (14)	0.0410 (16)
C1	0.0664 (16)	0.0493 (14)	0.0562 (14)	0.0225 (13)	0.0151 (13)	0.0193 (12)

C2	0.0573 (14)	0.0414 (13)	0.0528 (13)	0.0167 (11)	0.0177 (11)	0.0152 (10)
C3	0.0581 (14)	0.0496 (14)	0.0562 (14)	0.0177 (12)	0.0194 (12)	0.0133 (11)
C4	0.0658 (16)	0.0581 (15)	0.0681 (16)	0.0203 (13)	0.0112 (13)	0.0147 (12)
C5A	0.074 (4)	0.073 (5)	0.076 (6)	0.019 (3)	0.003 (3)	0.012 (3)
C6A	0.067 (4)	0.072 (4)	0.085 (4)	0.000 (3)	0.003 (3)	0.008 (3)
C7A	0.090 (9)	0.074 (8)	0.098 (9)	0.010 (6)	0.028 (7)	0.037 (6)
C8	0.0640 (16)	0.0470 (15)	0.0688 (16)	0.0117 (13)	0.0176 (13)	0.0156 (12)
C9	0.0635 (15)	0.0458 (13)	0.0590 (14)	0.0177 (12)	0.0229 (13)	0.0178 (12)
C10	0.0803 (18)	0.0627 (17)	0.0696 (17)	0.0299 (15)	0.0245 (14)	0.0343 (14)
C11	0.0681 (16)	0.0712 (18)	0.0609 (15)	0.0365 (14)	0.0223 (13)	0.0306 (13)
C12	0.096 (2)	0.098 (2)	0.086 (2)	0.0341 (19)	0.0211 (19)	0.0535 (19)
C13	0.108 (3)	0.134 (3)	0.083 (2)	0.048 (2)	0.010 (2)	0.060 (2)
C14	0.0760 (19)	0.084 (2)	0.0749 (19)	0.0317 (16)	0.0091 (16)	0.0260 (16)
C15	0.0719 (17)	0.0702 (17)	0.0687 (17)	0.0327 (15)	0.0147 (14)	0.0295 (14)
C16	0.108 (2)	0.0462 (16)	0.097 (2)	0.0126 (15)	-0.0073 (17)	0.0308 (15)
C17A	0.155 (6)	0.062 (5)	0.118 (5)	0.040 (4)	0.010 (4)	0.050 (4)
C5B	0.071 (11)	0.066 (9)	0.081 (14)	0.023 (8)	-0.005 (7)	0.001 (8)
C6B	0.073 (9)	0.074 (9)	0.098 (12)	0.025 (6)	0.004 (7)	0.006 (7)
C7B	0.060 (13)	0.027 (10)	0.083 (18)	-0.002 (9)	0.000 (11)	-0.019 (10)
C17B	0.151 (12)	0.052 (7)	0.076 (8)	0.055 (8)	-0.032 (7)	0.002 (5)

Geometric parameters (\AA , $^\circ$)

S1—C8	1.733 (3)	C16—C17A	1.436 (8)
S1—C9	1.711 (3)	C16—C17B	1.537 (14)
O1—C1	1.320 (3)	C4—H4A	0.9700
O1—C16	1.460 (3)	C4—H4B	0.9700
O2—C1	1.219 (3)	C5A—H5A	0.9700
O3—C10	1.207 (4)	C5A—H5B	0.9700
N1—C9	1.386 (3)	C5B—H5D	0.9700
N1—C10	1.349 (3)	C5B—H5C	0.9700
N2—C13	1.319 (6)	C6A—H6B	0.9700
N2—C14	1.315 (4)	C6A—H6A	0.9700
N1—H1	0.8600	C6B—H6D	0.9700
C1—C2	1.464 (4)	C6B—H6C	0.9700
C2—C3	1.444 (3)	C7A—H7B	0.9700
C2—C9	1.374 (4)	C7A—H7A	0.9700
C3—C4	1.506 (3)	C7B—H7D	0.9700
C3—C8	1.347 (4)	C7B—H7C	0.9700
C4—C5B	1.53 (2)	C12—H12	0.9300
C4—C5A	1.542 (11)	C13—H13	0.9300
C5A—C6A	1.500 (13)	C14—H14	0.9300
C5B—C6B	1.52 (3)	C15—H15	0.9300
C6A—C7A	1.538 (14)	C16—H16A	0.9700
C6B—C7B	1.54 (3)	C16—H16B	0.9700
C7A—C8	1.489 (12)	C17A—H17C	0.9600
C7B—C8	1.54 (2)	C17A—H17A	0.9600
C10—C11	1.498 (3)	C17A—H17B	0.9600

C11—C12	1.380 (4)	C17B—H17D	0.9600
C11—C15	1.367 (4)	C17B—H17E	0.9600
C12—C13	1.388 (5)	C17B—H17F	0.9600
C14—C15	1.376 (4)		
C8—S1—C9	90.88 (13)	H5A—C5A—H5B	108.00
C1—O1—C16	116.6 (2)	C4—C5B—H5C	109.00
C9—N1—C10	126.8 (2)	C4—C5B—H5D	109.00
C13—N2—C14	115.6 (3)	C6B—C5B—H5C	109.00
C10—N1—H1	117.00	C6B—C5B—H5D	109.00
C9—N1—H1	117.00	H5C—C5B—H5D	108.00
O1—C1—C2	113.0 (2)	H6A—C6A—H6B	108.00
O2—C1—C2	124.7 (2)	C5A—C6A—H6A	110.00
O1—C1—O2	122.3 (2)	C5A—C6A—H6B	110.00
C3—C2—C9	111.7 (2)	C7A—C6A—H6A	110.00
C1—C2—C3	128.5 (2)	C7A—C6A—H6B	110.00
C1—C2—C9	119.8 (2)	C7B—C6B—H6C	110.00
C4—C3—C8	121.2 (2)	C7B—C6B—H6D	110.00
C2—C3—C8	111.9 (2)	C5B—C6B—H6D	110.00
C2—C3—C4	126.9 (2)	C5B—C6B—H6C	110.00
C3—C4—C5B	110.8 (10)	H6C—C6B—H6D	108.00
C3—C4—C5A	112.2 (4)	C6A—C7A—H7A	110.00
C4—C5A—C6A	111.4 (7)	C6A—C7A—H7B	110.00
C4—C5B—C6B	111.8 (14)	H7A—C7A—H7B	108.00
C5A—C6A—C7A	110.2 (8)	C8—C7A—H7B	110.00
C5B—C6B—C7B	107.5 (18)	C8—C7A—H7A	110.00
C6A—C7A—C8	108.7 (8)	C6B—C7B—H7C	110.00
C6B—C7B—C8	107.5 (13)	C8—C7B—H7D	110.00
C3—C8—C7B	125.6 (8)	C8—C7B—H7C	110.00
C3—C8—C7A	126.1 (5)	C6B—C7B—H7D	110.00
S1—C8—C3	112.96 (19)	H7C—C7B—H7D	109.00
S1—C8—C7A	121.0 (5)	C11—C12—H12	121.00
S1—C8—C7B	121.3 (8)	C13—C12—H12	121.00
S1—C9—N1	123.36 (19)	C12—C13—H13	118.00
S1—C9—C2	112.64 (19)	N2—C13—H13	118.00
N1—C9—C2	124.0 (2)	C15—C14—H14	118.00
N1—C10—C11	115.4 (2)	N2—C14—H14	118.00
O3—C10—C11	122.7 (2)	C11—C15—H15	120.00
O3—C10—N1	121.8 (2)	C14—C15—H15	120.00
C10—C11—C15	124.5 (2)	O1—C16—H16A	111.00
C10—C11—C12	118.4 (3)	O1—C16—H16B	111.00
C12—C11—C15	117.1 (2)	C17A—C16—H16A	111.00
C11—C12—C13	118.4 (3)	C17A—C16—H16B	111.00
N2—C13—C12	124.8 (3)	H16A—C16—H16B	109.00
N2—C14—C15	124.4 (3)	C17B—C16—H16A	89.00
C11—C15—C14	119.7 (3)	C17B—C16—H16B	126.00
O1—C16—C17B	108.9 (7)	C16—C17A—H17A	109.00
O1—C16—C17A	105.7 (4)	C16—C17A—H17B	109.00

C3—C4—H4A	109.00	C16—C17A—H17C	109.00
C3—C4—H4B	109.00	H17A—C17A—H17B	109.00
C5A—C4—H4A	109.00	H17A—C17A—H17C	109.00
C5A—C4—H4B	109.00	H17B—C17A—H17C	110.00
H4A—C4—H4B	108.00	C16—C17B—H17D	110.00
C5B—C4—H4A	128.00	C16—C17B—H17E	110.00
C5B—C4—H4B	89.00	C16—C17B—H17F	110.00
C4—C5A—H5A	109.00	H17D—C17B—H17E	109.00
C4—C5A—H5B	109.00	H17D—C17B—H17F	109.00
C6A—C5A—H5A	109.00	H17E—C17B—H17F	109.00
C6A—C5A—H5B	109.00		
C9—S1—C8—C3	0.1 (2)	C3—C2—C9—S1	-0.2 (3)
C9—S1—C8—C7A	178.7 (6)	C3—C2—C9—N1	179.9 (2)
C8—S1—C9—N1	-180.0 (2)	C2—C3—C4—C5A	169.0 (5)
C8—S1—C9—C2	0.1 (2)	C8—C3—C4—C5A	-10.4 (6)
C16—O1—C1—O2	0.6 (4)	C2—C3—C8—S1	-0.2 (3)
C16—O1—C1—C2	-178.1 (2)	C2—C3—C8—C7A	-178.7 (6)
C1—O1—C16—C17A	165.1 (4)	C4—C3—C8—S1	179.3 (2)
C10—N1—C9—S1	-5.8 (4)	C4—C3—C8—C7A	0.8 (7)
C10—N1—C9—C2	174.1 (2)	C3—C4—C5A—C6A	41.5 (9)
C9—N1—C10—O3	3.3 (4)	C4—C5A—C6A—C7A	-63.6 (10)
C9—N1—C10—C11	-175.2 (2)	C5A—C6A—C7A—C8	51.1 (10)
C14—N2—C13—C12	-0.6 (6)	C6A—C7A—C8—S1	160.6 (5)
C13—N2—C14—C15	0.6 (5)	C6A—C7A—C8—C3	-21.0 (10)
O1—C1—C2—C3	-1.7 (4)	O3—C10—C11—C12	-6.1 (4)
O1—C1—C2—C9	177.4 (2)	O3—C10—C11—C15	175.4 (3)
O2—C1—C2—C3	179.6 (3)	N1—C10—C11—C12	172.3 (3)
O2—C1—C2—C9	-1.3 (4)	N1—C10—C11—C15	-6.2 (4)
C1—C2—C3—C4	-0.1 (4)	C10—C11—C12—C13	-177.6 (3)
C1—C2—C3—C8	179.4 (3)	C15—C11—C12—C13	1.0 (5)
C9—C2—C3—C4	-179.3 (2)	C10—C11—C15—C14	177.5 (3)
C9—C2—C3—C8	0.2 (3)	C12—C11—C15—C14	-1.0 (4)
C1—C2—C9—S1	-179.41 (19)	C11—C12—C13—N2	-0.3 (6)
C1—C2—C9—N1	0.6 (4)	N2—C14—C15—C11	0.2 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···O2	0.86	1.99	2.650 (3)	132
C7A—H7A···O3 ⁱ	0.97	2.56	3.323 (12)	136

Symmetry code: (i) $-x+1, -y, -z+1$.