

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[*(1,10-phenanthroline-κ²N,N')*lanthanum(III)]-μ-(5-bromo-2-hydroxybenzoato)-κ²O¹:O^{1'}-di-μ-chlorido]

Wen-Qing Zhu,* Jin-Ping Wang, Yu-Kun Yuan and Zhuo Li

College of Environment and Chemical Engineering, Xi'an Polytechnic University, Xi'an, Shaanxi 710048, People's Republic of China

Correspondence e-mail: wqzhu2008@163.com


Received 14 May 2012; accepted 5 June 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.011$ Å; R factor = 0.034; wR factor = 0.125; data-to-parameter ratio = 14.2.

In the title complex, $[\text{La}(\text{C}_7\text{H}_4\text{BrO}_3)\text{Cl}_2(\text{C}_{12}\text{H}_8\text{N}_2)]_n$, the La^{III} ion is eight-coordinated by two carboxylate O atoms from two 5-bromosalicylate ligands, two N atoms from a chelating 1,10-phenanthroline ligand and four bridging Cl atoms in a distorted square-antiprismatic geometry. The La^{III} ions are linked by bridging carboxylate groups and chloride anions into a chain along [100]. An intramolecular O—H...O hydrogen bond is formed in the 5-bromosalicylate ligand. π – π interactions between the pyridine and benzene rings and between the benzene rings are observed [centroid–centroid distances = 3.794 (5) and 3.804 (4) Å].

Related literature

For background to rare earth carboxylates, see: Ali *et al.* (2004); Costes *et al.* (2002); Kaur *et al.* (2010); Yin & Sun (2004). For complexes with salicylate ligands, see: Hu *et al.* (2005); Yin *et al.* (2004).



Experimental

Crystal data

$[\text{La}(\text{C}_7\text{H}_4\text{BrO}_3)\text{Cl}_2(\text{C}_{12}\text{H}_8\text{N}_2)]$
 $M_r = 606.02$
 Orthorhombic, *Pbca*
 $a = 8.2957$ (10) Å
 $b = 22.104$ (3) Å
 $c = 22.110$ (3) Å
 $V = 4054.3$ (9) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 4.36$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.24 \times 0.14$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\text{min}} = 0.352$, $T_{\text{max}} = 0.584$
 19025 measured reflections
 3598 independent reflections
 2849 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.125$
 $S = 1.00$
 3598 reflections
 254 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.20$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H3}\cdots\text{O2}$	0.82	1.93	2.588 (6)	137

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

This project was supported by the Shaanxi Science Research Project (No. 2011 K07–23), the Scientific Research Special Plan of Shaanxi Department of Education (No. 11 J K0566) and the Foundation of Xi'an Polytechnic University (No. BS1013).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2550).

References

- Ali, H. M., Khamis, N. A. & Yamin, B. M. (2004). *Acta Cryst.* **E60**, m1708–m1709.
 Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Costes, J.-P., Clemente-Juan, J. M., Dahan, F., Nicodème, F. & Verelst, M. (2002). *Angew. Chem. Int. Ed.* **41**, 323–325.
 Hu, M., Geng, C., Li, S., Du, Y., Jiang, Y. & Liu, Z. (2005). *J. Organomet. Chem.* **690**, 3118–3124.
 Kaur, G., Dwivedi, Y. & Rai, S. B. (2010). *Opt. Commun.* **283**, 3441–3447.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yin, M.-C., Ai, C.-C., Yuan, L.-J., Wang, C.-W. & Sun, J.-T. (2004). *J. Mol. Struct.* **691**, 33–37.
 Yin, M.-C. & Sun, J.-T. (2004). *J. Alloys Compd.* **381**, 50–57.

supporting information

Acta Cryst. (2012). E68, m896 [https://doi.org/10.1107/S1600536812025500]

***catena*-Poly[[*(1,10-phenanthroline- κ^2 N,N')*lanthanum(III)]- μ -(5-bromo-2-hydroxybenzoato)- κ^2 O¹:O^{1'}-di- μ -chlorido]**

Wen-Qing Zhu, Jin-Ping Wang, Yu-Kun Yuan and Zhuo Li

S1. Comment

Over the past years, much attention has been paid to the rare earth carboxylates, owing to their novel structures and potential applications in a wide range of materials science, such as superconductor, magnetic materials and luminescent probes (Ali *et al.*, 2004; Costes *et al.*, 2002; Kaur *et al.*, 2010; Yin & Sun, 2004). Because salicylic acid and its derivatives have been known for a long time to possess anti-inflammatory activity and have also been considered of interest from a structural point of view (Hu *et al.*, 2005; Yin *et al.*, 2004), a new lanthanum bromosalicylate complex was synthesized and its crystal structure is reported here.

S2. Experimental

An ethanol solution (5 ml) of LaCl₃·6H₂O (0.5 mmol, 0.177 g) was added dropwise to an ethanol solution (5 ml) of 5-bromosalicylic acid (0.5 mmol, 0.109 g) under stirring and then an ethanol solution (5 ml) of 1,10-phenanthroline (0.5 mmol, 0.084 g) was added dropwise. The mixture was stirred for about 13 min at room temperature and sealed in a Teflon-lined stainless autoclave and heated to 120°C for 60 h. After filtered, colorless flake-shaped crystals were obtained.

S3. Refinement

H atoms were positioned geometrically and refined as riding atoms, with O—H = 0.82 and C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{O})$. The highest residual electron density was found 1.20 Å from Br1 atom and the deepest hole 1.34 Å from La1 atom.




Figure 1

The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity. [Symmetry codes: (a) $x + 1/2, -y + 1/2, -z + 1$; (b) $x - 1/2, -y + 1/2, -z + 1$.]




Figure 2

The one-dimensional chain structure in the title compound. Hydrogen bonds are shown as dashed lines.

catena-Poly[[[(1,10-phenanthroline- κ^2N,N')lanthanum(III)]- μ -(5-bromo-2-hydroxybenzoato)- $\kappa^2O^1:O^1'$ -di- μ -chlorido]

Crystal data

[La(C₇H₄BrO₃)Cl₂(C₁₂H₈N₂)]

$M_r = 606.02$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 8.2957$ (10) Å

$b = 22.104$ (3) Å

$c = 22.110$ (3) Å

$V = 4054.3$ (9) Å³

$Z = 8$

$F(000) = 2320$

$D_x = 1.986$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3026 reflections

$\theta = 2.6$ – 24.6°

$\mu = 4.36$ mm⁻¹

$T = 296$ K

Flake, colorless

$0.30 \times 0.24 \times 0.14$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.352$, $T_{\max} = 0.584$

19025 measured reflections

3598 independent reflections

2849 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.065$

$\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 1.8^\circ$

$h = -9 \rightarrow 7$

$k = -26 \rightarrow 26$

$l = -26 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.125$

$S = 1.00$

3598 reflections

254 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.080P)^2 + 0.7837P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 1.20$ e Å⁻³

$\Delta\rho_{\min} = -1.34$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
La1	0.78367 (4)	0.242463 (15)	0.537359 (15)	0.02456 (15)
Br1	0.86777 (13)	-0.05046 (5)	0.66282 (5)	0.0877 (4)
Cl1	0.57034 (19)	0.18099 (7)	0.45631 (7)	0.0352 (4)
Cl2	0.46216 (18)	0.27765 (8)	0.57361 (7)	0.0365 (4)

N1	0.6683 (7)	0.1654 (3)	0.6202 (2)	0.0401 (14)
N2	0.8088 (7)	0.2724 (3)	0.6548 (2)	0.0396 (14)
O1	0.9829 (5)	0.16436 (19)	0.5594 (2)	0.0389 (11)
O2	1.2182 (5)	0.1607 (2)	0.5117 (2)	0.0423 (12)
O3	1.3579 (7)	0.0573 (2)	0.4954 (3)	0.0615 (15)
H3	1.3282	0.0899	0.4818	0.092*
C1	0.5984 (9)	0.1136 (3)	0.6047 (4)	0.0510 (19)
H1	0.5907	0.1048	0.5637	0.061*
C2	0.5357 (10)	0.0711 (4)	0.6451 (4)	0.067 (2)
H2	0.4911	0.0349	0.6317	0.080*
C3	0.5428 (11)	0.0852 (5)	0.7052 (4)	0.075 (3)
H3A	0.5001	0.0585	0.7334	0.090*
C4	0.6131 (10)	0.1391 (4)	0.7249 (3)	0.059 (2)
C5	0.6262 (12)	0.1591 (5)	0.7881 (3)	0.082 (3)
H5	0.5853	0.1341	0.8184	0.098*
C6	0.6954 (12)	0.2124 (5)	0.8043 (4)	0.077 (3)
H6	0.7022	0.2228	0.8450	0.093*
C7	0.7564 (11)	0.2516 (4)	0.7606 (4)	0.056 (2)
C8	0.8293 (11)	0.3067 (4)	0.7754 (4)	0.065 (2)
H8	0.8383	0.3182	0.8157	0.078*
C9	0.8855 (11)	0.3424 (4)	0.7323 (4)	0.062 (2)
H9	0.9332	0.3792	0.7419	0.074*
C10	0.8721 (9)	0.3240 (3)	0.6727 (3)	0.0482 (19)
H10	0.9107	0.3501	0.6431	0.058*
C11	0.7488 (9)	0.2351 (3)	0.6985 (3)	0.0417 (17)
C12	0.6755 (8)	0.1795 (3)	0.6812 (3)	0.0417 (17)
C13	1.0177 (9)	-0.0158 (3)	0.6085 (3)	0.0481 (18)
C14	1.1393 (10)	-0.0516 (3)	0.5865 (4)	0.056 (2)
H14	1.1465	-0.0922	0.5971	0.067*
C15	1.2497 (11)	-0.0257 (4)	0.5485 (4)	0.060 (2)
H15	1.3323	-0.0496	0.5331	0.072*
C16	1.2430 (9)	0.0348 (3)	0.5321 (3)	0.0442 (18)
C17	1.1162 (8)	0.0710 (3)	0.5553 (3)	0.0316 (14)
C18	1.0040 (8)	0.0436 (3)	0.5932 (3)	0.0397 (16)
H18	0.9185	0.0662	0.6083	0.048*
C19	1.1035 (8)	0.1364 (3)	0.5407 (3)	0.0352 (15)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
La1	0.0211 (2)	0.0260 (2)	0.0266 (2)	0.00183 (13)	-0.00042 (14)	0.00254 (13)
Br1	0.0732 (7)	0.0716 (7)	0.1182 (9)	-0.0105 (5)	0.0171 (6)	0.0530 (6)
Cl1	0.0248 (8)	0.0350 (9)	0.0459 (9)	0.0010 (6)	-0.0002 (7)	-0.0084 (7)
Cl2	0.0264 (8)	0.0551 (10)	0.0282 (8)	0.0097 (7)	-0.0002 (6)	-0.0023 (7)
N1	0.031 (3)	0.045 (3)	0.044 (3)	0.003 (3)	0.000 (3)	0.012 (3)
N2	0.032 (3)	0.052 (4)	0.035 (3)	0.011 (3)	-0.001 (2)	0.004 (3)
O1	0.034 (3)	0.039 (3)	0.043 (3)	0.012 (2)	0.009 (2)	0.012 (2)
O2	0.031 (3)	0.032 (3)	0.064 (3)	0.001 (2)	0.010 (2)	0.018 (2)

O3	0.051 (3)	0.045 (3)	0.088 (4)	0.017 (3)	0.032 (3)	0.013 (3)
C1	0.051 (5)	0.048 (4)	0.054 (4)	-0.001 (4)	-0.003 (4)	0.013 (4)
C2	0.059 (6)	0.061 (5)	0.081 (6)	-0.013 (4)	-0.005 (5)	0.032 (5)
C3	0.059 (6)	0.095 (7)	0.071 (6)	-0.014 (5)	0.005 (5)	0.050 (5)
C4	0.046 (5)	0.077 (6)	0.054 (5)	0.007 (4)	0.011 (4)	0.030 (4)
C5	0.077 (7)	0.137 (10)	0.032 (5)	0.018 (7)	0.012 (4)	0.044 (5)
C6	0.076 (7)	0.111 (9)	0.044 (5)	0.008 (6)	-0.004 (5)	0.010 (6)
C7	0.057 (6)	0.085 (7)	0.027 (4)	0.026 (4)	0.002 (4)	0.010 (4)
C8	0.074 (6)	0.079 (7)	0.043 (5)	0.020 (5)	-0.020 (4)	-0.022 (5)
C9	0.071 (6)	0.068 (6)	0.045 (5)	0.013 (5)	-0.016 (4)	-0.011 (4)
C10	0.046 (5)	0.055 (5)	0.043 (4)	0.008 (4)	-0.002 (3)	-0.013 (3)
C11	0.031 (4)	0.058 (5)	0.036 (4)	0.016 (3)	0.000 (3)	0.005 (3)
C12	0.023 (3)	0.064 (5)	0.038 (4)	0.017 (3)	0.007 (3)	0.018 (3)
C13	0.047 (5)	0.039 (4)	0.058 (5)	-0.008 (3)	-0.002 (4)	0.014 (3)
C14	0.060 (5)	0.032 (4)	0.076 (6)	0.000 (4)	0.000 (5)	0.010 (4)
C15	0.063 (5)	0.033 (4)	0.084 (6)	0.020 (4)	0.000 (5)	-0.004 (4)
C16	0.038 (4)	0.032 (4)	0.062 (5)	0.004 (3)	-0.001 (4)	-0.004 (3)
C17	0.030 (4)	0.024 (3)	0.041 (4)	0.002 (3)	-0.007 (3)	0.002 (3)
C18	0.034 (4)	0.036 (4)	0.048 (4)	0.001 (3)	-0.001 (3)	0.010 (3)
C19	0.032 (4)	0.037 (4)	0.037 (4)	0.004 (3)	-0.007 (3)	0.002 (3)

Geometric parameters (Å, °)

La1—O1	2.439 (4)	C3—H3A	0.9300
La1—O2 ⁱ	2.461 (4)	C4—C12	1.415 (10)
La1—N1	2.678 (5)	C4—C5	1.469 (12)
La1—N2	2.687 (6)	C5—C6	1.359 (14)
La1—C11	2.8618 (16)	C5—H5	0.9300
La1—C12	2.8915 (16)	C6—C7	1.393 (14)
La1—C12 ⁱⁱ	2.9001 (16)	C6—H6	0.9300
La1—C11 ⁱⁱ	2.9219 (16)	C7—C8	1.397 (12)
Br1—C13	1.891 (7)	C7—C11	1.422 (11)
Cl1—La1 ⁱ	2.9219 (16)	C8—C9	1.321 (12)
Cl2—La1 ⁱ	2.9001 (16)	C8—H8	0.9300
N1—C1	1.328 (9)	C9—C10	1.383 (10)
N1—C12	1.386 (9)	C9—H9	0.9300
N2—C10	1.317 (9)	C10—H10	0.9300
N2—C11	1.365 (10)	C11—C12	1.424 (11)
O1—C19	1.246 (8)	C13—C18	1.361 (9)
O2—C19	1.266 (8)	C13—C14	1.369 (10)
O2—La1 ⁱⁱ	2.461 (4)	C14—C15	1.368 (12)
O3—C16	1.348 (9)	C14—H14	0.9300
O3—H3	0.8200	C15—C16	1.386 (12)
C1—C2	1.395 (10)	C15—H15	0.9300
C1—H1	0.9300	C16—C17	1.418 (9)
C2—C3	1.365 (13)	C17—C18	1.392 (9)
C2—H2	0.9300	C17—C19	1.485 (8)
C3—C4	1.396 (13)	C18—H18	0.9300

O1—La1—O2 ⁱ	148.59 (15)	C3—C4—C12	118.5 (8)
O1—La1—N1	69.84 (15)	C3—C4—C5	125.7 (8)
O2 ⁱ —La1—N1	140.90 (16)	C12—C4—C5	115.7 (9)
O1—La1—N2	85.91 (16)	C6—C5—C4	122.8 (8)
O2 ⁱ —La1—N2	103.23 (17)	C6—C5—H5	118.6
N1—La1—N2	61.55 (19)	C4—C5—H5	118.6
O1—La1—C11	102.01 (12)	C5—C6—C7	120.7 (9)
O2 ⁱ —La1—C11	90.03 (11)	C5—C6—H6	119.6
N1—La1—C11	84.55 (13)	C7—C6—H6	119.6
N2—La1—C11	140.34 (13)	C6—C7—C8	122.5 (9)
O1—La1—C12	139.46 (11)	C6—C7—C11	119.6 (9)
O2 ⁱ —La1—C12	71.62 (11)	C8—C7—C11	117.8 (8)
N1—La1—C12	69.63 (12)	C9—C8—C7	120.4 (8)
N2—La1—C12	74.78 (12)	C9—C8—H8	119.8
C11—La1—C12	74.39 (4)	C7—C8—H8	119.8
O1—La1—C12 ⁱⁱ	73.42 (11)	C8—C9—C10	118.8 (8)
O2 ⁱ —La1—C12 ⁱⁱ	82.71 (12)	C8—C9—H9	120.6
N1—La1—C12 ⁱⁱ	131.57 (13)	C10—C9—H9	120.6
N2—La1—C12 ⁱⁱ	144.64 (13)	N2—C10—C9	125.0 (8)
C11—La1—C12 ⁱⁱ	73.33 (4)	N2—C10—H10	117.5
C12—La1—C12 ⁱⁱ	138.31 (4)	C9—C10—H10	117.5
O1—La1—C11 ⁱⁱ	81.30 (11)	N2—C11—C7	120.9 (8)
O2 ⁱ —La1—C11 ⁱⁱ	72.36 (11)	N2—C11—C12	119.0 (7)
N1—La1—C11 ⁱⁱ	128.79 (13)	C7—C11—C12	120.0 (7)
N2—La1—C11 ⁱⁱ	75.41 (13)	N1—C12—C4	120.5 (7)
C11—La1—C11 ⁱⁱ	143.93 (3)	N1—C12—C11	118.4 (6)
C12—La1—C11 ⁱⁱ	125.57 (5)	C4—C12—C11	121.1 (7)
C12 ⁱⁱ —La1—C11 ⁱⁱ	73.37 (4)	C18—C13—C14	122.0 (7)
La1—C11—La1 ⁱ	101.44 (5)	C18—C13—Br1	119.5 (6)
La1—C12—La1 ⁱ	101.25 (5)	C14—C13—Br1	118.4 (5)
C1—N1—C12	117.6 (6)	C15—C14—C13	118.0 (7)
C1—N1—La1	121.9 (5)	C15—C14—H14	121.0
C12—N1—La1	120.4 (4)	C13—C14—H14	121.0
C10—N2—C11	117.0 (7)	C14—C15—C16	122.4 (8)
C10—N2—La1	122.4 (5)	C14—C15—H15	118.8
C11—N2—La1	120.6 (5)	C16—C15—H15	118.8
C19—O1—La1	146.0 (4)	O3—C16—C15	119.0 (7)
C19—O2—La1 ⁱⁱ	139.2 (4)	O3—C16—C17	122.2 (6)
C16—O3—H3	109.5	C15—C16—C17	118.7 (8)
N1—C1—C2	125.3 (8)	C18—C17—C16	117.9 (6)
N1—C1—H1	117.4	C18—C17—C19	120.5 (6)
C2—C1—H1	117.4	C16—C17—C19	121.7 (6)
C3—C2—C1	116.9 (8)	C13—C18—C17	120.9 (7)
C3—C2—H2	121.5	C13—C18—H18	119.5
C1—C2—H2	121.5	C17—C18—H18	119.5
C2—C3—C4	121.1 (8)	O1—C19—O2	124.2 (6)
C2—C3—H3A	119.5	O1—C19—C17	117.9 (6)

C4—C3—H3A	119.5	O2—C19—C17	117.9 (6)
O1—La1—C11—La1 ⁱ	161.75 (11)	C2—C3—C4—C5	179.5 (9)
O2 ⁱ —La1—C11—La1 ⁱ	-47.48 (12)	C3—C4—C5—C6	-179.6 (10)
N1—La1—C11—La1 ⁱ	93.73 (13)	C12—C4—C5—C6	-0.9 (14)
N2—La1—C11—La1 ⁱ	63.6 (2)	C4—C5—C6—C7	0.9 (16)
Cl2—La1—C11—La1 ⁱ	23.40 (5)	C5—C6—C7—C8	-179.7 (10)
Cl2 ⁱⁱ —La1—C11—La1 ⁱ	-129.86 (6)	C5—C6—C7—C11	-1.2 (14)
Cl1 ⁱⁱⁱ —La1—C11—La1 ⁱ	-106.56 (8)	C6—C7—C8—C9	180.0 (9)
O1—La1—Cl2—La1 ⁱ	-113.89 (18)	C11—C7—C8—C9	1.5 (13)
O2 ⁱ —La1—Cl2—La1 ⁱ	71.80 (13)	C7—C8—C9—C10	-0.6 (13)
N1—La1—Cl2—La1 ⁱ	-113.49 (14)	C11—N2—C10—C9	1.6 (11)
N2—La1—Cl2—La1 ⁱ	-178.30 (15)	La1—N2—C10—C9	178.7 (6)
Cl1—La1—Cl2—La1 ⁱ	-23.57 (5)	C8—C9—C10—N2	-1.0 (13)
Cl2 ⁱⁱ —La1—Cl2—La1 ⁱ	16.84 (4)	C10—N2—C11—C7	-0.5 (10)
Cl1 ⁱⁱⁱ —La1—Cl2—La1 ⁱ	122.73 (5)	La1—N2—C11—C7	-177.7 (5)
O1—La1—N1—C1	-84.2 (5)	C10—N2—C11—C12	178.4 (6)
O2 ⁱ —La1—N1—C1	104.1 (6)	La1—N2—C11—C12	1.2 (8)
N2—La1—N1—C1	179.4 (6)	C6—C7—C11—N2	-179.5 (8)
Cl1—La1—N1—C1	20.8 (5)	C8—C7—C11—N2	-0.9 (12)
Cl2—La1—N1—C1	96.1 (5)	C6—C7—C11—C12	1.6 (12)
Cl2 ⁱⁱ —La1—N1—C1	-41.2 (6)	C8—C7—C11—C12	-179.9 (7)
Cl1 ⁱⁱⁱ —La1—N1—C1	-144.0 (5)	C1—N1—C12—C4	1.2 (9)
O1—La1—N1—C12	97.8 (5)	La1—N1—C12—C4	179.3 (5)
O2 ⁱ —La1—N1—C12	-73.9 (5)	C1—N1—C12—C11	-179.5 (6)
N2—La1—N1—C12	1.4 (4)	La1—N1—C12—C11	-1.4 (8)
Cl1—La1—N1—C12	-157.2 (5)	C3—C4—C12—N1	-0.7 (11)
Cl2—La1—N1—C12	-81.9 (4)	C5—C4—C12—N1	-179.5 (7)
Cl2 ⁱⁱ —La1—N1—C12	140.8 (4)	C3—C4—C12—C11	-179.9 (7)
Cl1 ⁱⁱⁱ —La1—N1—C12	38.0 (5)	C5—C4—C12—C11	1.3 (11)
O1—La1—N2—C10	112.4 (5)	N2—C11—C12—N1	0.2 (10)
O2 ⁱ —La1—N2—C10	-37.2 (5)	C7—C11—C12—N1	179.1 (6)
N1—La1—N2—C10	-178.4 (6)	N2—C11—C12—C4	179.4 (6)
Cl1—La1—N2—C10	-143.7 (5)	C7—C11—C12—C4	-1.7 (11)
Cl2—La1—N2—C10	-103.6 (5)	C18—C13—C14—C15	0.5 (12)
Cl2 ⁱⁱ —La1—N2—C10	58.9 (6)	Br1—C13—C14—C15	-178.3 (6)
Cl1 ⁱⁱⁱ —La1—N2—C10	30.3 (5)	C13—C14—C15—C16	0.5 (13)
O1—La1—N2—C11	-70.6 (5)	C14—C15—C16—O3	178.7 (8)
O2 ⁱ —La1—N2—C11	139.9 (5)	C14—C15—C16—C17	-0.5 (13)
N1—La1—N2—C11	-1.3 (5)	O3—C16—C17—C18	-179.6 (7)
Cl1—La1—N2—C11	33.4 (6)	C15—C16—C17—C18	-0.4 (10)
Cl2—La1—N2—C11	73.4 (5)	O3—C16—C17—C19	-0.6 (11)
Cl2 ⁱⁱ —La1—N2—C11	-124.0 (5)	C15—C16—C17—C19	178.6 (7)
Cl1 ⁱⁱⁱ —La1—N2—C11	-152.6 (5)	C14—C13—C18—C17	-1.4 (11)
O2 ⁱ —La1—O1—C19	-32.9 (9)	Br1—C13—C18—C17	177.4 (5)
N1—La1—O1—C19	157.1 (8)	C16—C17—C18—C13	1.3 (10)
N2—La1—O1—C19	-141.8 (8)	C19—C17—C18—C13	-177.7 (6)
Cl1—La1—O1—C19	77.5 (8)	La1—O1—C19—O2	39.6 (12)

C12—La1—O1—C19	157.5 (7)	La1—O1—C19—C17	-142.6 (6)
C12 ⁱⁱ —La1—O1—C19	9.2 (8)	La1 ⁱⁱ —O2—C19—O1	-9.6 (11)
C11 ⁱⁱⁱ —La1—O1—C19	-65.9 (8)	La1 ⁱⁱⁱ —O2—C19—C17	172.6 (4)
C12—N1—C1—C2	-2.0 (11)	C18—C17—C19—O1	-5.8 (9)
La1—N1—C1—C2	179.9 (6)	C16—C17—C19—O1	175.2 (6)
N1—C1—C2—C3	2.1 (13)	C18—C17—C19—O2	172.2 (6)
C1—C2—C3—C4	-1.4 (14)	C16—C17—C19—O2	-6.8 (9)
C2—C3—C4—C12	0.8 (13)		

Symmetry codes: (i) $x-1/2, -y+1/2, -z+1$; (ii) $x+1/2, -y+1/2, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O3—H3...O2	0.82	1.93	2.588 (6)	137