

Poly[μ_2 -aqua- μ_4 -(2-{3-[(6-chloropyridin-3-yl)methyl]-2-oxoimidazolidin-1-yl}-acetato)-sodium]

Rajni Kant,^{a*} Vivek K. Gupta,^a Kamini Kapoor,^a Chetan S. Shripanavar,^b Kaushik Banerjee^c and Madhukar B. Deshmukh^b

^aX-ray Crystallography Laboratory, Post-Graduate Department of Physics & Electronics, University of Jammu, Jammu Tawi 180 006, India, ^bDepartment of Chemistry, Shivaji University, Kolhapur, 416 004, India, and ^cNational Research Centre for Grapes, Pune 412 307, India
Correspondence e-mail: rkvk.paper11@gmail.com


Received 22 May 2012; accepted 1 June 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.046; wR factor = 0.112; data-to-parameter ratio = 14.0.

In the title compound, $[\text{Na}(\text{C}_{11}\text{H}_{11}\text{ClN}_3\text{O}_3)(\text{H}_2\text{O})]_n$, there are two independent Na^{I} ions, one of which lies on an inversion center and is coordinated in a slightly distorted octahedral environment. The other Na^{I} ion lies on a twofold rotation axis and is coordinated in a slightly distorted trigonal-bipyramidal coordination environment. In the organic ligand, the imidazolidine ring adopts a half-chair conformation. The Na^{I} ions bridge organic ligands and water molecules, forming a two-dimensional structure parallel to (100). There are intermolecular $\text{O}-\text{H}\cdots\text{O}$ and weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds within the two-dimensional structure.

Related literature

For background to the insecticidal applications of imidacloprid {systematic name: *N*-[1-[(6-chloro-3-pyridyl)methyl]-4,5-dihydroimidazol-2-yl]nitramide}, see: Legocki & Polec (2008); Kovganko & Kashkan (2004); Zhao *et al.* (2009); Tanner *et al.* (2010); Xu *et al.* (2010). For ring conformations, see: Duax & Norton (1975). For related structures, see: Kapoor *et al.* (2011, 2012); Kant *et al.* (2012).



Experimental

Crystal data

$[\text{Na}(\text{C}_{11}\text{H}_{11}\text{ClN}_3\text{O}_3)(\text{H}_2\text{O})]$
 $M_r = 309.68$
Monoclinic, $C2/c$
 $a = 45.655(2)$ Å
 $b = 4.9113(2)$ Å
 $c = 12.5205(7)$ Å
 $\beta = 102.184(5)^\circ$

$V = 2744.2(2)$ Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.33$ mm⁻¹
 $T = 293$ K
 $0.3 \times 0.2 \times 0.1$ mm

Data collection

Oxford Diffraction Xcalibur
Sapphire3 diffractometer
Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford
Diffraction, 2010)
 $T_{\text{min}} = 0.836$, $T_{\text{max}} = 1.000$

9498 measured reflections
2678 independent reflections
1909 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.112$
 $S = 1.02$
2678 reflections
191 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.44$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.46$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1W}-\text{H1W}\cdots\text{O12}^{\text{i}}$	0.80 (4)	2.02 (4)	2.826 (3)	179 (3)
$\text{O1W}-\text{H2W}\cdots\text{O15}^{\text{ii}}$	0.84 (4)	2.02 (4)	2.822 (2)	158 (3)
$\text{C10}-\text{H10B}\cdots\text{O12}^{\text{iii}}$	0.97	2.54	3.279 (3)	133
$\text{C13}-\text{H13B}\cdots\text{O16}^{\text{ii}}$	0.97	2.49	3.265 (3)	137

Symmetry codes: (i) $-x + 2, y + 1, -z + \frac{1}{2}$; (ii) $x, y + 1, z$; (iii) $x, -y + 1, z - \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008);

molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009).

RK acknowledges the Department of Science & Technology for the single-crystal X-ray diffractometer sanctioned as a National Facility under project No. SR/S2/CMP-47/2003. He is also thankful to the University of Jammu, Jammu, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5481).

References

- Duax, W. L. & Norton, D. A. (1975). *Atlas of Steroid Structures*, Vol. 1. New York: Plenum Press.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Kant, R., Gupta, V. K., Kapoor, K., Deshmukh, M. B. & Shripanavar, C. S. (2012). *Acta Cryst.* **E68**, o147.
- Kapoor, K., Gupta, V. K., Deshmukh, M. B., Shripanavar, C. S. & Kant, R. (2012). *Acta Cryst.* **E68**, o469.
- Kapoor, K., Gupta, V. K., Kant, R., Deshmukh, M. B. & Sripanavar, C. S. (2011). *X-ray Struct. Anal. Online*, **27**, 55–56.
- Kovganko, N. V. & Kashkan, Zh. N. (2004). *Russ. J. Org. Chem.* **40**, 1709–1726.
- Legocki, J. & Polec, I. (2008). *Pestycydy*, **1–2**, 143–159.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tanner, G. (2010). Masters Thesis "Development of a Method for the Analysis of Neonicotinoid Insecticide Residues in Honey using LCMS/MS and Investigations of Neonicotinoid Insecticides in Matrices of Importance in Apiculture", Austrian Agency for Health and Food Safety, Vienna.
- Xu, T., Wei, K.-Y., Wang, J., Ma, H.-X. & Li, J. (2010). *XUETAL J. AOAC Int.* **93**, 12–18.
- Zhao, Y., Li, Y., Wang, S. & Li, Z. (2009). *ARKIVOC*, **xi**, 152–164.

supporting information

Acta Cryst. (2012). E68, m891–m892 [https://doi.org/10.1107/S1600536812025007]

Poly[μ_2 -aqua- μ_4 -(2-{3-[(6-chloropyridin-3-yl)methyl]-2-oxoimidazolidin-1-yl}acetato)-sodium]

Rajni Kant, Vivek K. Gupta, Kamini Kapoor, Chetan S. Shripanavar, Kaushik Banerjee and Madhukar B. Deshmukh

S1. Comment

For the development of nicotinoid insecticides the crucial turning-point could be traced back to the work done by the scientists from Nihon Tokushu Noyaku Seizo K and Nippon Bayer (Legocki & Polec, 2008). Insects become resistant to insecticides due to continuous use and hence it is imperative to introduce new molecules having novel mode of action (Kovganko & Kashkan, 2004). The outstanding development of neonicotinoid insecticides has been achieved for the modern crop protection, consumer products, and animal health markets between 1990 and today reflects the enormous importance of this chemical class (Zhao *et al.*, 2009). Neonicotinoids have low toxicity toward mammals and no teratogenic or mutagenic effects (Xu *et al.*, 2010). The biological activity and agricultural uses of neonicotinoid insecticides are enormous (Zhao *et al.*, 2009). From investigations it is revealed that the neonicotinoids are converted into numerous and variable metabolites in plants as well as in mammals (Tanner *et al.*, 2010).

The asymmetric unit is shown in Fig. 1. The bond lengths and angles observed in (I) show normal values and are comparable to those in related structures (Kapoor *et al.*, 2011; Kant *et al.*, 2012). There are two independent Na¹ ions, one of which lies on an inversion center and is coordinated in a slightly distorted octahedral environment. The other Na¹ ion lies on a twofold rotation axis and is coordinated in a slightly distorted trigonal bipyramidal coordination environment. In the organic ligand the imidazole ring adopts *half-chair* conformation (asymmetry parameter: $\Delta C_2(C9-C10) = 2.31$). The Na¹ ions bridge organic ligands and solvent water molecules to form a two-dimensional structure parallel to (100). There are intermolecular O—H \cdots O and weak C—H \cdots O hydrogen bonds within the two-dimensional structure.

S2. Experimental

Ethyl[3-[(6-chloropyridin-3-yl)methyl]-2-(nitroimino)imidazolidin-1-yl]acetate (0.341 g m, 0.001 mol) was dissolved in 5 ml methanol and 5 ml 1 N NaOH solution was added. The reaction mixture was refluxed on a water bath at 343K for 12 h, and then cooled. The compound was re-precipitated upon neutralization with 1 N HCl. The compound was dissolved in methanol and crystallized in a fume hood at room temperature by the process of slow evaporation.

m.p. 575 K IR (KBr) ν_{\max} : 3421, 3300, 2872, 2930, 1668, 1606 cm⁻¹. LC—MS/MS: 270, 252, 224, 149, 126 m/z.

S3. Refinement

All H atoms except water H atoms were positioned geometrically and were treated as riding on their parent C atoms, with C—H distances of 0.93–0.97 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Water H atoms were found in a difference map and isotropically refined




Figure 1

The asymmetric unit with ellipsoids are drawn at the 40% probability level. H atoms are shown as small spheres of arbitrary radii.




Figure 2

Part of the crystal structure. The broken lines show the intermolecular C—H...O and O—H...O interactions. Only H atoms involved in hydrogen bonds have been shown.

Poly[μ_2 -aqua- μ_4 -(2-{3-[(6-chloropyridin-3-yl)methyl]-2-oxoimidazolidin-1-yl}acetato)-sodium]

Crystal data

[Na(C₁₁H₁₁ClN₃O₃)(H₂O)]

$M_r = 309.68$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 45.655$ (2) Å

$b = 4.9113$ (2) Å

$c = 12.5205$ (7) Å

$\beta = 102.184$ (5)°

$V = 2744.2$ (2) Å³

$Z = 8$

$F(000) = 1280$

$D_x = 1.499$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5286 reflections

$\theta = 3.6$ – 29.0 °

$\mu = 0.33$ mm⁻¹

$T = 293$ K

Block, white

$0.3 \times 0.2 \times 0.1$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 16.1049 pixels mm⁻¹
 ω scan
Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford Diffraction, 2010)
 $T_{\min} = 0.836$, $T_{\max} = 1.000$

9498 measured reflections
2678 independent reflections
1909 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 3.6^\circ$
 $h = -56 \rightarrow 52$
 $k = -6 \rightarrow 5$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.112$
 $S = 1.02$
2678 reflections
191 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0459P)^2 + 2.523P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. *CrysAlis PRO*, Oxford Diffraction Ltd., Version 1.171.34.40 (release 27-08-2010 *CrysAlis171.NET*) (compiled Aug 27 2010, 11:50:40) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Na1	1.0000	1.0000	0.0000	0.0267 (3)
Na2	1.0000	1.0028 (2)	0.2500	0.0286 (3)
Cl1	0.76939 (2)	0.7478 (2)	0.18991 (10)	0.0963 (4)
N1	0.81725 (5)	0.4509 (6)	0.2281 (2)	0.0652 (7)
C2	0.81005 (7)	0.2440 (7)	0.0194 (3)	0.0685 (9)
H2	0.8072	0.1756	-0.0513	0.082*
C3	0.83467 (6)	0.1665 (5)	0.0972 (2)	0.0447 (7)
C4	0.83683 (7)	0.2781 (6)	0.1992 (3)	0.0570 (8)
H4	0.8534	0.2292	0.2529	0.068*
C5	0.78956 (7)	0.4245 (8)	0.0471 (3)	0.0769 (11)
H5	0.7728	0.4802	-0.0043	0.092*
C6	0.79453 (7)	0.5174 (7)	0.1510 (3)	0.0614 (8)

C7	0.85914 (6)	-0.0155 (5)	0.0735 (2)	0.0491 (7)
H7A	0.8676	-0.1189	0.1387	0.059*
H7B	0.8505	-0.1439	0.0167	0.059*
N8	0.88292 (4)	0.1344 (4)	0.03879 (16)	0.0377 (5)
C9	0.87821 (7)	0.2620 (6)	-0.0683 (2)	0.0549 (8)
H9A	0.8602	0.3723	-0.0825	0.066*
H9B	0.8770	0.1275	-0.1257	0.066*
C10	0.90605 (6)	0.4362 (6)	-0.0581 (2)	0.0512 (7)
H10A	0.9219	0.3394	-0.0829	0.061*
H10B	0.9018	0.6046	-0.0989	0.061*
N11	0.91385 (4)	0.4864 (4)	0.05842 (16)	0.0331 (5)
C12	0.90251 (5)	0.2884 (5)	0.11303 (19)	0.0299 (5)
O12	0.90778 (4)	0.2517 (4)	0.21182 (13)	0.0437 (5)
C13	0.94038 (5)	0.6403 (5)	0.1062 (2)	0.0360 (6)
H13A	0.9397	0.6836	0.1812	0.043*
H13B	0.9401	0.8108	0.0668	0.043*
C14	0.96972 (5)	0.4954 (4)	0.10520 (16)	0.0228 (5)
O15	0.99295 (3)	0.6404 (3)	0.12325 (12)	0.0268 (4)
O16	0.96893 (3)	0.2446 (3)	0.09003 (13)	0.0302 (4)
O1W	1.03482 (4)	1.2101 (4)	0.15005 (15)	0.0364 (4)
H1W	1.0512 (7)	1.220 (6)	0.189 (3)	0.055 (9)*
H2W	1.0263 (8)	1.361 (8)	0.152 (3)	0.082 (12)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Na1	0.0343 (7)	0.0252 (6)	0.0226 (6)	-0.0001 (5)	0.0104 (5)	-0.0014 (5)
Na2	0.0387 (7)	0.0243 (6)	0.0236 (6)	0.000	0.0084 (5)	0.000
Cl1	0.0693 (6)	0.1034 (8)	0.1248 (9)	0.0315 (5)	0.0398 (6)	0.0245 (6)
N1	0.0462 (15)	0.0806 (19)	0.0682 (18)	0.0095 (14)	0.0111 (13)	0.0045 (15)
C2	0.0438 (18)	0.095 (3)	0.059 (2)	-0.0028 (18)	-0.0058 (15)	-0.0005 (19)
C3	0.0307 (13)	0.0456 (15)	0.0556 (18)	-0.0117 (12)	0.0042 (12)	0.0062 (13)
C4	0.0397 (16)	0.073 (2)	0.0546 (19)	0.0058 (15)	0.0004 (13)	0.0096 (16)
C5	0.0401 (18)	0.104 (3)	0.079 (3)	0.0158 (18)	-0.0045 (17)	0.019 (2)
C6	0.0423 (17)	0.067 (2)	0.078 (2)	0.0010 (15)	0.0184 (16)	0.0143 (18)
C7	0.0412 (15)	0.0387 (14)	0.0650 (19)	-0.0123 (13)	0.0060 (13)	-0.0015 (14)
N8	0.0363 (11)	0.0375 (11)	0.0369 (12)	-0.0057 (10)	0.0024 (9)	-0.0023 (9)
C9	0.0507 (17)	0.074 (2)	0.0349 (16)	-0.0085 (15)	-0.0032 (13)	-0.0032 (14)
C10	0.0439 (15)	0.074 (2)	0.0327 (15)	-0.0043 (14)	0.0015 (12)	0.0151 (14)
N11	0.0278 (10)	0.0322 (10)	0.0386 (11)	-0.0006 (9)	0.0057 (8)	0.0036 (9)
C12	0.0242 (11)	0.0327 (12)	0.0327 (14)	0.0049 (10)	0.0057 (10)	-0.0007 (10)
O12	0.0328 (9)	0.0672 (12)	0.0308 (10)	-0.0053 (8)	0.0057 (7)	0.0033 (9)
C13	0.0331 (13)	0.0251 (11)	0.0517 (16)	0.0000 (10)	0.0133 (11)	-0.0013 (11)
C14	0.0302 (11)	0.0213 (10)	0.0173 (10)	0.0011 (10)	0.0063 (9)	0.0016 (9)
O15	0.0288 (8)	0.0230 (7)	0.0283 (8)	-0.0029 (7)	0.0056 (6)	-0.0017 (6)
O16	0.0329 (9)	0.0182 (7)	0.0410 (9)	-0.0002 (6)	0.0116 (7)	-0.0023 (6)
O1W	0.0315 (10)	0.0405 (11)	0.0347 (10)	-0.0025 (9)	0.0011 (8)	-0.0037 (8)

Geometric parameters (Å, °)

Na1—O16 ⁱ	2.3239 (14)	C5—C6	1.351 (5)
Na1—O16 ⁱⁱ	2.3239 (14)	C5—H5	0.9300
Na1—O15 ⁱⁱⁱ	2.4112 (14)	C7—N8	1.452 (3)
Na1—O15	2.4112 (14)	C7—H7A	0.9700
Na1—O1W ⁱⁱⁱ	2.4208 (18)	C7—H7B	0.9700
Na1—O1W	2.4208 (18)	N8—C12	1.373 (3)
Na1—Na2 ⁱⁱⁱ	3.1302 (2)	N8—C9	1.454 (3)
Na1—Na2	3.1302 (2)	C9—C10	1.515 (4)
Na2—O15	2.3611 (16)	C9—H9A	0.9700
Na2—O15 ^{iv}	2.3611 (16)	C9—H9B	0.9700
Na2—O1W	2.4432 (18)	C10—N11	1.448 (3)
Na2—O1W ^{iv}	2.4432 (18)	C10—H10A	0.9700
Na2—O16 ^v	2.4969 (16)	C10—H10B	0.9700
Na2—O16 ⁱⁱ	2.4968 (16)	N11—C12	1.353 (3)
Na2—Na1 ^{iv}	3.1302 (2)	N11—C13	1.446 (3)
Na2—H2W	2.58 (3)	C12—O12	1.223 (3)
Cl1—C6	1.753 (3)	C13—C14	1.519 (3)
N1—C6	1.301 (4)	C13—H13A	0.9700
N1—C4	1.336 (4)	C13—H13B	0.9700
C2—C3	1.377 (4)	C14—O16	1.246 (2)
C2—C5	1.385 (5)	C14—O15	1.257 (2)
C2—H2	0.9300	O16—Na1 ^{vi}	2.3239 (14)
C3—C4	1.374 (4)	O16—Na2 ^{vi}	2.4968 (16)
C3—C7	1.509 (4)	O1W—H1W	0.81 (3)
C4—H4	0.9300	O1W—H2W	0.84 (4)
O16 ⁱ —Na1—O16 ⁱⁱ	180.00 (5)	C6—N1—C4	115.7 (3)
O16 ⁱ —Na1—O15 ⁱⁱⁱ	83.72 (5)	C3—C2—C5	119.6 (3)
O16 ⁱⁱ —Na1—O15 ⁱⁱⁱ	96.28 (5)	C3—C2—H2	120.2
O16 ⁱ —Na1—O15	96.28 (5)	C5—C2—H2	120.2
O16 ⁱⁱ —Na1—O15	83.72 (5)	C4—C3—C2	115.8 (3)
O15 ⁱⁱⁱ —Na1—O15	180.00 (7)	C4—C3—C7	120.6 (2)
O16 ⁱ —Na1—O1W ⁱⁱⁱ	76.78 (6)	C2—C3—C7	123.5 (3)
O16 ⁱⁱ —Na1—O1W ⁱⁱⁱ	103.22 (6)	N1—C4—C3	125.7 (3)
O15 ⁱⁱⁱ —Na1—O1W ⁱⁱⁱ	88.29 (6)	N1—C4—H4	117.1
O15—Na1—O1W ⁱⁱⁱ	91.71 (6)	C3—C4—H4	117.1
O16 ⁱ —Na1—O1W	103.22 (6)	C6—C5—C2	118.2 (3)
O16 ⁱⁱ —Na1—O1W	76.78 (6)	C6—C5—H5	120.9
O15 ⁱⁱⁱ —Na1—O1W	91.71 (6)	C2—C5—H5	120.9
O15—Na1—O1W	88.29 (6)	N1—C6—C5	125.1 (3)
O1W ⁱⁱⁱ —Na1—O1W	180.00 (8)	N1—C6—Cl1	115.0 (3)
O16 ⁱ —Na1—Na2 ⁱⁱⁱ	51.96 (4)	C5—C6—Cl1	119.9 (3)
O16 ⁱⁱ —Na1—Na2 ⁱⁱⁱ	128.04 (4)	N8—C7—C3	112.9 (2)
O15 ⁱⁱⁱ —Na1—Na2 ⁱⁱⁱ	48.32 (4)	N8—C7—H7A	109.0
O15—Na1—Na2 ⁱⁱⁱ	131.68 (4)	C3—C7—H7A	109.0
O1W ⁱⁱⁱ —Na1—Na2 ⁱⁱⁱ	50.26 (4)	N8—C7—H7B	109.0

O1W—Na1—Na2 ⁱⁱⁱ	129.74 (4)	C3—C7—H7B	109.0
O16 ⁱ —Na1—Na2	128.04 (4)	H7A—C7—H7B	107.8
O16 ⁱⁱ —Na1—Na2	51.96 (4)	C12—N8—C7	119.9 (2)
O15 ⁱⁱⁱ —Na1—Na2	131.68 (4)	C12—N8—C9	109.6 (2)
O15—Na1—Na2	48.32 (4)	C7—N8—C9	121.3 (2)
O1W ⁱⁱⁱ —Na1—Na2	129.74 (4)	N8—C9—C10	102.0 (2)
O1W—Na1—Na2	50.26 (4)	N8—C9—H9A	111.4
Na2 ⁱⁱⁱ —Na1—Na2	180.00 (4)	C10—C9—H9A	111.4
O15—Na2—O15 ^{iv}	82.15 (8)	N8—C9—H9B	111.4
O15—Na2—O1W	88.91 (5)	C10—C9—H9B	111.4
O15 ^{iv} —Na2—O1W	130.33 (6)	H9A—C9—H9B	109.2
O15—Na2—O1W ^{iv}	130.33 (6)	N11—C10—C9	101.8 (2)
O15 ^{iv} —Na2—O1W ^{iv}	88.91 (6)	N11—C10—H10A	111.4
O1W—Na2—O1W ^{iv}	130.75 (10)	C9—C10—H10A	111.4
O15—Na2—O16 ^v	150.63 (6)	N11—C10—H10B	111.4
O15 ^{iv} —Na2—O16 ^v	81.11 (5)	C9—C10—H10B	111.4
O1W—Na2—O16 ^v	83.79 (6)	H10A—C10—H10B	109.3
O1W ^{iv} —Na2—O16 ^v	73.25 (6)	C12—N11—C13	122.9 (2)
O15—Na2—O16 ⁱⁱ	81.11 (5)	C12—N11—C10	110.4 (2)
O15 ^{iv} —Na2—O16 ⁱⁱ	150.63 (6)	C13—N11—C10	120.80 (19)
O1W—Na2—O16 ⁱⁱ	73.25 (6)	O12—C12—N11	127.1 (2)
O1W ^{iv} —Na2—O16 ⁱⁱ	83.79 (6)	O12—C12—N8	124.4 (2)
O16 ^v —Na2—O16 ⁱⁱ	123.21 (8)	N11—C12—N8	108.5 (2)
O15—Na2—Na1 ^{iv}	129.79 (5)	N11—C13—C14	114.54 (18)
O15 ^{iv} —Na2—Na1 ^{iv}	49.71 (3)	N11—C13—H13A	108.6
O1W—Na2—Na1 ^{iv}	130.64 (5)	C14—C13—H13A	108.6
O1W ^{iv} —Na2—Na1 ^{iv}	49.63 (4)	N11—C13—H13B	108.6
O16 ^v —Na2—Na1 ^{iv}	47.14 (3)	C14—C13—H13B	108.6
O16 ⁱⁱ —Na2—Na1 ^{iv}	133.19 (4)	H13A—C13—H13B	107.6
O15—Na2—Na1	49.71 (3)	O16—C14—O15	125.69 (19)
O15 ^{iv} —Na2—Na1	129.79 (5)	O16—C14—C13	117.84 (18)
O1W—Na2—Na1	49.63 (4)	O15—C14—C13	116.43 (18)
O1W ^{iv} —Na2—Na1	130.65 (5)	C14—O15—Na2	122.53 (12)
O16 ^v —Na2—Na1	133.19 (4)	C14—O15—Na1	121.41 (12)
O16 ⁱⁱ —Na2—Na1	47.14 (3)	Na2—O15—Na1	81.97 (5)
Na1 ^{iv} —Na2—Na1	179.49 (4)	C14—O16—Na1 ^{vi}	125.73 (13)
O15—Na2—H2W	101.9 (8)	C14—O16—Na2 ^{vi}	110.77 (13)
O15 ^{iv} —Na2—H2W	145.3 (8)	Na1 ^{vi} —O16—Na2 ^{vi}	80.89 (5)
O1W—Na2—H2W	19.0 (8)	Na1—O1W—Na2	80.11 (5)
O1W ^{iv} —Na2—H2W	112.1 (8)	Na1—O1W—H1W	151 (2)
O16 ^v —Na2—H2W	79.3 (8)	Na2—O1W—H1W	110 (2)
O16 ⁱⁱ —Na2—H2W	62.4 (8)	Na1—O1W—H2W	100 (2)
Na1 ^{iv} —Na2—H2W	125.2 (8)	Na2—O1W—H2W	90 (2)
Na1—Na2—H2W	55.2 (8)	H1W—O1W—H2W	108 (3)
O16 ⁱ —Na1—Na2—O15	−59.31 (6)	C13—N11—C12—N8	−163.81 (19)
O16 ⁱⁱ —Na1—Na2—O15	120.69 (6)	C10—N11—C12—N8	−10.9 (3)
O15 ⁱⁱⁱ —Na1—Na2—O15	180.000 (3)	C7—N8—C12—O12	23.0 (3)

O1W ⁱⁱⁱ —Na1—Na2—O15	46.52 (8)	C9—N8—C12—O12	170.2 (2)
O1W—Na1—Na2—O15	-133.48 (8)	C7—N8—C12—N11	-155.4 (2)
O16 ⁱ —Na1—Na2—O15 ^{iv}	-39.26 (7)	C9—N8—C12—N11	-8.1 (3)
O16 ⁱⁱ —Na1—Na2—O15 ^{iv}	140.74 (7)	C12—N11—C13—C14	80.5 (3)
O15 ⁱⁱⁱ —Na1—Na2—O15 ^{iv}	-159.94 (9)	C10—N11—C13—C14	-69.7 (3)
O15—Na1—Na2—O15 ^{iv}	20.06 (9)	N11—C13—C14—O16	-17.1 (3)
O1W ⁱⁱⁱ —Na1—Na2—O15 ^{iv}	66.58 (8)	N11—C13—C14—O15	164.93 (18)
O1W—Na1—Na2—O15 ^{iv}	-113.42 (8)	O16—C14—O15—Na2	-140.66 (17)
O16 ⁱ —Na1—Na2—O1W	74.17 (8)	C13—C14—O15—Na2	37.1 (2)
O16 ⁱⁱ —Na1—Na2—O1W	-105.83 (8)	O16—C14—O15—Na1	118.07 (19)
O15 ⁱⁱⁱ —Na1—Na2—O1W	-46.52 (8)	C13—C14—O15—Na1	-64.2 (2)
O15—Na1—Na2—O1W	133.48 (8)	O15 ^{iv} —Na2—O15—C14	73.13 (14)
O1W ⁱⁱⁱ —Na1—Na2—O1W	180.000 (2)	O1W—Na2—O15—C14	-155.86 (15)
O16 ⁱ —Na1—Na2—O1W ^{iv}	-172.30 (8)	O1W ^{iv} —Na2—O15—C14	-8.69 (18)
O16 ⁱⁱ —Na1—Na2—O1W ^{iv}	7.70 (8)	O16 ^v —Na2—O15—C14	128.87 (15)
O15 ⁱⁱⁱ —Na1—Na2—O1W ^{iv}	67.01 (9)	O16 ⁱⁱ —Na2—O15—C14	-82.65 (15)
O15—Na1—Na2—O1W ^{iv}	-112.99 (9)	Na1 ^{iv} —Na2—O15—C14	57.82 (16)
O1W ⁱⁱⁱ —Na1—Na2—O1W ^{iv}	-66.47 (14)	Na1—Na2—O15—C14	-122.29 (16)
O1W—Na1—Na2—O1W ^{iv}	113.53 (14)	O15 ^{iv} —Na2—O15—Na1	-164.57 (7)
O16 ⁱ —Na1—Na2—O16 ^v	81.15 (10)	O1W—Na2—O15—Na1	-33.57 (6)
O16 ⁱⁱ —Na1—Na2—O16 ^v	-98.85 (10)	O1W ^{iv} —Na2—O15—Na1	113.60 (7)
O15 ⁱⁱⁱ —Na1—Na2—O16 ^v	-39.54 (8)	O16 ^v —Na2—O15—Na1	-108.84 (10)
O15—Na1—Na2—O16 ^v	140.46 (8)	O16 ⁱⁱ —Na2—O15—Na1	39.65 (4)
O1W ⁱⁱⁱ —Na1—Na2—O16 ^v	-173.02 (8)	Na1 ^{iv} —Na2—O15—Na1	-179.885 (9)
O1W—Na1—Na2—O16 ^v	6.98 (8)	O16 ⁱ —Na1—O15—C14	-99.57 (14)
O16 ⁱ —Na1—Na2—O16 ⁱⁱ	180.000 (1)	O16 ⁱⁱ —Na1—O15—C14	80.43 (14)
O15 ⁱⁱⁱ —Na1—Na2—O16 ⁱⁱ	59.31 (6)	O1W ⁱⁱⁱ —Na1—O15—C14	-22.69 (15)
O15—Na1—Na2—O16 ⁱⁱ	-120.69 (6)	O1W—Na1—O15—C14	157.31 (15)
O1W ⁱⁱⁱ —Na1—Na2—O16 ⁱⁱ	-74.17 (8)	Na2 ⁱⁱⁱ —Na1—O15—C14	-56.62 (15)
O1W—Na1—Na2—O16 ⁱⁱ	105.83 (8)	Na2—Na1—O15—C14	123.38 (15)
C5—C2—C3—C4	0.2 (4)	O16 ⁱ —Na1—O15—Na2	137.04 (5)
C5—C2—C3—C7	-176.3 (3)	O16 ⁱⁱ —Na1—O15—Na2	-42.96 (5)
C6—N1—C4—C3	-0.6 (5)	O1W ⁱⁱⁱ —Na1—O15—Na2	-146.07 (6)
C2—C3—C4—N1	0.3 (5)	O1W—Na1—O15—Na2	33.93 (6)
C7—C3—C4—N1	176.9 (3)	Na2 ⁱⁱⁱ —Na1—O15—Na2	180.000 (2)
C3—C2—C5—C6	-0.4 (5)	O15—C14—O16—Na1 ^{vi}	-35.2 (3)
C4—N1—C6—C5	0.4 (5)	C13—C14—O16—Na1 ^{vi}	147.11 (16)
C4—N1—C6—Cl1	-179.1 (2)	O15—C14—O16—Na2 ^{vi}	58.5 (2)
C2—C5—C6—N1	0.1 (6)	C13—C14—O16—Na2 ^{vi}	-119.20 (17)
C2—C5—C6—Cl1	179.6 (3)	O16 ⁱ —Na1—O1W—Na2	-128.89 (6)
C4—C3—C7—N8	-87.5 (3)	O16 ⁱⁱ —Na1—O1W—Na2	51.11 (6)
C2—C3—C7—N8	88.8 (3)	O15 ⁱⁱⁱ —Na1—O1W—Na2	147.16 (6)
C3—C7—N8—C12	71.7 (3)	O15—Na1—O1W—Na2	-32.83 (6)
C3—C7—N8—C9	-71.6 (3)	Na2 ⁱⁱⁱ —Na1—O1W—Na2	180.000 (1)
C12—N8—C9—C10	22.3 (3)	O15—Na2—O1W—Na1	33.61 (5)
C7—N8—C9—C10	169.0 (2)	O15 ^{iv} —Na2—O1W—Na1	112.35 (8)
N8—C9—C10—N11	-26.8 (3)	O1W ^{iv} —Na2—O1W—Na1	-113.33 (6)
C9—C10—N11—C12	24.0 (3)	O16 ^v —Na2—O1W—Na1	-174.89 (6)

C9—C10—N11—C13	177.6 (2)	O16 ⁱⁱ —Na2—O1W—Na1	-47.43 (5)
C13—N11—C12—O12	17.9 (3)	Na1 ^{iv} —Na2—O1W—Na1	179.44 (4)
C10—N11—C12—O12	170.9 (2)		

Symmetry codes: (i) $-x+2, -y+1, -z$; (ii) $x, y+1, z$; (iii) $-x+2, -y+2, -z$; (iv) $-x+2, y, -z+1/2$; (v) $-x+2, y+1, -z+1/2$; (vi) $x, y-1, z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H1W \cdots O12 ^v	0.80 (4)	2.02 (4)	2.826 (3)	179 (3)
O1W—H2W \cdots O15 ⁱⁱ	0.84 (4)	2.02 (4)	2.822 (2)	158 (3)
C10—H10B \cdots O12 ^{vii}	0.97	2.54	3.279 (3)	133
C13—H13B \cdots O16 ⁱⁱ	0.97	2.49	3.265 (3)	137

Symmetry codes: (ii) $x, y+1, z$; (v) $-x+2, y+1, -z+1/2$; (vii) $x, -y+1, z-1/2$.