

Potassium *N*-bromo-2-nitrobenzene-sulfonamide monohydrate

B. Thimme Gowda,^{a*} Sabine Foro^b and H. S. Spandana^a

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 4 October 2012; accepted 8 October 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.053; wR factor = 0.145; data-to-parameter ratio = 14.7.

In the title compound, $\text{K}^+\cdot\text{C}_6\text{H}_4\text{BrN}_2\text{O}_4\text{S}^-\cdot\text{H}_2\text{O}$, the K^+ ion is hepta-coordinated by two O atoms from two different water molecules, three sulfonyl O atoms from three *N*-bromo-2-nitrobenzenesulfonamide anions and two nitro O atoms from two *N*-bromo-2-nitrobenzenesulfonamide anions. The S–N distance of 1.576 (4) Å is consistent with an S=N double bond. The crystal structure is stabilized by intermolecular O–H···N and O–H···Br hydrogen bonds which link the molecules into polymeric layers running parallel to the *bc* plane.

Related literature

For the preparation of metal salts of *N*-haloarylsulfonamides, see: Gowda & Mahadevappa (1983); Usha & Gowda (2006). For studies on the effect of substituents and metal ions on the structures of *N*-haloarylsulfonamides, see: George *et al.* (2000); Gowda *et al.* (2011a,b); Olmstead & Power (1986). For positioning of water H atoms, see: Nardelli (1999).



Experimental

Crystal data

$\text{K}^+\cdot\text{C}_6\text{H}_4\text{BrN}_2\text{O}_4\text{S}^-\cdot\text{H}_2\text{O}$

$M_r = 337.20$

Monoclinic, $P2_1/c$
 $a = 13.034(2)\text{ \AA}$
 $b = 12.815(2)\text{ \AA}$
 $c = 6.7741(9)\text{ \AA}$
 $\beta = 100.65(1)^\circ$
 $V = 1112.0(3)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 4.27\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.48 \times 0.48 \times 0.24\text{ mm}$

Data collection

Oxford Diffraction Xcalibur diffractometer with Sapphire CCD detector
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009
 $T_{\min} = 0.234$, $T_{\max} = 0.428$
3896 measured reflections
2236 independent reflections
1847 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.145$
 $S = 1.06$
2236 reflections
152 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.79\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5–H51···N1 ⁱ	0.84 (2)	2.13 (3)	2.926 (5)	157 (5)
O5–H52···Br1 ⁱⁱ	0.84 (2)	2.85 (4)	3.509 (4)	137 (4)

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the University Grants Commission, Government of India, New Delhi, for a one-time grant to Faculty/Professors under UGC-BSR.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2509).

References

- George, E., Vivekanandan, S. & Sivakumar, K. (2000). *Acta Cryst. C* **56**, 1208–1209.
- Gowda, B. T., Foro, S. & Shakuntala, K. (2011a). *Acta Cryst. E* **67**, m926.
- Gowda, B. T., Foro, S. & Shakuntala, K. (2011b). *Acta Cryst. E* **67**, m1015.
- Gowda, B. T. & Mahadevappa, D. S. (1983). *Talanta*, **30**, 359–362.
- Nardelli, M. (1999). *J. Appl. Cryst.* **32**, 563–571.
- Olmstead, M. M. & Power, P. P. (1986). *Inorg. Chem.* **25**, 4057–4058.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Usha, K. M. & Gowda, B. T. (2006). *J. Chem. Sci.* **118**, 351–359.

supporting information

Acta Cryst. (2012). E68, m1358 [doi:10.1107/S1600536812042080]

Potassium *N*-bromo-2-nitrobenzenesulfonamidate monohydrate

B. Thimme Gowda, Sabine Foro and H. S. Spandana

S1. Comment

In the present work, to explore the effect of substituents on the crystal structures of metal salts of *N*-haloaryl-sulfonamides (George *et al.*, 2000; Gowda *et al.*, 2011*a,b*; Olmstead & Power, 1986), the crystal structure of potassium *N*-bromo-2-nitro-benzenesulfonamidate monohydrate (**I**) has been determined (Fig. 1). The structure of (**I**) resembles those of potassium *N*-bromo-2-chloro-benzenesulfonamidate sesquihydrate (**II**) (Gowda *et al.*, 2011*a*), potassium *N*-bromo-2-methyl-benzenesulfonamidate sesquihydrate (**III**) (Gowda *et al.*, 2011*b*), and sodium *N*-chloro-aryl-sulfonamides (George *et al.*, 2000; Olmstead & Power, 1986).

In the title compound (**I**), the K^+ ion is hepta coordinated by two O atoms from two different water molecules, three sulfonyl O atoms from three *N*-bromo-2-nitro-benzenesulfonamidate anions and two nitro O atoms from two *N*-bromo-2-nitro-benzenesulfonamidate anions (Fig. 2.). This is in contrast to K^+ ion hepta coordination by three O atoms from water molecules and by four sulfonyl O atoms of three *N*-bromo-2-chloro-benzenesulfonamide anions in (**II**) and three *N*-bromo-2-methyl-benzenesulfonamide anions in (**III**).

The S—N distance of 1.576 (4) Å in (**I**) is consistent with an S—N double bond and is in agreement with the observed values of 1.582 (4) Å in (**II**) and 1.577 (5) Å in (**III**).

The packing diagram consists of a two-dimensional polymeric layer running parallel to the *bc* plane (Fig. 2). The molecular packing is stabilized by O5—H51···N1 and O5—H52···Br1 hydrogen bonds (Table 1).

S2. Experimental

The title compound was prepared by a method similar to the one described by Gowda & Mahadevappa (Gowda & Mahadevappa, 1983) and Usha & Gowda (Usha & Gowda, 2006). 2 g of 2-nitrobenzenesulfonamide was dissolved with stirring in 40 ml of 5*M* KOH at room temperature. The resultant solution was cooled in ice and 4 ml of liquid bromine was added drop wise with constant stirring. The resultant potassium salt of *N*-bromo-2-nitro-benzenesulfonamidate monohydrate was filtered under suction, washed quickly with a minimum quantity of ice cold water. The purity of the compound was checked by determining its melting point (175–177°C) and estimating, iodometrically, the amount of active bromine present in it. It was further characterized from its infrared spectrum. The characteristic absorptions observed are 3624.3, 3333.0, 3192.2, 2978.1, 2922.2, 2075.4, 1626.0, 1602.9, 1477.5, 1452.4, 1242.2, 1122.6, 1060.9, 937.4, 817.8, 686.7, 640.4, 578.6, 549.6, 524.6 and 470.6 cm⁻¹.

Prism like yellow single crystals of the title compound used in the X-ray diffraction studies were obtained from its aqueous solution at room temperature.

S3. Refinement

H atoms bonded to C were positioned with idealized geometry using a riding model with the aromatic C—H = 0.93 Å. The O-bound H atoms were located in a difference map and were refined with restrained geometry (Nardelli, 1999), *viz.*

O—H distances were restrained to 0.85 (2) Å and the H—H distance was restrained to 1.365 Å, thus leading to the angle of 107°. All H atoms were refined with isotropic displacement parameters set at 1.2 U_{eq} of the parent atom. The residual electron-density features are located in the region of Br1. The highest peak and the deepest hole are 0.80 and 0.84 Å from Br1, respectively.






Figure 1

Molecular structure of the title compound, showing the atom labelling scheme for the asymmetric unit and extended to show the coordination geometry for the K^+ ion. The displacement ellipsoids are drawn at the 50% probability level. The H atoms are represented as small spheres of arbitrary radii.

**Figure 2**

Bridging of potassium cations, *N*-bromo-2-nitrobenzenesulfonamide anions and water molecules in the structure of the title compound.

**Figure 3**

Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

Potassium *N*-bromo-2-nitrobenzenesulfonamide monohydrate

Crystal data



$M_r = 337.20$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 13.034(2)$ Å

$b = 12.815(2)$ Å

$c = 6.7741(9)$ Å

$\beta = 100.65(1)^\circ$

$V = 1112.0(3)$ Å³

$Z = 4$

Data collection

Oxford Diffraction Xcalibur

diffractometer with Sapphire CCD detector

Radiation source: fine-focus sealed tube

Graphite monochromator

Rotation method data acquisition using ω scans.

$F(000) = 664$

$D_x = 2.014 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2399 reflections

$\theta = 3.1\text{--}27.8^\circ$

$\mu = 4.27 \text{ mm}^{-1}$

$T = 293$ K

Prism, yellow

$0.48 \times 0.48 \times 0.24$ mm

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.234$, $T_{\max} = 0.428$

3896 measured reflections

2236 independent reflections

1847 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$
 $\theta_{\text{max}} = 26.4^\circ, \theta_{\text{min}} = 3.2^\circ$
 $h = -16 \rightarrow 12$

$k = -16 \rightarrow 11$
 $l = -6 \rightarrow 8$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.145$
 $S = 1.06$
2236 reflections
152 parameters
3 restraints
Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0909P)^2 + 0.1082P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 0.79 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -1.21 \text{ e } \text{\AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.082 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2873 (3)	-0.0123 (3)	0.3340 (6)	0.0272 (9)
C2	0.3529 (3)	0.0693 (3)	0.3079 (6)	0.0271 (8)
C3	0.4505 (3)	0.0546 (4)	0.2573 (6)	0.0346 (10)
H3	0.4930	0.1113	0.2430	0.042*
C4	0.4833 (4)	-0.0463 (4)	0.2286 (7)	0.0419 (11)
H4	0.5488	-0.0583	0.1975	0.050*
C5	0.4173 (4)	-0.1291 (4)	0.2469 (7)	0.0425 (11)
H5	0.4377	-0.1966	0.2221	0.051*
C6	0.3214 (4)	-0.1127 (3)	0.3015 (7)	0.0368 (10)
H6	0.2791	-0.1695	0.3167	0.044*
Br1	0.24804 (4)	-0.10055 (4)	0.78959 (7)	0.0470 (3)
K1	0.09562 (9)	0.13045 (8)	0.88211 (15)	0.0418 (3)
N1	0.1510 (3)	-0.1035 (3)	0.5411 (6)	0.0384 (9)
N2	0.3214 (3)	0.1792 (3)	0.3257 (5)	0.0316 (8)
O1	0.1685 (3)	0.0941 (2)	0.5295 (5)	0.0395 (8)
O2	0.0840 (3)	-0.0085 (2)	0.2371 (5)	0.0442 (8)
O3	0.2363 (3)	0.2061 (3)	0.2308 (5)	0.0454 (8)
O4	0.3826 (3)	0.2369 (3)	0.4304 (6)	0.0520 (9)
S1	0.16307 (8)	-0.00155 (7)	0.41728 (15)	0.0294 (3)
O5	-0.0180 (3)	0.2758 (3)	0.6363 (6)	0.0494 (9)

H51	-0.066 (3)	0.240 (3)	0.566 (7)	0.059*
H52	-0.044 (4)	0.326 (3)	0.689 (8)	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.027 (2)	0.0276 (19)	0.0253 (18)	0.0017 (17)	0.0016 (16)	-0.0018 (16)
C2	0.026 (2)	0.0279 (18)	0.0271 (18)	0.0015 (17)	0.0053 (16)	0.0028 (16)
C3	0.032 (2)	0.042 (2)	0.032 (2)	0.000 (2)	0.0101 (18)	0.0059 (19)
C4	0.036 (3)	0.054 (3)	0.038 (2)	0.014 (2)	0.015 (2)	0.001 (2)
C5	0.047 (3)	0.036 (2)	0.045 (3)	0.020 (2)	0.012 (2)	-0.003 (2)
C6	0.041 (3)	0.030 (2)	0.040 (2)	0.0028 (19)	0.009 (2)	-0.0020 (18)
Br1	0.0516 (4)	0.0461 (4)	0.0432 (4)	-0.0034 (2)	0.0087 (2)	0.0096 (2)
K1	0.0378 (6)	0.0498 (6)	0.0380 (5)	0.0041 (5)	0.0076 (4)	-0.0024 (5)
N1	0.036 (2)	0.034 (2)	0.047 (2)	-0.0106 (16)	0.0122 (19)	0.0010 (16)
N2	0.032 (2)	0.0279 (17)	0.0369 (18)	-0.0005 (15)	0.0100 (16)	0.0052 (15)
O1	0.046 (2)	0.0313 (15)	0.0463 (18)	-0.0007 (13)	0.0206 (16)	-0.0100 (13)
O2	0.0293 (17)	0.0527 (19)	0.0465 (18)	-0.0035 (15)	-0.0039 (15)	-0.0014 (16)
O3	0.045 (2)	0.0325 (17)	0.0564 (19)	0.0096 (15)	0.0025 (16)	0.0113 (15)
O4	0.048 (2)	0.0379 (17)	0.070 (2)	-0.0146 (16)	0.0112 (19)	-0.0130 (17)
S1	0.0254 (6)	0.0274 (5)	0.0360 (5)	-0.0033 (4)	0.0069 (4)	-0.0032 (4)
O5	0.043 (2)	0.0443 (19)	0.063 (2)	-0.0011 (16)	0.0146 (18)	0.0010 (17)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.383 (6)	K1—O2 ⁱⁱ	2.806 (3)
C1—C6	1.391 (5)	K1—O3 ⁱⁱⁱ	2.877 (4)
C1—S1	1.815 (4)	K1—O2 ⁱⁱⁱ	3.018 (4)
C2—C3	1.390 (6)	K1—O3 ⁱ	3.081 (4)
C2—N2	1.479 (5)	K1—H51	3.06 (5)
C3—C4	1.386 (6)	N1—S1	1.576 (4)
C3—H3	0.9300	N2—O4	1.215 (5)
C4—C5	1.385 (7)	N2—O3	1.225 (5)
C4—H4	0.9300	O1—S1	1.437 (3)
C5—C6	1.384 (7)	O2—S1	1.448 (3)
C5—H5	0.9300	O2—K1 ⁱⁱ	2.806 (3)
C6—H6	0.9300	O2—K1 ^{iv}	3.018 (4)
Br1—N1	1.910 (4)	O3—K1 ^{iv}	2.877 (4)
Br1—K1	3.6829 (12)	O3—K1 ^v	3.081 (4)
K1—O5	2.743 (4)	O5—K1 ^v	2.746 (4)
K1—O5 ⁱ	2.746 (4)	O5—H51	0.844 (19)
K1—O1	2.768 (3)	O5—H52	0.841 (19)
C2—C1—C6	117.1 (4)	O2 ⁱⁱⁱ —K1—O3 ⁱ	141.88 (10)
C2—C1—S1	126.2 (3)	O5—K1—Br1	133.65 (9)
C6—C1—S1	116.7 (3)	O5 ⁱ —K1—Br1	145.95 (9)
C1—C2—C3	123.0 (4)	O1—K1—Br1	55.66 (7)
C1—C2—N2	121.5 (4)	O2 ⁱⁱ —K1—Br1	87.19 (8)

C3—C2—N2	115.4 (4)	O3 ⁱⁱⁱ —K1—Br1	97.38 (7)
C4—C3—C2	118.7 (4)	O2 ⁱⁱⁱ —K1—Br1	76.67 (7)
C4—C3—H3	120.6	O3 ⁱ —K1—Br1	96.72 (7)
C2—C3—H3	120.6	O5—K1—H51	15.6 (6)
C5—C4—C3	119.3 (4)	O5 ⁱ —K1—H51	81.7 (9)
C5—C4—H4	120.4	O1—K1—H51	77.0 (10)
C3—C4—H4	120.4	O2 ⁱⁱ —K1—H51	68.0 (7)
C6—C5—C4	120.9 (4)	O3 ⁱⁱⁱ —K1—H51	132.2 (6)
C6—C5—H5	119.5	O2 ⁱⁱⁱ —K1—H51	134.5 (10)
C4—C5—H5	119.5	O3 ⁱ —K1—H51	80.1 (8)
C5—C6—C1	120.9 (4)	Br1—K1—H51	125.1 (8)
C5—C6—H6	119.5	S1—N1—Br1	109.8 (2)
C1—C6—H6	119.5	O4—N2—O3	124.7 (4)
N1—Br1—K1	82.87 (12)	O4—N2—C2	117.7 (4)
O5—K1—O5 ⁱ	77.92 (8)	O3—N2—C2	117.6 (3)
O5—K1—O1	79.82 (11)	S1—O1—K1	127.53 (18)
O5 ⁱ —K1—O1	157.72 (11)	S1—O2—K1 ⁱⁱ	134.68 (19)
O5—K1—O2 ⁱⁱ	82.86 (11)	S1—O2—K1 ^{iv}	120.16 (18)
O5 ⁱ —K1—O2 ⁱⁱ	84.62 (11)	K1 ⁱⁱ —O2—K1 ^{iv}	105.16 (10)
O1—K1—O2 ⁱⁱ	93.37 (11)	N2—O3—K1 ^{iv}	135.6 (3)
O5—K1—O3 ⁱⁱⁱ	117.39 (11)	N2—O3—K1 ^v	123.6 (3)
O5 ⁱ —K1—O3 ⁱⁱⁱ	70.94 (11)	K1 ^{iv} —O3—K1 ^v	100.02 (11)
O1—K1—O3 ⁱⁱⁱ	119.80 (11)	O1—S1—O2	117.1 (2)
O2 ⁱⁱ —K1—O3 ⁱⁱⁱ	142.64 (11)	O1—S1—N1	115.2 (2)
O5—K1—O2 ⁱⁱⁱ	141.57 (11)	O2—S1—N1	105.8 (2)
O5 ⁱ —K1—O2 ⁱⁱⁱ	69.28 (11)	O1—S1—C1	105.67 (19)
O1—K1—O2 ⁱⁱⁱ	131.59 (10)	O2—S1—C1	105.7 (2)
O2 ⁱⁱ —K1—O2 ⁱⁱⁱ	74.84 (10)	N1—S1—C1	106.5 (2)
O3 ⁱⁱⁱ —K1—O2 ⁱⁱⁱ	70.30 (9)	K1—O5—K1 ^v	112.63 (13)
O5—K1—O3 ⁱ	67.90 (10)	K1—O5—H51	104 (4)
O5 ⁱ —K1—O3 ⁱ	109.52 (11)	K1 ^v —O5—H51	108 (4)
O1—K1—O3 ⁱ	60.50 (9)	K1—O5—H52	118 (4)
O2 ⁱⁱ —K1—O3 ⁱ	143.02 (10)	K1 ^v —O5—H52	104 (4)
O3 ⁱⁱⁱ —K1—O3 ⁱ	73.50 (8)	H51—O5—H52	110 (3)
C6—C1—C2—C3	-1.9 (6)	O4—N2—O3—K1 ^{iv}	156.5 (3)
S1—C1—C2—C3	175.3 (3)	C2—N2—O3—K1 ^{iv}	-22.1 (5)
C6—C1—C2—N2	176.0 (4)	O4—N2—O3—K1 ^v	-35.7 (5)
S1—C1—C2—N2	-6.8 (6)	C2—N2—O3—K1 ^v	145.7 (3)
C1—C2—C3—C4	1.0 (6)	K1—O1—S1—O2	102.7 (3)
N2—C2—C3—C4	-177.0 (4)	K1—O1—S1—N1	-22.6 (3)
C2—C3—C4—C5	1.3 (7)	K1—O1—S1—C1	-139.9 (2)
C3—C4—C5—C6	-2.8 (7)	K1 ⁱⁱ —O2—S1—O1	-114.6 (3)
C4—C5—C6—C1	1.8 (8)	K1 ^{iv} —O2—S1—O1	65.8 (3)
C2—C1—C6—C5	0.5 (7)	K1 ⁱⁱ —O2—S1—N1	15.4 (3)
S1—C1—C6—C5	-177.0 (4)	K1 ^{iv} —O2—S1—N1	-164.3 (2)
N1—Br1—K1—O5	-28.49 (17)	K1 ⁱⁱ —O2—S1—C1	128.1 (3)
N1—Br1—K1—O5 ⁱ	124.97 (19)	K1 ^{iv} —O2—S1—C1	-51.6 (2)

N1—Br1—K1—O1	−47.15 (14)	Br1—N1—S1—O1	−46.5 (3)
N1—Br1—K1—O2 ⁱⁱ	48.87 (14)	Br1—N1—S1—O2	−177.5 (2)
N1—Br1—K1—O3 ⁱⁱⁱ	−168.38 (14)	Br1—N1—S1—C1	70.3 (2)
N1—Br1—K1—O2 ⁱⁱⁱ	123.95 (13)	C2—C1—S1—O1	−24.4 (4)
N1—Br1—K1—O3 ⁱ	−94.22 (13)	C6—C1—S1—O1	152.8 (3)
K1—Br1—N1—S1	63.0 (2)	C2—C1—S1—O2	100.4 (4)
C1—C2—N2—O4	131.2 (4)	C6—C1—S1—O2	−82.4 (4)
C3—C2—N2—O4	−50.8 (5)	C2—C1—S1—N1	−147.4 (4)
C1—C2—N2—O3	−50.2 (5)	C6—C1—S1—N1	29.8 (4)
C3—C2—N2—O3	127.9 (4)	O5 ⁱ —K1—O5—K1 ^v	140.66 (16)
O5—K1—O1—S1	−120.6 (3)	O1—K1—O5—K1 ^v	−38.60 (13)
O5 ⁱ —K1—O1—S1	−122.5 (3)	O2 ⁱⁱ —K1—O5—K1 ^v	−133.34 (15)
O2 ⁱⁱ —K1—O1—S1	−38.5 (3)	O3 ⁱⁱⁱ —K1—O5—K1 ^v	79.80 (16)
O3 ⁱⁱⁱ —K1—O1—S1	123.5 (2)	O2 ⁱⁱⁱ —K1—O5—K1 ^v	172.23 (12)
O2 ⁱⁱⁱ —K1—O1—S1	34.2 (3)	O3 ⁱ —K1—O5—K1 ^v	23.55 (11)
O3 ⁱ —K1—O1—S1	169.1 (3)	Br1—K1—O5—K1 ^v	−54.17 (18)
Br1—K1—O1—S1	45.8 (2)		

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x, -y, -z+1$; (iii) $x, y, z+1$; (iv) $x, y, z-1$; (v) $x, -y+1/2, z-1/2$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O5—H51 ⁱⁱ —N1 ⁱⁱ	0.84 (2)	2.13 (3)	2.926 (5)	157 (5)
O5—H52 ^{vi} —Br1 ^{vi}	0.84 (2)	2.85 (4)	3.509 (4)	137 (4)

Symmetry codes: (ii) $-x, -y, -z+1$; (vi) $-x, y+1/2, -z+3/2$.