

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tris[4-(dimethylamino)pyridinium] hexakis(thiocyanato- κN)ferrate(III) monohydrate

Susanne Wöhlert,* Inke Jess and Christian Näther

Institut für Anorganische Chemie, Christian-Albrechts-Universität Kiel, Max-Eyth-Strasse 2, 24118 Kiel, Germany

Correspondence e-mail: swoehlert@ac.uni-kiel.de


Received 13 November 2012; accepted 3 December 2012

 Key indicators: single-crystal X-ray study; $T = 180$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.042; wR factor = 0.104; data-to-parameter ratio = 17.1.

In the title compound, $(\text{C}_7\text{H}_{11}\text{N}_2)_3[\text{Fe}(\text{NCS})_6] \cdot \text{H}_2\text{O}$, the Fe^{III} cation is coordinated by six terminal N -bonded thiocyanate anions into a discrete threefold negatively charged complex. Charge balance is achieved by three protonated 4-(dimethylamino)pyridine cations. The asymmetric unit consists of one Fe^{III} cation, six thiocyanate anions, three 4-(dimethylamino)pyridinium cations and one water molecule, all of them located in general positions.

Related literature

For general background to our work on the synthesis and characterization of coordination compounds based on transition metal thiocyanates and neutral N -donor co-ligands such as pyridine, see: Boeckmann & Näther (2011, 2012).



Experimental

Crystal data

$(\text{C}_7\text{H}_{11}\text{N}_2)_3[\text{Fe}(\text{NCS})_6] \cdot \text{H}_2\text{O}$
 $M_r = 791.88$
 Triclinic, $P\bar{1}$
 $a = 11.5780$ (7) Å
 $b = 11.7620$ (7) Å
 $c = 16.5450$ (11) Å
 $\alpha = 81.260$ (7)°
 $\beta = 71.550$ (7)°

$\gamma = 62.950$ (6)°
 $V = 1903.4$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.77$ mm⁻¹
 $T = 180$ K
 $0.13 \times 0.08 \times 0.06$ mm

Data collection

Stoe IPDS-1 diffractometer
 Absorption correction: numerical
 (X -SHAPE and X -RED32; Stoe & Cie, 2008)
 $T_{\min} = 0.808$, $T_{\max} = 0.947$

13617 measured reflections
 7356 independent reflections
 4936 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.104$
 $S = 0.95$
 7356 reflections

431 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34$ e Å⁻³
 $\Delta\rho_{\min} = -0.50$ e Å⁻³

Data collection: X -AREA (Stoe & Cie, 2008); cell refinement: X -AREA; data reduction: X -AREA; program(s) used to solve structure: $SHELXS97$ (Sheldrick, 2008); program(s) used to refine structure: $SHELXL97$ (Sheldrick, 2008); molecular graphics: XP in $SHELXTL$ (Sheldrick, 2008) and $DIAMOND$ (Brandenburg, 2012); software used to prepare material for publication: $XCIF$ in $SHELXTL$.

We gratefully acknowledge financial support by the State of Schleswig-Holstein and the Deutsche Forschungsgemeinschaft (project 720/3-1). We thank Professor Dr Wolfgang Bensch for access to his experimental facility.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HP2051).

References

- Boeckmann, J. & Näther, C. (2011). *Dalton Trans.* **39**, 11019–11026.
 Boeckmann, J. & Näther, C. (2012). *Polyhedron*, **31**, 587–595.
 Brandenburg, K. (2012). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stoe & Cie (2008). X -AREA, X -RED32 and X -SHAPE. Stoe & Cie, Darmstadt, Germany.

supporting information

Acta Cryst. (2013). E69, m33 [https://doi.org/10.1107/S1600536812049574]

Tris[4-(dimethylamino)pyridinium] hexakis(thiocyanato- κ N)ferrate(III) monohydrate

Susanne Wöhlert, Inke Jess and Christian Näther

S1. Comment

Recently we have reported on the synthesis and characterization of coordination compounds based on transition metal thiocyanates and neutral N-donor co-ligands like e.g. pyridine (Boeckmann & Näther, 2011, 2012). To investigate the influence of the co-ligand we tried to prepare similar compounds with N,N'-dimethylaminopyridine. Therefore, we have reacted iron(II) chloride tetrahydrate with potassium thiocyanate and N,N'-dimethylaminopyridine in water, which lead to the formation of single crystals of the title compound by accident. To identify the product of this reaction a single crystal structure determination was performed.


The crystal structure of the title compound consists of discrete $[\text{Fe}(\text{NCS})_6]^{3-}$ anions, three N,N'-dimethylamino-pyridinium cations as well as one water molecule (Fig. 1). In the discrete complexes the iron(III) cations are coordinated by six thiocyanato anions within slightly distorted octahedra with distances in the range of 2.052 (2) Å to 2.079 (2) Å and angles ranging from 87.92 (10)° to 91.80 (10)° and from 178.35 (10)° to 179.11 (10)° (Tab. 1). The building blocks are connected via intermolecular N—H \cdots S, N—H \cdots O and O—H \cdots S hydrogen bonding, in which the protonated N atom is involved (Tab. 2). These blocks are elongated in the direction of the crystallographic *a*-axis (Fig. 2).

S2. Experimental


$\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ and N,N'-dimethylaminopyridine were obtained from Sigma Aldrich. KNCS are obtained from Alfa Aesar. 0.3 mmol (59.6 mg) $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$, 0.6 mmol (58.3 mg) KNCS and 0.15 mmol (18.3 mg) dimethylaminopyridine were reacted with 1 mL H_2O in a snap cap vial. After one week red colored block-shaped single crystals of the title compound were obtained.

S3. Refinement

All C-H and N-H H atoms were located in difference map but were positioned with idealized geometry (methyl H atoms allowed to rotate but not to tip) and were refined isotropic with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$ (1.5 for the methyl H atoms) using a riding model with $C_{\text{aromatic}} = 0.95$ Å, $C_{\text{methyl}} = 0.98$ Å and N—H = 0.88 Å. The O-H H were located in difference map, their bond lengths were set to ideal values and finally they were refined isotropic with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$ using a riding model with O-H = 0.84 Å.

**Figure 1**

Crystal structure of the title compound with labeling and displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

Crystal structure of the title compound with view along the crystallographic *c*-axis. Intermolecular hydrogen bonding is shown as dashed lines.

Tris[4-(dimethylamino)pyridinium] hexakis(thiocyanato- κ N)ferrate(III) monohydrate

Crystal data

 $(C_7H_{11}N_2)_3[Fe(NCS)_6] \cdot H_2O$ $M_r = 791.88$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 11.5780$ (7) Å $b = 11.7620$ (7) Å $c = 16.5450$ (11) Å $\alpha = 81.260$ (7)° $\beta = 71.550$ (7)° $\gamma = 62.950$ (6)° $V = 1903.4$ (2) Å³ $Z = 2$ $F(000) = 822$ $D_x = 1.382$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 13617 reflections

 $\theta = 2.4$ – 26.0 ° $\mu = 0.77$ mm⁻¹ $T = 180$ K

Block, red

 $0.13 \times 0.08 \times 0.06$ mm

Data collection

Stoe IPDS-1

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi scan

Absorption correction: numerical

 $(X\text{-SHAPE}$ and $X\text{-RED32}$; Stoe & Cie, 2008) $T_{\min} = 0.808$, $T_{\max} = 0.947$

13617 measured reflections

7356 independent reflections

4936 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.075$ $\theta_{\max} = 26.0$ °, $\theta_{\min} = 2.4$ ° $h = -14 \rightarrow 14$ $k = -14 \rightarrow 14$ $l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.042$ $wR(F^2) = 0.104$ $S = 0.95$

7356 reflections

431 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.052P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.34$ e Å⁻³ $\Delta\rho_{\min} = -0.50$ e Å⁻³Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0045 (11)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.98269 (4)	0.75134 (4)	0.75439 (2)	0.01991 (12)
N1	1.1616 (3)	0.7621 (3)	0.68821 (16)	0.0302 (6)

C1	1.2706 (3)	0.7490 (3)	0.65317 (18)	0.0254 (6)
S1	1.42277 (9)	0.72966 (11)	0.60257 (6)	0.0479 (2)
N2	0.9072 (2)	0.8295 (2)	0.65142 (16)	0.0281 (5)
C2	0.8775 (3)	0.8660 (3)	0.58809 (18)	0.0226 (6)
S2	0.83393 (8)	0.91910 (8)	0.50092 (5)	0.0360 (2)
N3	0.8051 (3)	0.7391 (3)	0.82393 (16)	0.0291 (6)
C3	0.6995 (3)	0.7518 (3)	0.86786 (17)	0.0250 (6)
S3	0.55149 (8)	0.77201 (10)	0.93009 (5)	0.0430 (2)
N4	1.0601 (2)	0.5739 (2)	0.70384 (16)	0.0276 (5)
C4	1.0898 (3)	0.4916 (3)	0.65919 (17)	0.0223 (6)
S4	1.12768 (8)	0.37586 (7)	0.59908 (5)	0.02920 (18)
N5	1.0629 (2)	0.6720 (2)	0.85664 (16)	0.0280 (5)
C5	1.1078 (3)	0.6329 (3)	0.91418 (18)	0.0243 (6)
S5	1.17023 (8)	0.57453 (8)	0.99515 (5)	0.0352 (2)
N6	0.9021 (3)	0.9292 (2)	0.80598 (15)	0.0279 (5)
C6	0.8648 (3)	1.0174 (3)	0.84754 (17)	0.0233 (6)
S6	0.81802 (9)	1.14057 (7)	0.90350 (5)	0.0349 (2)
N10	0.8102 (3)	0.6969 (2)	1.06431 (16)	0.0300 (6)
H10	0.8080	0.6393	1.0374	0.036*
C11	0.8352 (3)	0.6680 (3)	1.14068 (19)	0.0287 (6)
H11	0.8516	0.5851	1.1642	0.034*
C12	0.8376 (3)	0.7546 (3)	1.18468 (17)	0.0232 (6)
H12	0.8539	0.7326	1.2390	0.028*
C13	0.8157 (3)	0.8783 (3)	1.14994 (16)	0.0198 (5)
C14	0.7897 (3)	0.9043 (3)	1.06899 (18)	0.0289 (6)
H14	0.7731	0.9860	1.0432	0.035*
C15	0.7884 (3)	0.8131 (3)	1.02853 (19)	0.0310 (7)
H15	0.7719	0.8313	0.9742	0.037*
C16	0.8466 (3)	0.9423 (3)	1.27260 (18)	0.0295 (6)
H16A	0.7616	0.9703	1.3185	0.044*
H16B	0.8965	0.9894	1.2756	0.044*
H16C	0.9012	0.8507	1.2789	0.044*
C17	0.7972 (4)	1.0927 (3)	1.1516 (2)	0.0345 (7)
H17A	0.8679	1.0823	1.0977	0.052*
H17B	0.8018	1.1450	1.1902	0.052*
H17C	0.7085	1.1347	1.1407	0.052*
N11	0.8174 (2)	0.9670 (2)	1.19072 (15)	0.0252 (5)
N20	0.8066 (3)	0.2006 (3)	0.55337 (16)	0.0324 (6)
H20	0.8113	0.1424	0.5231	0.039*
C21	0.7879 (3)	0.3170 (3)	0.51934 (19)	0.0337 (7)
H21	0.7819	0.3344	0.4625	0.040*
C22	0.7775 (3)	0.4096 (3)	0.56454 (19)	0.0290 (6)
H22	0.7631	0.4915	0.5395	0.035*
C23	0.7880 (3)	0.3847 (3)	0.64942 (17)	0.0221 (6)
C24	0.8093 (3)	0.2608 (3)	0.68218 (18)	0.0247 (6)
H24	0.8173	0.2392	0.7385	0.030*
C25	0.8183 (3)	0.1721 (3)	0.6331 (2)	0.0299 (7)
H25	0.8331	0.0889	0.6556	0.036*

C26	0.7592 (4)	0.6008 (3)	0.6599 (2)	0.0384 (8)
H26A	0.8331	0.5927	0.6081	0.058*
H26B	0.7588	0.6519	0.7017	0.058*
H26C	0.6729	0.6429	0.6458	0.058*
C27	0.8000 (3)	0.4446 (3)	0.78040 (19)	0.0327 (7)
H27A	0.7285	0.4235	0.8192	0.049*
H27B	0.7979	0.5188	0.8020	0.049*
H27C	0.8880	0.3716	0.7768	0.049*
N21	0.7785 (2)	0.4741 (2)	0.69564 (15)	0.0253 (5)
N30	0.4772 (3)	0.3824 (4)	0.7836 (2)	0.0598 (10)
H30	0.4722	0.4477	0.8067	0.072*
C31	0.5092 (4)	0.2694 (5)	0.8242 (3)	0.0573 (12)
H31	0.5254	0.2614	0.8781	0.069*
C32	0.5190 (3)	0.1663 (4)	0.7905 (2)	0.0459 (9)
H32	0.5446	0.0866	0.8199	0.055*
C33	0.4912 (3)	0.1771 (3)	0.7116 (2)	0.0328 (7)
C34	0.4594 (3)	0.2978 (3)	0.6703 (2)	0.0399 (8)
H34	0.4428	0.3095	0.6162	0.048*
C35	0.4524 (4)	0.3966 (4)	0.7073 (3)	0.0530 (10)
H35	0.4298	0.4773	0.6792	0.064*
C36	0.4696 (4)	0.0904 (4)	0.5952 (2)	0.0456 (9)
H36A	0.5426	0.1023	0.5505	0.068*
H36B	0.3832	0.1638	0.5954	0.068*
H36C	0.4665	0.0124	0.5842	0.068*
C37	0.5321 (4)	-0.0459 (4)	0.7194 (3)	0.0544 (10)
H37A	0.4772	-0.0370	0.7790	0.082*
H37B	0.6279	-0.0837	0.7172	0.082*
H37C	0.5163	-0.1014	0.6898	0.082*
N31	0.4946 (3)	0.0793 (3)	0.67822 (18)	0.0371 (6)
O1	0.4555 (3)	0.5701 (4)	0.8757 (3)	0.0986 (15)
H1O1	0.4927	0.6104	0.8866	0.148*
H2O1	0.4004	0.5649	0.9218	0.148*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0247 (2)	0.0191 (2)	0.0198 (2)	-0.01218 (16)	-0.00614 (15)	-0.00240 (14)
N1	0.0341 (14)	0.0365 (15)	0.0283 (13)	-0.0208 (12)	-0.0074 (11)	-0.0083 (11)
C1	0.0323 (16)	0.0266 (15)	0.0241 (14)	-0.0169 (12)	-0.0104 (12)	0.0002 (11)
S1	0.0299 (4)	0.0642 (6)	0.0518 (5)	-0.0269 (4)	-0.0040 (4)	-0.0013 (5)
N2	0.0338 (13)	0.0265 (13)	0.0273 (13)	-0.0148 (11)	-0.0110 (11)	0.0017 (10)
C2	0.0217 (13)	0.0208 (14)	0.0264 (15)	-0.0090 (11)	-0.0067 (11)	-0.0044 (11)
S2	0.0438 (4)	0.0392 (5)	0.0268 (4)	-0.0141 (4)	-0.0178 (3)	-0.0017 (3)
N3	0.0332 (13)	0.0349 (15)	0.0273 (13)	-0.0215 (11)	-0.0065 (11)	-0.0043 (11)
C3	0.0336 (16)	0.0305 (15)	0.0203 (13)	-0.0187 (13)	-0.0124 (12)	-0.0001 (11)
S3	0.0322 (4)	0.0695 (6)	0.0321 (4)	-0.0292 (4)	-0.0023 (3)	-0.0044 (4)
N4	0.0339 (13)	0.0223 (12)	0.0309 (13)	-0.0154 (11)	-0.0085 (10)	-0.0027 (11)
C4	0.0268 (13)	0.0198 (14)	0.0214 (13)	-0.0117 (11)	-0.0074 (11)	0.0029 (11)

S4	0.0450 (4)	0.0209 (4)	0.0222 (3)	-0.0143 (3)	-0.0089 (3)	-0.0030 (3)
N5	0.0342 (13)	0.0249 (13)	0.0293 (13)	-0.0135 (11)	-0.0136 (11)	0.0004 (10)
C5	0.0279 (14)	0.0224 (14)	0.0271 (15)	-0.0139 (12)	-0.0066 (12)	-0.0058 (12)
S5	0.0428 (4)	0.0408 (5)	0.0301 (4)	-0.0193 (4)	-0.0193 (3)	0.0007 (3)
N6	0.0359 (13)	0.0257 (13)	0.0258 (12)	-0.0173 (11)	-0.0066 (10)	-0.0022 (10)
C6	0.0322 (14)	0.0204 (14)	0.0190 (13)	-0.0138 (12)	-0.0069 (11)	0.0020 (11)
S6	0.0600 (5)	0.0223 (4)	0.0203 (4)	-0.0178 (4)	-0.0078 (3)	-0.0022 (3)
N10	0.0403 (14)	0.0309 (14)	0.0256 (13)	-0.0194 (11)	-0.0085 (10)	-0.0081 (11)
C11	0.0323 (15)	0.0252 (15)	0.0304 (15)	-0.0148 (12)	-0.0068 (12)	-0.0013 (12)
C12	0.0280 (14)	0.0235 (14)	0.0202 (13)	-0.0132 (11)	-0.0074 (11)	0.0019 (11)
C13	0.0214 (12)	0.0244 (14)	0.0160 (12)	-0.0131 (11)	-0.0026 (10)	-0.0015 (10)
C14	0.0411 (16)	0.0313 (16)	0.0210 (14)	-0.0206 (14)	-0.0116 (12)	0.0031 (12)
C15	0.0397 (16)	0.0410 (18)	0.0215 (14)	-0.0240 (14)	-0.0113 (12)	0.0018 (13)
C16	0.0463 (17)	0.0311 (16)	0.0231 (14)	-0.0233 (14)	-0.0173 (13)	0.0042 (12)
C17	0.0528 (19)	0.0267 (16)	0.0351 (17)	-0.0228 (15)	-0.0206 (15)	0.0050 (13)
N11	0.0372 (13)	0.0233 (12)	0.0227 (12)	-0.0169 (11)	-0.0136 (10)	0.0027 (10)
N20	0.0379 (14)	0.0362 (15)	0.0299 (13)	-0.0198 (12)	-0.0073 (11)	-0.0119 (11)
C21	0.0413 (17)	0.046 (2)	0.0214 (14)	-0.0245 (15)	-0.0095 (12)	-0.0028 (13)
C22	0.0389 (16)	0.0320 (16)	0.0243 (14)	-0.0217 (14)	-0.0130 (12)	0.0068 (12)
C23	0.0216 (13)	0.0240 (14)	0.0226 (13)	-0.0119 (11)	-0.0049 (10)	-0.0020 (11)
C24	0.0274 (14)	0.0264 (15)	0.0215 (13)	-0.0136 (12)	-0.0055 (11)	-0.0001 (11)
C25	0.0332 (15)	0.0250 (15)	0.0342 (16)	-0.0156 (13)	-0.0060 (12)	-0.0051 (13)
C26	0.056 (2)	0.0208 (15)	0.050 (2)	-0.0160 (15)	-0.0328 (17)	0.0054 (14)
C27	0.0446 (18)	0.0310 (16)	0.0286 (15)	-0.0173 (14)	-0.0153 (13)	-0.0037 (13)
N21	0.0325 (12)	0.0214 (12)	0.0271 (12)	-0.0126 (10)	-0.0135 (10)	-0.0003 (10)
N30	0.0410 (18)	0.078 (3)	0.065 (2)	-0.0339 (18)	0.0046 (16)	-0.029 (2)
C31	0.039 (2)	0.098 (4)	0.037 (2)	-0.035 (2)	-0.0031 (16)	-0.005 (2)
C32	0.0353 (17)	0.067 (3)	0.0351 (18)	-0.0254 (18)	-0.0102 (14)	0.0108 (17)
C33	0.0196 (13)	0.0413 (18)	0.0308 (16)	-0.0127 (13)	-0.0040 (11)	0.0096 (13)
C34	0.0363 (17)	0.0393 (19)	0.0416 (19)	-0.0166 (15)	-0.0113 (14)	0.0073 (15)
C35	0.0395 (19)	0.047 (2)	0.068 (3)	-0.0205 (18)	-0.0082 (18)	0.002 (2)
C36	0.0387 (18)	0.050 (2)	0.043 (2)	-0.0155 (17)	-0.0080 (15)	-0.0057 (17)
C37	0.043 (2)	0.037 (2)	0.069 (3)	-0.0123 (17)	-0.0146 (18)	0.0185 (19)
N31	0.0301 (13)	0.0305 (14)	0.0393 (15)	-0.0082 (11)	-0.0050 (11)	0.0044 (12)
O1	0.066 (2)	0.123 (3)	0.122 (3)	-0.065 (2)	0.020 (2)	-0.064 (3)

Geometric parameters (Å, °)

Fe1—N4	2.052 (2)	N20—H20	0.8800
Fe1—N6	2.059 (2)	C21—C22	1.353 (4)
Fe1—N2	2.061 (3)	C21—H21	0.9500
Fe1—N1	2.065 (3)	C22—C23	1.422 (4)
Fe1—N3	2.072 (3)	C22—H22	0.9500
Fe1—N5	2.079 (2)	C23—N21	1.338 (4)
N1—C1	1.157 (4)	C23—C24	1.413 (4)
C1—S1	1.618 (3)	C24—C25	1.365 (4)
N2—C2	1.165 (4)	C24—H24	0.9500
C2—S2	1.620 (3)	C25—H25	0.9500

N3—C3	1.162 (4)	C26—N21	1.461 (4)
C3—S3	1.625 (3)	C26—H26A	0.9800
N4—C4	1.164 (4)	C26—H26B	0.9800
C4—S4	1.621 (3)	C26—H26C	0.9800
N5—C5	1.162 (4)	C27—N21	1.464 (4)
C5—S5	1.634 (3)	C27—H27A	0.9800
N6—C6	1.165 (4)	C27—H27B	0.9800
C6—S6	1.620 (3)	C27—H27C	0.9800
N10—C11	1.345 (4)	N30—C31	1.339 (6)
N10—C15	1.347 (4)	N30—C35	1.351 (6)
N10—H10	0.8800	N30—H30	0.8800
C11—C12	1.353 (4)	C31—C32	1.355 (7)
C11—H11	0.9500	C31—H31	0.9500
C12—C13	1.419 (4)	C32—C33	1.415 (5)
C12—H12	0.9500	C32—H32	0.9500
C13—N11	1.336 (4)	C33—N31	1.330 (5)
C13—C14	1.423 (4)	C33—C34	1.418 (5)
C14—C15	1.356 (4)	C34—C35	1.354 (6)
C14—H14	0.9500	C34—H34	0.9500
C15—H15	0.9500	C35—H35	0.9500
C16—N11	1.454 (4)	C36—N31	1.467 (5)
C16—H16A	0.9800	C36—H36A	0.9800
C16—H16B	0.9800	C36—H36B	0.9800
C16—H16C	0.9800	C36—H36C	0.9800
C17—N11	1.466 (4)	C37—N31	1.457 (5)
C17—H17A	0.9800	C37—H37A	0.9800
C17—H17B	0.9800	C37—H37B	0.9800
C17—H17C	0.9800	C37—H37C	0.9800
N20—C25	1.344 (4)	O1—H1O1	0.8400
N20—C21	1.346 (4)	O1—H2O1	0.8401
N4—Fe1—N6	179.11 (11)	N20—C21—H21	119.3
N4—Fe1—N2	88.68 (10)	C22—C21—H21	119.3
N6—Fe1—N2	91.24 (10)	C21—C22—C23	120.1 (3)
N4—Fe1—N1	89.09 (10)	C21—C22—H22	119.9
N6—Fe1—N1	91.80 (10)	C23—C22—H22	119.9
N2—Fe1—N1	90.15 (10)	N21—C23—C24	121.8 (3)
N4—Fe1—N3	91.19 (10)	N21—C23—C22	121.6 (3)
N6—Fe1—N3	87.92 (10)	C24—C23—C22	116.6 (3)
N2—Fe1—N3	91.48 (10)	C25—C24—C23	120.1 (3)
N1—Fe1—N3	178.35 (10)	C25—C24—H24	119.9
N4—Fe1—N5	91.08 (10)	C23—C24—H24	119.9
N6—Fe1—N5	89.02 (10)	N20—C25—C24	121.1 (3)
N2—Fe1—N5	178.77 (10)	N20—C25—H25	119.5
N1—Fe1—N5	88.64 (10)	C24—C25—H25	119.5
N3—Fe1—N5	89.73 (10)	N21—C26—H26A	109.5
C1—N1—Fe1	170.0 (3)	N21—C26—H26B	109.5
N1—C1—S1	178.9 (3)	H26A—C26—H26B	109.5

C2—N2—Fe1	173.1 (2)	N21—C26—H26C	109.5
N2—C2—S2	178.9 (3)	H26A—C26—H26C	109.5
C3—N3—Fe1	168.7 (2)	H26B—C26—H26C	109.5
N3—C3—S3	178.9 (3)	N21—C27—H27A	109.5
C4—N4—Fe1	162.8 (2)	N21—C27—H27B	109.5
N4—C4—S4	178.3 (3)	H27A—C27—H27B	109.5
C5—N5—Fe1	176.9 (2)	N21—C27—H27C	109.5
N5—C5—S5	178.4 (3)	H27A—C27—H27C	109.5
C6—N6—Fe1	167.6 (2)	H27B—C27—H27C	109.5
N6—C6—S6	177.9 (3)	C23—N21—C26	120.9 (2)
C11—N10—C15	120.8 (2)	C23—N21—C27	121.1 (2)
C11—N10—H10	119.6	C26—N21—C27	117.7 (2)
C15—N10—H10	119.6	C31—N30—C35	120.3 (4)
N10—C11—C12	121.4 (3)	C31—N30—H30	119.9
N10—C11—H11	119.3	C35—N30—H30	119.9
C12—C11—H11	119.3	N30—C31—C32	121.8 (4)
C11—C12—C13	120.2 (3)	N30—C31—H31	119.1
C11—C12—H12	119.9	C32—C31—H31	119.1
C13—C12—H12	119.9	C31—C32—C33	120.1 (4)
N11—C13—C12	122.5 (2)	C31—C32—H32	120.0
N11—C13—C14	121.2 (3)	C33—C32—H32	120.0
C12—C13—C14	116.3 (2)	N31—C33—C32	121.8 (3)
C15—C14—C13	120.4 (3)	N31—C33—C34	121.9 (3)
C15—C14—H14	119.8	C32—C33—C34	116.3 (3)
C13—C14—H14	119.8	C35—C34—C33	120.4 (4)
N10—C15—C14	120.9 (3)	C35—C34—H34	119.8
N10—C15—H15	119.5	C33—C34—H34	119.8
C14—C15—H15	119.5	N30—C35—C34	121.1 (4)
N11—C16—H16A	109.5	N30—C35—H35	119.4
N11—C16—H16B	109.5	C34—C35—H35	119.4
H16A—C16—H16B	109.5	N31—C36—H36A	109.5
N11—C16—H16C	109.5	N31—C36—H36B	109.5
H16A—C16—H16C	109.5	H36A—C36—H36B	109.5
H16B—C16—H16C	109.5	N31—C36—H36C	109.5
N11—C17—H17A	109.5	H36A—C36—H36C	109.5
N11—C17—H17B	109.5	H36B—C36—H36C	109.5
H17A—C17—H17B	109.5	N31—C37—H37A	109.5
N11—C17—H17C	109.5	N31—C37—H37B	109.5
H17A—C17—H17C	109.5	H37A—C37—H37B	109.5
H17B—C17—H17C	109.5	N31—C37—H37C	109.5
C13—N11—C16	122.4 (2)	H37A—C37—H37C	109.5
C13—N11—C17	120.6 (2)	H37B—C37—H37C	109.5
C16—N11—C17	116.9 (2)	C33—N31—C37	121.7 (3)
C25—N20—C21	120.7 (3)	C33—N31—C36	121.3 (3)
C25—N20—H20	119.6	C37—N31—C36	116.8 (3)
C21—N20—H20	119.6	H1O1—O1—H2O1	105.9
N20—C21—C22	121.3 (3)		
