

## 3,4',5-Trichlorobiphenyl-4-yl 2,2,2-trichloroethyl sulfate

Hans-Joachim Lehmler,<sup>a\*</sup> Xianran He,<sup>a</sup> Michael W. Duffel<sup>b</sup> and Sean Parkin<sup>c</sup>

<sup>a</sup>The University of Iowa, Department of Occupational and Environmental Health, Iowa City, IA 52242, USA, <sup>b</sup>The University of Iowa, Department of Pharmaceutical Sciences and Experimental Therapeutics, Iowa City, IA 52242, USA, and <sup>c</sup>University of Kentucky, Department of Chemistry, Lexington, KY 40506-0055, USA  
Correspondence e-mail: hans-joachim-lehmler@uiowa.edu


Received 23 February 2013; accepted 22 March 2013

Key indicators: single-crystal X-ray study;  $T = 90\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.011\text{ \AA}$ ;  $R$  factor = 0.064;  $wR$  factor = 0.161; data-to-parameter ratio = 22.0.

Crystals of the title compound,  $\text{C}_{14}\text{H}_8\text{Cl}_6\text{O}_4\text{S}$ , are twinned by inversion, with unequal components [0.85 (3):0.15 (3)]. The asymmetric unit contains two independent molecules that are related by a pseudo-inversion center. The  $\text{C}_{\text{ar}}-\text{O}$  [1.393 (9) and 1.397 (9)  $\text{\AA}$ ] and ester  $\text{S}-\text{O}$  bond lengths [1.600 (5) and 1.590 (5)  $\text{\AA}$ ] of both molecules are comparable to the structurally related 2,3,5,5-trichlorobiphenyl-4-yl 2,2,2-trichloroethyl sulfate. The dihedral angles between the benzene rings in the two molecules are 37.8 (2) and 35.0 (2) $^{\circ}$ .

### Related literature

For related structures of biphenyl-4-yl ester 2,2,2-trichloroethyl esters of sulfuric acid, see: Li *et al.* (2008, 2010a,b,c). For a review of structures of sulfuric acid aryl mono esters, see: Brandao *et al.* (2005); Denehy *et al.* (2006). For additional background to sulfate metabolites of polychlorinated biphenyls, see: Liu *et al.* (2006, 2009); Wang *et al.* (2006); Dhakal *et al.* (2012); Zhai *et al.* (2013).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_8\text{Cl}_6\text{O}_4\text{S}$

$M_r = 484.96$

Orthorhombic,  $Pca2_1$   
 $a = 13.993 (3)\text{ \AA}$   
 $b = 9.1890 (18)\text{ \AA}$   
 $c = 28.778 (6)\text{ \AA}$   
 $V = 3700.3 (13)\text{ \AA}^3$

$Z = 8$   
Cu  $K\alpha$  radiation  
 $\mu = 9.71\text{ mm}^{-1}$   
 $T = 90\text{ K}$   
 $0.17 \times 0.09 \times 0.02\text{ mm}$

#### Data collection

Bruker X8 Proteum diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2006)  
 $T_{\min} = 0.504$ ,  $T_{\max} = 0.830$

45894 measured reflections  
6651 independent reflections  
6238 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.062$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.161$   
 $S = 1.15$   
6651 reflections  
302 parameters  
1 restraint

H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.96\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.85\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
3176 Friedel pairs  
Flack parameter: 0.15 (3)

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and local procedures.

This research was supported by grants ES05605, ES013661 and ES017425 from the National Institute of Environmental Health Sciences, NIH.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YK2088).

### References

- Brandao, T. A. S., Priebe, J. P., Damasceno, A. S., Bortoluzzi, A. J., Kirby, A. J. & Nome, F. (2005). *J. Mol. Struct.* **734**, 205–209.
- Bruker (2006). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Denehy, E., White, J. M. & Williams, S. J. (2006). *Chem. Commun.* pp. 314–316.
- Dhakal, K., He, X., Lehmler, H. J., Teesch, L. M., Duffel, M. W. & Robertson, L. W. (2012). *Chem. Res. Toxicol.* **25**, 2796–2804.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Li, X., Parkin, S., Duffel, M. W., Robertson, L. W. & Lehmler, H.-J. (2010a). *Acta Cryst. E* **66**, o1073.
- Li, X., Parkin, S., Duffel, M. W., Robertson, L. W. & Lehmler, H.-J. (2010b). *Environ. Int.* **36**, 843–848.
- Li, X., Parkin, S., Duffel, M. W., Robertson, L. W. & Lehmler, H.-J. (2010c). *Acta Cryst. E* **66**, o1615–o1616.
- Li, X., Parkin, S., Robertson, L. W. & Lehmler, H.-J. (2008). *Acta Cryst. E* **64**, o2464.
- Liu, Y., Apak, T. I., Lehmler, H.-J., Robertson, L. W. & Duffel, M. W. (2006). *Chem. Res. Toxicol.* **19**, 1420–1425.
- Liu, Y., Smart, J. T., Song, Y., Lehmler, H.-J., Robertson, L. W. & Duffel, M. W. (2009). *Drug Metab. Dispos.* **37**, 1065–1072.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, L.-Q., Lehmler, H.-J., Robertson, L. W. & James, M. O. (2006). *Chem. Biol. Interact.* **159**, 235–246.
- Zhai, G., Lehmler, H. J. & Schnoor, J. L. (2013). *Environ. Sci. Technol.* **47**, 557–562.

# supporting information

*Acta Cryst.* (2013). E69, o620 [doi:10.1107/S1600536813007976]

## 3,4',5-Trichlorobiphenyl-4-yl 2,2,2-trichloroethyl sulfate

Hans-Joachim Lehmler, Xianran He, Michael W. Duffel and Sean Parkin

### S1. Comment

Sulfuric acid monoesters of hydroxylated polychlorinated biphenyls (OHPCBs) are emerging as an important class of metabolites of polychlorinated biphenyls (PCBs). Two recent *in vivo* studies report the formation of PCB sulfates by rats (Dhakal *et al.*, 2012) and poplar plants (Zhai *et al.*, 2013). *In vitro* studies demonstrate that PCB sulfates are both substrates and inhibitors of mammalian cytosolic sulfotransferases (Liu *et al.*, 2006; Wang *et al.*, 2006; Liu *et al.*, 2009). Only limited structural information about sulfate mono- and diesters of hydroxylated PCBs is available to support structure-activity or structure-property relationship studies. Here we report the structure of the title compound, a biphenyl-4-yl 2,2,2-trichloroethyl sulfate with two chlorine substituents *ortho* to the sulfate group, to contribute to the number of available crystal structures.

The two independent molecules of the title compound in the asymmetric unit are related by a pseudo-inversion center. The length of the C<sub>aromatic</sub>—O bonds of the two molecules are 1.393 (9) and 1.397 (9) Å, respectively. These bond lengths are comparable to the C<sub>aromatic</sub>—O bond length (1.405 Å) reported for the structurally related 2',3,5,5'-trichloro-biphenyl-4-yl 2,2,2-trichloroethyl sulfate (Li *et al.*, 2010b). In contrast, biphenyl-4-yl 2,2,2-trichloroethyl sulfates without electronegative chlorine substituents *ortho* to the sulfate group have slightly longer C<sub>aromatic</sub>—O bond length ranging from 1.426 to 1.449 Å (Li *et al.*, 2008; Li *et al.*, 2010b; Li *et al.*, 2010a; Li *et al.*, 2010c).

The lengths of the PCB sulfate ester bond of the title compound (*i.e.*, S1—O1) are 1.600 (5) and 1.590 (5) Å. In contrast, biphenyl-4-yl 2,2,2-trichloroethyl sulfates without chlorine substituents *ortho* to the sulfate group typically have shorter sulfate ester bond lengths ranging from 1.563 to 1.586 Å (Li *et al.*, 2008; Li *et al.*, 2010b; Li *et al.*, 2010a; Li *et al.*, 2010c). Similar to aromatic sulfate monoesters (Branda *et al.*, 2005; Denehy *et al.*, 2006), this difference suggests that chlorine substituents *ortho* to the sulfate group decrease the stability of the S—O ester bond.

The dihedral angle of the biphenyl moiety of PCB derivatives is a structural parameter associated with the affinity of PCB derivatives for cellular target molecules. The two molecules of the title compound have solid state dihedral angles of 37.8 (2)° and 35.0 (2)°. Similarly, structurally related biphenyl-4-yl 2,2,2-trichloroethyl sulfates have dihedral angles ranging from 4.9° to 41.8° in the solid state (Li *et al.*, 2008; Li *et al.*, 2010a; Li *et al.*, 2010c). The fact that biphenyl-4-yl 2,2,2-trichloroethyl sulfates without *ortho* chlorine substituents adopt a range of dihedral angles can be explained by crystal packing effects, which force the biphenyl moiety to adopt an energetically less favorable conformation in the solid state.


### S2. Experimental

The title compound was synthesized from 3,4',5-trichlorobiphenyl-4-ol and 2,2,2-trichloroethyl sulfonyl chloride using 4-dimethylaminopyridine as catalyst as reported previously (Li *et al.*, 2008). Crystals suitable for X-ray diffraction analysis were obtained by slow evaporation of a methanolic solution.

**S3. Refinement**

H atoms were found in difference Fourier maps and subsequently placed in idealized positions with constrained distances of 0.99 Å ( $R_2\text{CH}_2$ ), 0.95 Å ( $\text{C}_{\text{sp}^2}\text{H}$ ), and with  $U_{\text{iso}}(\text{H})$  values set to either  $1.2U_{\text{eq}}$  or  $1.5U_{\text{eq}}$  ( $\text{RCH}_3$ , OH) of the attached atom.

The two independent molecules are related by a pseudo-inversion centre, which results in large correlations between the displacement parameters. In order to ensure satisfactory refinement, the displacement parameters of equivalent atoms in each molecule were constrained to be the same using the EADP command of *SHELXL97*.

**Figure 1**

View of the title compound showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**3,4',5-Trichlorobiphenyl-4-yl 2,2,2-trichloroethyl sulfate***Crystal data* $M_r = 484.96$ Orthorhombic,  $Pca2_1$ 

Hall symbol: P 2c -2ac

 $a = 13.993 (3)$  Å $b = 9.1890 (18)$  Å $c = 28.778 (6)$  Å $V = 3700.3 (13)$  Å<sup>3</sup> $Z = 8$  $F(000) = 1936$  $D_x = 1.741 \text{ Mg m}^{-3}$ Cu  $K\alpha$  radiation,  $\lambda = 1.54178$  Å

Cell parameters from 9992 reflections

 $\theta = 3.1\text{--}68.3^\circ$  $\mu = 9.71 \text{ mm}^{-1}$  $T = 90 \text{ K}$ 

Flake, colourless

 $0.17 \times 0.09 \times 0.02$  mm*Data collection*Bruker X8 Proteum  
diffractometer

Radiation source: fine-focus rotating anode

Graded multilayer optics monochromator

Detector resolution: 5.6 pixels mm<sup>-1</sup> $\varphi$  and  $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Bruker, 2006) $T_{\min} = 0.504$ ,  $T_{\max} = 0.830$ 

45894 measured reflections

6651 independent reflections

6238 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.062$  $\theta_{\max} = 68.4^\circ$ ,  $\theta_{\min} = 3.1^\circ$  $h = -14 \rightarrow 16$  $k = -10 \rightarrow 11$  $l = -34 \rightarrow 34$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.064$  $wR(F^2) = 0.161$  $S = 1.15$ 

6651 reflections

302 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0514P)^2 + 21.3733P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.96 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.85 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack (1983), 3176 Friedel pairs

Absolute structure parameter: 0.15 (3)

### Special details

**Experimental.** The crystal was twinned by inversion, but with unequal sized pieces of each component. The refined Flack parameter indicates major:minor fractions of 0.85 (3):0.15 (3).

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against all reflections. The weighted  $R$ -value  $wR$  and goodness of fit  $S$  are based on  $F^2$ . Conventional  $R$ -values  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -values based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -values based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1A	0.77740 (12)	0.4047 (2)	0.35290 (7)	0.0283 (2)
O1A	0.8439 (4)	0.2785 (5)	0.33235 (19)	0.0273 (7)
O2A	0.8158 (4)	0.4167 (6)	0.40377 (19)	0.0313 (7)
O3A	0.7985 (4)	0.5393 (6)	0.3314 (2)	0.0302 (7)
O4A	0.6838 (4)	0.3483 (6)	0.3531 (2)	0.0349 (7)
Cl1A	1.03327 (13)	0.3420 (2)	0.37524 (7)	0.0380 (3)
Cl2A	0.78298 (12)	0.2737 (2)	0.23480 (7)	0.0318 (2)
Cl3A	1.37897 (13)	0.5084 (2)	0.10448 (8)	0.0378 (3)
Cl4A	0.8960 (2)	0.4431 (3)	0.50012 (9)	0.0563 (4)
Cl5A	0.69326 (19)	0.4810 (2)	0.48583 (9)	0.0487 (4)
Cl6A	0.76595 (19)	0.2066 (2)	0.52014 (8)	0.0489 (4)
C1A	1.0641 (5)	0.3630 (7)	0.2372 (3)	0.0240 (9)
C2A	1.0822 (5)	0.3612 (8)	0.2847 (3)	0.0273 (9)
H2A	1.1456	0.3760	0.2956	0.033*
C3A	1.0090 (5)	0.3381 (8)	0.3166 (3)	0.0276 (9)
C4A	0.9162 (5)	0.3143 (8)	0.3011 (3)	0.0252 (9)
C5A	0.8987 (5)	0.3119 (7)	0.2542 (3)	0.0259 (9)
C6A	0.9704 (5)	0.3381 (8)	0.2223 (3)	0.0273 (9)
H6A	0.9560	0.3393	0.1900	0.033*
C7A	0.7997 (6)	0.2919 (9)	0.4344 (3)	0.0326 (10)
H7A1	0.8546	0.2241	0.4332	0.039*
H7A2	0.7412	0.2386	0.4251	0.039*
C8A	0.7887 (7)	0.3548 (10)	0.4822 (3)	0.0431 (12)
C1'A	1.1400 (5)	0.3932 (8)	0.2034 (3)	0.0284 (9)
C2'A	1.1233 (5)	0.4783 (9)	0.1639 (3)	0.0314 (11)
H2'A	1.0607	0.5138	0.1581	0.038*
C3'A	1.1959 (5)	0.5118 (9)	0.1333 (3)	0.0307 (10)
H3'A	1.1833	0.5691	0.1065	0.037*

C4'A	1.2877 (5)	0.4609 (9)	0.1420 (3)	0.0284 (10)
C5'A	1.3053 (5)	0.3752 (8)	0.1800 (3)	0.0293 (10)
H5'A	1.3677	0.3379	0.1850	0.035*
C6'A	1.2329 (5)	0.3424 (8)	0.2113 (3)	0.0285 (9)
H6'A	1.2464	0.2854	0.2380	0.034*
S1B	0.48992 (12)	0.0952 (2)	0.44025 (7)	0.0283 (2)
O1B	0.4243 (3)	0.2207 (5)	0.46086 (19)	0.0273 (7)
O2B	0.4512 (4)	0.0835 (6)	0.38944 (19)	0.0313 (7)
O3B	0.4667 (4)	-0.0367 (6)	0.4615 (2)	0.0302 (7)
O4B	0.5834 (4)	0.1524 (6)	0.4402 (2)	0.0349 (7)
Cl1B	0.23508 (13)	0.1574 (2)	0.41731 (7)	0.0380 (3)
Cl2B	0.48229 (12)	0.2267 (2)	0.55864 (7)	0.0318 (2)
Cl3B	-0.11866 (13)	0.0187 (2)	0.68821 (8)	0.0378 (3)
Cl4B	0.3715 (2)	0.0537 (3)	0.29351 (9)	0.0563 (4)
Cl5B	0.50048 (19)	0.2929 (2)	0.27257 (8)	0.0487 (4)
Cl6B	0.57633 (19)	0.0222 (3)	0.30941 (8)	0.0489 (4)
C1B	0.2034 (5)	0.1376 (8)	0.5547 (3)	0.0240 (9)
C2B	0.1855 (5)	0.1373 (8)	0.5077 (3)	0.0273 (9)
H2B	0.1222	0.1221	0.4967	0.033*
C3B	0.2590 (5)	0.1590 (8)	0.4759 (3)	0.0276 (9)
C4B	0.3515 (5)	0.1866 (8)	0.4923 (3)	0.0252 (9)
C5B	0.3684 (5)	0.1883 (8)	0.5392 (3)	0.0259 (9)
C6B	0.2963 (5)	0.1672 (8)	0.5714 (3)	0.0273 (9)
H6B	0.3089	0.1725	0.6038	0.033*
C7B	0.4686 (6)	0.2041 (9)	0.3589 (3)	0.0326 (10)
H7B1	0.5276	0.2556	0.3684	0.039*
H7B2	0.4146	0.2736	0.3604	0.039*
C8B	0.4791 (7)	0.1461 (10)	0.3102 (3)	0.0431 (12)
C1'B	0.1230 (5)	0.1095 (8)	0.5886 (3)	0.0284 (9)
C2'B	0.1395 (5)	0.0364 (9)	0.6294 (3)	0.0314 (11)
H2'B	0.2024	0.0042	0.6364	0.038*
C3'B	0.0659 (5)	0.0087 (8)	0.6607 (3)	0.0307 (10)
H3'B	0.0784	-0.0421	0.6888	0.037*
C4'B	-0.0243 (5)	0.0551 (9)	0.6506 (3)	0.0284 (10)
C5'B	-0.0442 (5)	0.1307 (8)	0.6098 (3)	0.0293 (10)
H5'B	-0.1072	0.1629	0.6031	0.035*
C6'B	0.0308 (5)	0.1580 (8)	0.5789 (3)	0.0285 (9)
H6'B	0.0188	0.2102	0.5510	0.034*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1A	0.0183 (5)	0.0276 (5)	0.0392 (6)	0.0017 (4)	0.0002 (4)	0.0039 (4)
O1A	0.0202 (14)	0.0205 (15)	0.0411 (17)	-0.0046 (12)	0.0037 (13)	0.0031 (12)
O2A	0.0285 (16)	0.0267 (16)	0.0386 (17)	-0.0036 (13)	0.0023 (13)	0.0059 (13)
O3A	0.0248 (17)	0.0217 (16)	0.0442 (18)	0.0022 (12)	0.0006 (13)	0.0063 (13)
O4A	0.0187 (15)	0.0370 (18)	0.0491 (19)	-0.0033 (13)	-0.0001 (14)	0.0041 (15)
Cl1A	0.0219 (5)	0.0543 (7)	0.0379 (6)	-0.0016 (5)	-0.0044 (5)	0.0037 (5)

Cl2A	0.0167 (5)	0.0340 (6)	0.0447 (6)	-0.0060 (4)	-0.0047 (4)	0.0021 (5)
Cl3A	0.0279 (6)	0.0455 (7)	0.0399 (6)	-0.0032 (5)	0.0037 (5)	0.0000 (6)
Cl4A	0.0802 (11)	0.0341 (6)	0.0545 (8)	-0.0139 (7)	-0.0252 (7)	0.0035 (5)
Cl5A	0.0709 (11)	0.0286 (7)	0.0465 (9)	0.0093 (7)	0.0109 (8)	0.0067 (7)
Cl6A	0.0715 (11)	0.0299 (8)	0.0454 (9)	0.0031 (7)	0.0054 (8)	0.0030 (7)
C1A	0.018 (2)	0.0134 (18)	0.040 (2)	0.0007 (16)	0.0015 (18)	0.0006 (17)
C2A	0.0137 (19)	0.023 (2)	0.045 (3)	-0.0001 (16)	-0.0014 (17)	0.0013 (19)
C3A	0.019 (2)	0.020 (2)	0.044 (2)	0.0014 (17)	-0.0017 (18)	0.0002 (18)
C4A	0.0122 (18)	0.0156 (19)	0.048 (3)	-0.0009 (15)	-0.0002 (18)	0.0014 (17)
C5A	0.017 (2)	0.0120 (18)	0.048 (3)	0.0000 (16)	0.0010 (18)	0.0004 (17)
C6A	0.020 (2)	0.022 (2)	0.040 (2)	-0.0008 (17)	-0.0039 (18)	-0.0016 (18)
C7A	0.033 (2)	0.027 (2)	0.038 (2)	0.001 (2)	-0.002 (2)	0.0039 (19)
C8A	0.060 (3)	0.028 (2)	0.041 (3)	-0.002 (2)	-0.007 (3)	0.000 (2)
C1'A	0.017 (2)	0.024 (2)	0.044 (2)	0.0023 (17)	-0.0018 (18)	-0.0069 (19)
C2'A	0.017 (2)	0.034 (3)	0.043 (2)	0.0000 (19)	-0.0038 (19)	0.000 (2)
C3'A	0.025 (2)	0.030 (2)	0.037 (2)	0.002 (2)	-0.005 (2)	0.000 (2)
C4'A	0.019 (2)	0.026 (2)	0.040 (2)	-0.0023 (18)	0.0028 (18)	-0.0037 (19)
C5'A	0.019 (2)	0.026 (2)	0.043 (3)	0.0008 (18)	-0.0034 (18)	-0.003 (2)
C6'A	0.019 (2)	0.023 (2)	0.042 (3)	-0.0013 (17)	-0.0001 (19)	0.001 (2)
S1B	0.0183 (5)	0.0276 (5)	0.0392 (6)	0.0017 (4)	0.0002 (4)	0.0039 (4)
O1B	0.0202 (14)	0.0205 (15)	0.0411 (17)	-0.0046 (12)	0.0037 (13)	0.0031 (12)
O2B	0.0285 (16)	0.0267 (16)	0.0386 (17)	-0.0036 (13)	0.0023 (13)	0.0059 (13)
O3B	0.0248 (17)	0.0217 (16)	0.0442 (18)	0.0022 (12)	0.0006 (13)	0.0063 (13)
O4B	0.0187 (15)	0.0370 (18)	0.0491 (19)	-0.0033 (13)	-0.0001 (14)	0.0041 (15)
Cl1B	0.0219 (5)	0.0543 (7)	0.0379 (6)	-0.0016 (5)	-0.0044 (5)	0.0037 (5)
Cl2B	0.0167 (5)	0.0340 (6)	0.0447 (6)	-0.0060 (4)	-0.0047 (4)	0.0021 (5)
Cl3B	0.0279 (6)	0.0455 (7)	0.0399 (6)	-0.0032 (5)	0.0037 (5)	0.0000 (6)
Cl4B	0.0802 (11)	0.0341 (6)	0.0545 (8)	-0.0139 (7)	-0.0252 (7)	0.0035 (5)
Cl5B	0.0709 (11)	0.0286 (7)	0.0465 (9)	0.0093 (7)	0.0109 (8)	0.0067 (7)
Cl6B	0.0715 (11)	0.0299 (8)	0.0454 (9)	0.0031 (7)	0.0054 (8)	0.0030 (7)
C1B	0.018 (2)	0.0134 (18)	0.040 (2)	0.0007 (16)	0.0015 (18)	0.0006 (17)
C2B	0.0137 (19)	0.023 (2)	0.045 (3)	-0.0001 (16)	-0.0014 (17)	0.0013 (19)
C3B	0.019 (2)	0.020 (2)	0.044 (2)	0.0014 (17)	-0.0017 (18)	0.0002 (18)
C4B	0.0122 (18)	0.0156 (19)	0.048 (3)	-0.0009 (15)	-0.0002 (18)	0.0014 (17)
C5B	0.017 (2)	0.0120 (18)	0.048 (3)	0.0000 (16)	0.0010 (18)	0.0004 (17)
C6B	0.020 (2)	0.022 (2)	0.040 (2)	-0.0008 (17)	-0.0039 (18)	-0.0016 (18)
C7B	0.033 (2)	0.027 (2)	0.038 (2)	0.001 (2)	-0.002 (2)	0.0039 (19)
C8B	0.060 (3)	0.028 (2)	0.041 (3)	-0.002 (2)	-0.007 (3)	0.000 (2)
C1'B	0.017 (2)	0.024 (2)	0.044 (2)	0.0023 (17)	-0.0018 (18)	-0.0069 (19)
C2'B	0.017 (2)	0.034 (3)	0.043 (2)	0.0000 (19)	-0.0038 (19)	0.000 (2)
C3'B	0.025 (2)	0.030 (2)	0.037 (2)	0.002 (2)	-0.005 (2)	0.000 (2)
C4'B	0.019 (2)	0.026 (2)	0.040 (2)	-0.0023 (18)	0.0028 (18)	-0.0037 (19)
C5'B	0.019 (2)	0.026 (2)	0.043 (3)	0.0008 (18)	-0.0034 (18)	-0.003 (2)
C6'B	0.019 (2)	0.023 (2)	0.042 (3)	-0.0013 (17)	-0.0001 (19)	0.001 (2)

Geometric parameters ( $\text{\AA}$ ,  $\text{^{\circ}}$ )

S1A—O4A	1.408 (5)	S1B—O3B	1.396 (6)
S1A—O3A	1.414 (6)	S1B—O4B	1.409 (5)
S1A—O2A	1.563 (6)	S1B—O2B	1.563 (6)
S1A—O1A	1.600 (5)	S1B—O1B	1.590 (5)
O1A—C4A	1.393 (9)	O1B—C4B	1.397 (9)
O2A—C7A	1.464 (9)	O2B—C7B	1.434 (9)
C11A—C3A	1.722 (9)	C11B—C3B	1.720 (8)
C12A—C5A	1.749 (7)	C12B—C5B	1.726 (7)
C13A—C4'A	1.730 (8)	C13B—C4'B	1.739 (8)
C14A—C8A	1.783 (10)	C14B—C8B	1.793 (10)
C15A—C8A	1.772 (10)	C15B—C8B	1.755 (9)
C16A—C8A	1.774 (9)	C16B—C8B	1.774 (10)
C1A—C2A	1.391 (11)	C1B—C2B	1.378 (11)
C1A—C6A	1.400 (10)	C1B—C6B	1.412 (10)
C1A—C1'A	1.467 (10)	C1B—C1'B	1.510 (10)
C2A—C3A	1.390 (11)	C2B—C3B	1.389 (11)
C2A—H2A	0.9500	C2B—H2B	0.9500
C3A—C4A	1.390 (10)	C3B—C4B	1.401 (10)
C4A—C5A	1.373 (11)	C4B—C5B	1.370 (11)
C5A—C6A	1.381 (11)	C5B—C6B	1.384 (11)
C6A—H6A	0.9500	C6B—H6B	0.9500
C7A—C8A	1.500 (12)	C7B—C8B	1.509 (12)
C7A—H7A1	0.9900	C7B—H7B1	0.9900
C7A—H7A2	0.9900	C7B—H7B2	0.9900
C1'A—C6'A	1.399 (10)	C1'B—C2'B	1.373 (12)
C1'A—C2'A	1.400 (12)	C1'B—C6'B	1.393 (10)
C2'A—C3'A	1.380 (12)	C2'B—C3'B	1.391 (12)
C2'A—H2'A	0.9500	C2'B—H2'B	0.9500
C3'A—C4'A	1.389 (11)	C3'B—C4'B	1.364 (10)
C3'A—H3'A	0.9500	C3'B—H3'B	0.9500
C4'A—C5'A	1.369 (12)	C4'B—C5'B	1.392 (12)
C5'A—C6'A	1.388 (11)	C5'B—C6'B	1.399 (11)
C5'A—H5'A	0.9500	C5'B—H5'B	0.9500
C6'A—H6'A	0.9500	C6'B—H6'B	0.9500
O4A—S1A—O3A	121.2 (3)	O3B—S1B—O4B	122.7 (3)
O4A—S1A—O2A	109.9 (3)	O3B—S1B—O2B	105.6 (3)
O3A—S1A—O2A	106.0 (3)	O4B—S1B—O2B	110.3 (3)
O4A—S1A—O1A	106.0 (3)	O3B—S1B—O1B	109.4 (3)
O3A—S1A—O1A	110.5 (3)	O4B—S1B—O1B	105.4 (3)
O2A—S1A—O1A	101.4 (3)	O2B—S1B—O1B	101.4 (3)
C4A—O1A—S1A	119.3 (4)	C4B—O1B—S1B	120.0 (4)
C7A—O2A—S1A	117.1 (5)	C7B—O2B—S1B	117.5 (5)
C2A—C1A—C6A	118.1 (7)	C2B—C1B—C6B	120.1 (7)
C2A—C1A—C1'A	121.5 (7)	C2B—C1B—C1'B	119.9 (6)
C6A—C1A—C1'A	120.3 (7)	C6B—C1B—C1'B	120.0 (7)

C3A—C2A—C1A	121.1 (7)	C1B—C2B—C3B	120.8 (7)
C3A—C2A—H2A	119.5	C1B—C2B—H2B	119.6
C1A—C2A—H2A	119.5	C3B—C2B—H2B	119.6
C4A—C3A—C2A	120.0 (8)	C2B—C3B—C4B	119.3 (8)
C4A—C3A—Cl1A	120.1 (6)	C2B—C3B—Cl1B	119.9 (6)
C2A—C3A—Cl1A	119.8 (6)	C4B—C3B—Cl1B	120.7 (6)
C5A—C4A—C3A	119.0 (7)	C5B—C4B—O1B	120.6 (6)
C5A—C4A—O1A	120.1 (6)	C5B—C4B—C3B	119.5 (7)
C3A—C4A—O1A	120.5 (7)	O1B—C4B—C3B	119.8 (7)
C4A—C5A—C6A	121.5 (7)	C4B—C5B—C6B	122.2 (7)
C4A—C5A—Cl2A	118.8 (6)	C4B—C5B—Cl2B	118.7 (6)
C6A—C5A—Cl2A	119.7 (6)	C6B—C5B—Cl2B	119.0 (6)
C5A—C6A—C1A	120.3 (8)	C5B—C6B—C1B	118.1 (7)
C5A—C6A—H6A	119.9	C5B—C6B—H6B	121.0
C1A—C6A—H6A	119.9	C1B—C6B—H6B	121.0
O2A—C7A—C8A	105.4 (6)	O2B—C7B—C8B	108.2 (7)
O2A—C7A—H7A1	110.7	O2B—C7B—H7B1	110.0
C8A—C7A—H7A1	110.7	C8B—C7B—H7B1	110.0
O2A—C7A—H7A2	110.7	O2B—C7B—H7B2	110.0
C8A—C7A—H7A2	110.7	C8B—C7B—H7B2	110.0
H7A1—C7A—H7A2	108.8	H7B1—C7B—H7B2	108.4
C7A—C8A—Cl5A	112.5 (6)	C7B—C8B—Cl5B	108.6 (6)
C7A—C8A—Cl6A	106.7 (6)	C7B—C8B—Cl6B	108.2 (6)
Cl5A—C8A—Cl6A	109.3 (5)	Cl5B—C8B—Cl6B	110.8 (6)
C7A—C8A—Cl4A	110.8 (7)	C7B—C8B—Cl4B	109.5 (7)
Cl5A—C8A—Cl4A	108.6 (5)	Cl5B—C8B—Cl4B	110.0 (5)
Cl6A—C8A—Cl4A	108.8 (5)	Cl6B—C8B—Cl4B	109.7 (5)
C6'A—C1'A—C2'A	118.2 (7)	C2'B—C1'B—C6'B	118.9 (7)
C6'A—C1'A—C1A	120.1 (7)	C2'B—C1'B—C1B	120.7 (6)
C2'A—C1'A—C1A	121.6 (6)	C6'B—C1'B—C1B	120.4 (7)
C3'A—C2'A—C1'A	121.3 (7)	C1'B—C2'B—C3'B	121.2 (7)
C3'A—C2'A—H2'A	119.3	C1'B—C2'B—H2'B	119.4
C1'A—C2'A—H2'A	119.3	C3'B—C2'B—H2'B	119.4
C2'A—C3'A—C4'A	119.3 (8)	C4'B—C3'B—C2'B	119.4 (8)
C2'A—C3'A—H3'A	120.3	C4'B—C3'B—H3'B	120.3
C4'A—C3'A—H3'A	120.3	C2'B—C3'B—H3'B	120.3
C5'A—C4'A—C3'A	120.3 (7)	C3'B—C4'B—C5'B	121.3 (8)
C5'A—C4'A—Cl3A	120.7 (6)	C3'B—C4'B—Cl3B	120.7 (7)
C3'A—C4'A—Cl3A	118.9 (7)	C5'B—C4'B—Cl3B	118.0 (6)
C4'A—C5'A—C6'A	120.7 (7)	C4'B—C5'B—C6'B	118.5 (7)
C4'A—C5'A—H5'A	119.6	C4'B—C5'B—H5'B	120.8
C6'A—C5'A—H5'A	119.6	C6'B—C5'B—H5'B	120.8
C5'A—C6'A—C1'A	120.0 (8)	C1'B—C6'B—C5'B	120.6 (8)
C5'A—C6'A—H6'A	120.0	C1'B—C6'B—H6'B	119.7
C1'A—C6'A—H6'A	120.0	C5'B—C6'B—H6'B	119.7
O4A—S1A—O1A—C4A	138.7 (5)	O3B—S1B—O1B—C4B	-4.9 (6)
O3A—S1A—O1A—C4A	5.6 (6)	O4B—S1B—O1B—C4B	-138.6 (6)

O2A—S1A—O1A—C4A	−106.4 (5)	O2B—S1B—O1B—C4B	106.3 (6)
O4A—S1A—O2A—C7A	45.2 (6)	O3B—S1B—O2B—C7B	−177.9 (5)
O3A—S1A—O2A—C7A	177.8 (5)	O4B—S1B—O2B—C7B	−43.3 (6)
O1A—S1A—O2A—C7A	−66.7 (5)	O1B—S1B—O2B—C7B	68.0 (6)
C6A—C1A—C2A—C3A	−1.2 (11)	C6B—C1B—C2B—C3B	3.1 (11)
C1'A—C1A—C2A—C3A	177.5 (7)	C1'B—C1B—C2B—C3B	−178.2 (7)
C1A—C2A—C3A—C4A	0.9 (11)	C1B—C2B—C3B—C4B	−2.3 (11)
C1A—C2A—C3A—Cl1A	−178.0 (6)	C1B—C2B—C3B—Cl1B	179.7 (6)
C2A—C3A—C4A—C5A	1.0 (11)	S1B—O1B—C4B—C5B	91.6 (8)
Cl1A—C3A—C4A—C5A	179.9 (5)	S1B—O1B—C4B—C3B	−92.2 (7)
C2A—C3A—C4A—O1A	174.4 (6)	C2B—C3B—C4B—C5B	1.6 (11)
Cl1A—C3A—C4A—O1A	−6.7 (10)	Cl1B—C3B—C4B—C5B	179.5 (6)
S1A—O1A—C4A—C5A	−92.0 (7)	C2B—C3B—C4B—O1B	−174.6 (6)
S1A—O1A—C4A—C3A	94.7 (7)	Cl1B—C3B—C4B—O1B	3.3 (10)
C3A—C4A—C5A—C6A	−2.6 (11)	O1B—C4B—C5B—C6B	174.4 (6)
O1A—C4A—C5A—C6A	−176.0 (6)	C3B—C4B—C5B—C6B	−1.8 (11)
C3A—C4A—C5A—Cl2A	177.1 (5)	O1B—C4B—C5B—Cl2B	−2.3 (10)
O1A—C4A—C5A—Cl2A	3.7 (9)	C3B—C4B—C5B—Cl2B	−178.5 (5)
C4A—C5A—C6A—C1A	2.3 (11)	C4B—C5B—C6B—C1B	2.5 (11)
Cl2A—C5A—C6A—C1A	−177.4 (5)	Cl2B—C5B—C6B—C1B	179.2 (5)
C2A—C1A—C6A—C5A	−0.4 (11)	C2B—C1B—C6B—C5B	−3.2 (11)
C1'A—C1A—C6A—C5A	−179.1 (6)	C1'B—C1B—C6B—C5B	178.2 (6)
S1A—O2A—C7A—C8A	−148.3 (6)	S1B—O2B—C7B—C8B	148.2 (6)
O2A—C7A—C8A—Cl5A	58.7 (8)	O2B—C7B—C8B—Cl5B	−180.0 (5)
O2A—C7A—C8A—Cl6A	178.5 (5)	O2B—C7B—C8B—Cl6B	−59.6 (8)
O2A—C7A—C8A—Cl4A	−63.2 (7)	O2B—C7B—C8B—Cl4B	59.9 (8)
C2A—C1A—C1'A—C6'A	36.2 (11)	C2B—C1B—C1'B—C2'B	145.8 (8)
C6A—C1A—C1'A—C6'A	−145.1 (7)	C6B—C1B—C1'B—C2'B	−35.5 (11)
C2A—C1A—C1'A—C2'A	−140.7 (8)	C2B—C1B—C1'B—C6'B	−34.4 (11)
C6A—C1A—C1'A—C2'A	38.0 (11)	C6B—C1B—C1'B—C6'B	144.2 (7)
C6'A—C1'A—C2'A—C3'A	0.1 (12)	C6'B—C1'B—C2'B—C3'B	0.8 (12)
C1A—C1'A—C2'A—C3'A	177.0 (7)	C1B—C1'B—C2'B—C3'B	−179.4 (7)
C1'A—C2'A—C3'A—C4'A	−0.5 (12)	C1'B—C2'B—C3'B—C4'B	0.0 (13)
C2'A—C3'A—C4'A—C5'A	1.7 (12)	C2'B—C3'B—C4'B—C5'B	−0.5 (12)
C2'A—C3'A—C4'A—Cl3A	−177.9 (6)	C2'B—C3'B—C4'B—Cl3B	178.5 (6)
C3'A—C4'A—C5'A—C6'A	−2.4 (12)	C3'B—C4'B—C5'B—C6'B	0.3 (12)
Cl3A—C4'A—C5'A—C6'A	177.1 (6)	Cl3B—C4'B—C5'B—C6'B	−178.8 (6)
C4'A—C5'A—C6'A—C1'A	2.0 (12)	C2'B—C1'B—C6'B—C5'B	−1.1 (12)
C2'A—C1'A—C6'A—C5'A	−0.8 (11)	C1B—C1'B—C6'B—C5'B	179.2 (7)
C1A—C1'A—C6'A—C5'A	−177.8 (7)	C4'B—C5'B—C6'B—C1'B	0.5 (11)