

Tetrakis(μ -3-chlorobenzoato- κ^2 O:O')-bis[(*N,N*-diethylnicotinamide- κ N¹)-copper(II)]

Nihat Bozkurt,^a Tuncay Tuñç,^b Nagihan Çaylak Delibaş,^c Hacı Necefoğlu^a and Tuncer Hökelek^{d*}

^aDepartment of Chemistry, Kafkas University, 36100 Kars, Turkey, ^bAksaray University, Science Education Department, 68100, Aksaray, Turkey, ^cDepartment of Physics, Sakarya University, 54187 Esentepe, Sakarya, Turkey, and ^dDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey
Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 27 June 2013; accepted 28 June 2013

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.031; wR factor = 0.083; data-to-parameter ratio = 16.7.

In the title centrosymmetric binuclear Cu^{II} complex, $[\text{Cu}_2(\text{C}_7\text{H}_4\text{ClO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$, the two Cu^{II} cations [$\text{Cu}\cdots\text{Cu} = 2.6314(4)$ Å] are bridged by four 3-chlorobenzoate (CB) anions. The four carboxylate O atoms around each Cu^{II} cation form a distorted square-planar arrangement, the distorted square-pyramidal coordination geometry being completed by the pyridine N atom of the *N,N*-diethylnicotinamide (DNA) molecule. The dihedral angle between the benzene ring and the carboxylate group is $4.49(11)^\circ$ in one of the independent CB ligands and $12.00(10)^\circ$ in the other. The benzene rings of the independent CB ligands are oriented at a dihedral angle of $84.13(6)^\circ$. In the crystal, weak $\text{C}\cdots\text{H}\cdots\text{O}$ hydrogen bonds link the binuclear complex molecules into supramolecular chains running along [101].

Related literature

For niacin, see: Krishnamachari (1974). For *N,N*-diethylnicotinamide, see: Bigoli *et al.* (1972). For related structures, see: Speier & Fulop (1989); Usabaliev *et al.* (1980); Hökelek *et al.* (1995); Hökelek *et al.* (2009*a,b,c*, 2011); Necefoğlu *et al.* (2010*a,b*); Aydın *et al.* (2012).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_7\text{H}_4\text{ClO}_2)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2]$
 $M_r = 1105.75$
Monoclinic, $P2_1/c$
 $a = 12.6077(4)$ Å
 $b = 16.7569(5)$ Å
 $c = 12.1402(4)$ Å
 $\beta = 99.647(2)^\circ$

$V = 2528.54(14)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.11$ mm⁻¹
 $T = 296$ K
 $0.35 \times 0.25 \times 0.20$ mm

Data collection

Bruker SMART BREEZE CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2012)
 $T_{\text{min}} = 0.711$, $T_{\text{max}} = 0.724$

78293 measured reflections
5166 independent reflections
4702 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.083$
 $S = 1.04$
5166 reflections
309 parameters

85 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.70$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—O1	1.9791 (13)	Cu1—O4 ⁱ	1.9703 (13)
Cu1—O2 ⁱ	1.9688 (13)	Cu1—N1	2.1454 (14)
Cu1—O3	1.9755 (13)		

Symmetry code: (i) $-x + 1, -y + 2, -z + 1$.

Table 2

Hydrogen-bond geometry (Å, °).

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
C17—H17 ⁱⁱ ⋯O5 ⁱⁱ	0.93	2.45	3.221 (3)	140

Symmetry code: (ii) $-x + 2, -y + 2, -z + 2$.

Data collection: APEX2 (Bruker, 2012); cell refinement: SAINT (Bruker, 2012); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: WinGX (Farrugia, 2012) and PLATON (Spek, 2009).

The authors acknowledge the Aksaray University, Science and Technology Application and Research Center, Aksaray, Turkey, for the use of the Bruker SMART BREEZE CCD diffractometer (purchased under grant No. 2010K120480 of the State Planning Organization).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5716).

References

- Aydın, Ö., Çaylak Delibaş, N., Necefoğlu, H. & Hökelek, T. (2012). *Acta Cryst.* **E68**, m409–m410.
- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst.* **B28**, 962–966.
- Bruker (2012). *APEX2, SAINT and SADABS*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Hökelek, T., Dal, H., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009c). *Acta Cryst.* **E65**, m1582–m1583.
- Hökelek, T., Necefoğlu, H. & Balcı, M. (1995). *Acta Cryst.* **C51**, 2020–2023.
- Hökelek, T., Sağlam, E. G., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2011). *Acta Cryst.* **E67**, m28–m29.
- Hökelek, T., Yılmaz, F., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009a). *Acta Cryst.* **E65**, m955–m956.
- Hökelek, T., Yılmaz, F., Tercan, B., Aybirdi, Ö. & Necefoğlu, H. (2009b). *Acta Cryst.* **E65**, m1328–m1329.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Necefoğlu, H., Çimen, E., Tercan, B., Dal, H. & Hökelek, T. (2010a). *Acta Cryst.* **E66**, m334–m335.
- Necefoğlu, H., Çimen, E., Tercan, B., Dal, H. & Hökelek, T. (2010b). *Acta Cryst.* **E66**, m485–m486.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Speier, G. & Fulop, V. (1989). *J. Chem. Soc. Dalton Trans.* pp. 2331–2333.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Usubaliev, B. T., Movsumov, E. M., Musaev, F. N., Nadzhafov, G. N., Amiraslanov, I. R. & Mamedov, Kh. S. (1980). *Koord. Khim.* **6**, 1091–1096.

supporting information

Acta Cryst. (2013). E69, m431–m432 [doi:10.1107/S1600536813017881]

Tetrakis(μ -3-chlorobenzoato- κ^2 O:O')bis[(*N,N*-diethylnicotinamide- κ N¹)copper(II)]

Nihat Bozkurt, Tuncay Tunç, Nagihan Çaylak Delibaş, Hacali Necefoğlu and Tuncer Hökelek

S1. Comment

As a part of our ongoing investigation on transition metal complexes of nicotinamide (NA), one form of niacin (Krishnamachari, 1974), and/or the nicotinic acid derivative *N,N*-diethylnicotinamide (DENA), an important respiratory stimulant (Bigoli *et al.*, 1972), the title compound was synthesized and its crystal structure is reported herein.

The title compound is a binuclear compound, consisting of four chlorobenzoate (CB) ligands. The structures of similar complexes of the Cu²⁺, Zn²⁺ and Co²⁺ ions, [Cu(C₆H₅COO)₂(C₅H₅N)]₂ (Usubaliev *et al.*, 1980); [Cu(C₆H₅CO₂)₂(Py)]₂ (Speier & Fulop, 1989); [Cu₂(C₆H₅COO)₄(C₁₀H₁₄N₂O)₂] (Hökelek *et al.*, 1995); [Cu₂(C₈H₇O₂)₄(C₆H₆N₂O)₂] (Necefoğlu *et al.*, 2010a); [Cu₂(C₇H₄IO₂)₄(H₂O)₂] (Aydın *et al.*, 2012); [Zn₂(C₁₁H₁₄NO₂)₄(C₁₀H₁₄N₂O)₂] (Hökelek *et al.*, 2009a); [Zn₂(C₈H₈NO₂)₄(C₁₀H₁₄N₂O)₂].2H₂O (Hökelek *et al.*, 2009b); [Zn₂(C₉H₁₀NO₂)₄(C₁₀H₁₄N₂O)₂] (Hökelek *et al.*, 2009c); [Zn₂(C₈H₇O₂)₄(C₁₀H₁₄N₂O)₂] (Necefoğlu *et al.*, 2010b) and [Co₂(C₁₁H₁₄NO₂)₄(C₁₀H₁₄N₂O)₂] (Hökelek *et al.*, 2011) have also been determined. In these structures, the benzoate ion acts as a bidentate ligand.

The title dimeric complex, [Cu₂(CB)₄(DENA)₂], has a centre of symmetry and two Cu^{II} atoms are surrounded by four CB groups and two DENA molecules. The CB groups act as bridging ligands. The Cu1...Cu1a [symmetry code: (a) 1 - x, - y, 1 - z] distance is 2.6314 (4) Å. The average Cu-O distance is 1.9734 (13) Å (Table 1), and four O atoms of the bridging CB ligands around each Cu atom form a distorted square plane. The Cu atom lies 1.3997 (2) Å below the least-squares plane. The average O-Cu-O bond angle is 89.41 (6)°. A distorted square-pyramidal arrangement around each Cu atom is completed by the DENA N atom at 2.1454 (14) Å from the Cu atom (Table 1). The N1-Cu1...Cu1a angle is 173.85 (4)° and the dihedral angle between plane through Cu1, O1, O2, C1, Cu1a, O1a, O2a, C1a and the plane through Cu1, O3, O4, C8, Cu1a, O3a, O4a, C8a is 89.76(1.45)°. The dihedral angles between the planar carboxylate groups [(O1/O2/C1) and (O3/O4/C8)] and the adjacent benzene rings A (C2—C7) and B (C9—C14) are 4.49 (11) and 12.00 (10)°, respectively, while that between rings A, B and C (N1/C15—C19) are A/B = 84.13 (6), A/C = 57.94 (6) and B/C = 37.95 (7)°.

In the crystal structure, weak intermolecular C—H...O interactions (Table 2) link the molecules into infinite chains through the centrosymmetric dimers running along the [1 0 1] direction.

S2. Experimental

The title compound was prepared by the reaction of CuSO₄·5H₂O (1.25 g, 5 mmol) in H₂O (100 ml) and diethylnicotinamide (1.78 g, 10 mmol) in H₂O (20 ml) with sodium 3-chlorobenzoate (1.79 g, 10 mmol) in H₂O (100 ml). The mixture was set aside to crystallize at ambient temperature for five days, giving green single crystals.

S3. Refinement

The C-bound H-atoms were positioned geometrically with C—H = 0.93, 0.97 and 0.96 Å, for aromatic, methylene and methyl H-atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.5$ for methyl H-atoms and $k = 1.2$ for all other H-atoms.




Figure 1

The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level [symmetry code: (a) $1 - x, -y, 1 - z$]. Hydrogen atoms have been omitted for clarity.




Figure 2

A view of the packing of the title compound. Hydrogen bonds are shown as dashed lines. Hydrogen atoms not involved in hydrogen bonding have been omitted for clarity.

Tetrakis(μ -3-chlorobenzoato- κ^2 O:O')bis[(*N,N*-diethylnicotinamide- κ N')copper(II)]*Crystal data*[Cu₂(C₇H₄ClO₂)₄(C₁₀H₁₄N₂O)₂] $M_r = 1105.75$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 12.6077$ (4) Å $b = 16.7569$ (5) Å $c = 12.1402$ (4) Å $\beta = 99.647$ (2)° $V = 2528.54$ (14) Å³ $Z = 2$ $F(000) = 1132$ $D_x = 1.452$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9380 reflections

 $\theta = 2.5$ – 29.9 ° $\mu = 1.11$ mm⁻¹ $T = 296$ K

Block, green

 $0.35 \times 0.25 \times 0.20$ mm*Data collection*

Bruker SMART BREEZE CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2012)

 $T_{\min} = 0.711$, $T_{\max} = 0.724$

78293 measured reflections

5166 independent reflections

4702 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.026$ $\theta_{\text{max}} = 26.4$ °, $\theta_{\text{min}} = 1.6$ ° $h = -15 \rightarrow 15$ $k = -20 \rightarrow 20$ $l = -15 \rightarrow 15$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.031$ $wR(F^2) = 0.083$ $S = 1.04$

5166 reflections

309 parameters

85 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0449P)^2 + 1.301P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.002$ $\Delta\rho_{\text{max}} = 0.70$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³*Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.580773 (15)	0.980853 (12)	0.578926 (15)	0.02838 (8)
Cl1	0.53514 (7)	0.79998 (5)	-0.03442 (5)	0.0761 (2)
Cl2	0.92843 (7)	1.31020 (5)	0.55213 (9)	0.1053 (3)
O1	0.63814 (10)	0.91758 (8)	0.46442 (10)	0.0409 (3)

O2	0.49984 (11)	0.95064 (9)	0.33241 (11)	0.0458 (3)
O3	0.64805 (10)	1.07718 (8)	0.52693 (11)	0.0422 (3)
O4	0.51122 (10)	1.10927 (8)	0.39242 (11)	0.0405 (3)
O5	1.04739 (13)	1.05214 (11)	0.83585 (15)	0.0658 (5)
N1	0.70894 (11)	0.96124 (10)	0.71646 (12)	0.0347 (3)
N2	1.06037 (13)	0.95243 (12)	0.71450 (16)	0.0502 (4)
C1	0.58602 (14)	0.91427 (10)	0.36697 (15)	0.0343 (4)
C2	0.62814 (15)	0.86185 (11)	0.28384 (15)	0.0373 (4)
C3	0.57253 (17)	0.85814 (12)	0.17511 (16)	0.0433 (4)
H3	0.5123	0.8900	0.1529	0.052*
C4	0.6080 (2)	0.80633 (14)	0.10029 (17)	0.0518 (5)
C5	0.6965 (2)	0.75909 (16)	0.1303 (2)	0.0661 (7)
H5	0.7186	0.7242	0.0790	0.079*
C6	0.7521 (2)	0.76385 (16)	0.2371 (2)	0.0678 (7)
H6	0.8127	0.7322	0.2581	0.081*
C7	0.71904 (18)	0.81551 (13)	0.31447 (19)	0.0493 (5)
H7	0.7579	0.8188	0.3865	0.059*
C8	0.60290 (14)	1.12046 (10)	0.44801 (14)	0.0319 (3)
C9	0.66516 (15)	1.19115 (11)	0.41815 (16)	0.0368 (4)
C10	0.75758 (17)	1.21469 (12)	0.48927 (19)	0.0463 (5)
H10	0.7814	1.1867	0.5549	0.056*
C11	0.81362 (19)	1.28015 (14)	0.4613 (2)	0.0620 (6)
C12	0.7809 (2)	1.32217 (15)	0.3639 (3)	0.0749 (8)
H12	0.8202	1.3659	0.3461	0.090*
C13	0.6896 (2)	1.29872 (15)	0.2938 (3)	0.0700 (7)
H13	0.6668	1.3267	0.2280	0.084*
C14	0.63081 (19)	1.23348 (13)	0.32036 (19)	0.0506 (5)
H14	0.5685	1.2182	0.2727	0.061*
C15	0.81050 (14)	0.97600 (11)	0.70628 (15)	0.0361 (4)
H15	0.8250	0.9911	0.6366	0.043*
C16	0.89570 (14)	0.97002 (11)	0.79394 (15)	0.0353 (4)
C17	0.87265 (15)	0.94853 (12)	0.89783 (15)	0.0412 (4)
H17	0.9273	0.9452	0.9595	0.049*
C18	0.76805 (17)	0.93223 (14)	0.90842 (16)	0.0468 (5)
H18	0.7512	0.9167	0.9770	0.056*
C19	0.68845 (15)	0.93925 (13)	0.81604 (16)	0.0425 (4)
H19	0.6178	0.9282	0.8237	0.051*
C20	1.00821 (15)	0.99464 (13)	0.78250 (17)	0.0429 (4)
C21	1.02241 (19)	0.87796 (16)	0.6595 (2)	0.0649 (7)
H211	0.9638	0.8574	0.6937	0.078*
H212	1.0803	0.8391	0.6717	0.078*
C22	0.9840 (3)	0.8867 (3)	0.5347 (3)	0.1069 (13)
H221	0.9232	0.9220	0.5219	0.160*
H222	0.9635	0.8354	0.5030	0.160*
H223	1.0410	0.9083	0.5003	0.160*
C23	1.17124 (19)	0.97631 (17)	0.7067 (2)	0.0660 (7)
H231	1.1878	0.9589	0.6353	0.079*
H232	1.1761	1.0341	0.7092	0.079*

C24	1.25301 (19)	0.9423 (2)	0.7983 (3)	0.0759 (8)
H241	1.2468	0.8852	0.7983	0.114*
H242	1.3238	0.9570	0.7864	0.114*
H243	1.2409	0.9630	0.8689	0.114*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02465 (11)	0.03345 (13)	0.02565 (12)	-0.00047 (7)	0.00020 (8)	-0.00112 (8)
C11	0.1050 (5)	0.0886 (5)	0.0378 (3)	-0.0216 (4)	0.0206 (3)	-0.0159 (3)
C12	0.0775 (5)	0.0876 (5)	0.1497 (8)	-0.0475 (4)	0.0157 (5)	-0.0400 (6)
O1	0.0411 (7)	0.0475 (8)	0.0346 (7)	0.0051 (6)	0.0077 (5)	-0.0066 (6)
O2	0.0428 (7)	0.0565 (8)	0.0370 (7)	0.0102 (6)	0.0033 (6)	-0.0131 (6)
O3	0.0388 (7)	0.0412 (7)	0.0432 (7)	-0.0097 (6)	-0.0030 (5)	0.0079 (6)
O4	0.0352 (7)	0.0427 (7)	0.0420 (7)	-0.0059 (5)	0.0018 (5)	0.0078 (6)
O5	0.0505 (9)	0.0705 (11)	0.0739 (11)	-0.0157 (8)	0.0032 (8)	-0.0230 (9)
N1	0.0292 (7)	0.0439 (8)	0.0293 (7)	0.0024 (6)	0.0000 (6)	0.0020 (6)
N2	0.0353 (8)	0.0583 (11)	0.0583 (11)	-0.0018 (8)	0.0114 (7)	-0.0050 (9)
C1	0.0339 (9)	0.0340 (9)	0.0370 (9)	-0.0054 (7)	0.0115 (7)	-0.0037 (7)
C2	0.0426 (10)	0.0342 (9)	0.0386 (9)	-0.0065 (7)	0.0175 (8)	-0.0034 (7)
C3	0.0485 (11)	0.0447 (11)	0.0396 (10)	-0.0073 (9)	0.0152 (8)	-0.0041 (8)
C4	0.0713 (14)	0.0516 (12)	0.0372 (10)	-0.0171 (11)	0.0227 (10)	-0.0097 (9)
C5	0.0912 (19)	0.0568 (14)	0.0586 (14)	0.0074 (13)	0.0371 (13)	-0.0140 (12)
C6	0.0772 (17)	0.0645 (16)	0.0668 (15)	0.0230 (13)	0.0275 (13)	-0.0051 (13)
C7	0.0532 (12)	0.0502 (12)	0.0475 (11)	0.0052 (9)	0.0175 (9)	-0.0023 (9)
C8	0.0336 (8)	0.0320 (8)	0.0313 (8)	-0.0012 (7)	0.0089 (7)	-0.0025 (7)
C9	0.0388 (9)	0.0325 (9)	0.0422 (10)	-0.0010 (7)	0.0160 (7)	-0.0024 (7)
C10	0.0464 (11)	0.0413 (10)	0.0532 (12)	-0.0095 (9)	0.0141 (9)	-0.0081 (9)
C11	0.0547 (13)	0.0458 (12)	0.0905 (18)	-0.0178 (10)	0.0269 (12)	-0.0182 (12)
C12	0.0827 (19)	0.0418 (12)	0.112 (2)	-0.0138 (12)	0.0507 (17)	0.0072 (14)
C13	0.0880 (19)	0.0502 (14)	0.0798 (18)	0.0066 (13)	0.0367 (15)	0.0238 (13)
C14	0.0570 (13)	0.0455 (11)	0.0523 (12)	0.0032 (9)	0.0178 (10)	0.0091 (9)
C15	0.0328 (9)	0.0463 (10)	0.0284 (8)	0.0019 (7)	0.0030 (7)	0.0019 (7)
C16	0.0298 (8)	0.0374 (9)	0.0368 (9)	0.0048 (7)	-0.0002 (7)	-0.0041 (7)
C17	0.0390 (10)	0.0477 (11)	0.0323 (9)	0.0063 (8)	-0.0071 (7)	0.0005 (8)
C18	0.0478 (11)	0.0610 (13)	0.0305 (9)	0.0047 (9)	0.0032 (8)	0.0074 (9)
C19	0.0341 (9)	0.0564 (12)	0.0366 (9)	-0.0003 (8)	0.0053 (7)	0.0065 (9)
C20	0.0325 (9)	0.0496 (11)	0.0430 (10)	0.0018 (8)	-0.0034 (8)	-0.0001 (9)
C21	0.0471 (12)	0.0638 (15)	0.0857 (17)	0.0030 (11)	0.0162 (12)	-0.0228 (13)
C22	0.087 (2)	0.141 (3)	0.086 (2)	0.022 (2)	-0.0043 (18)	-0.052 (2)
C23	0.0429 (12)	0.0784 (18)	0.0805 (18)	-0.0069 (11)	0.0214 (12)	0.0006 (14)
C24	0.0368 (12)	0.092 (2)	0.100 (2)	0.0002 (12)	0.0133 (12)	-0.0083 (17)

Geometric parameters (Å, °)

Cu1—Cu1 ⁱ	2.6314 (4)	C9—C10	1.386 (3)
Cu1—O1	1.9791 (13)	C9—C14	1.388 (3)
Cu1—O2 ⁱ	1.9688 (13)	C10—C11	1.378 (3)

Cu1—O3	1.9755 (13)	C10—H10	0.9300
Cu1—O4 ⁱ	1.9703 (13)	C11—C12	1.378 (4)
Cu1—N1	2.1454 (14)	C12—H12	0.9300
C11—C4	1.740 (2)	C13—C12	1.369 (4)
C12—C11	1.741 (3)	C13—H13	0.9300
O1—C1	1.255 (2)	C14—C13	1.388 (3)
O2—C1	1.255 (2)	C14—H14	0.9300
O2—Cu1 ⁱ	1.9688 (13)	C15—H15	0.9300
O3—C8	1.259 (2)	C16—C15	1.383 (2)
O4—Cu1 ⁱ	1.9703 (13)	C16—C20	1.506 (3)
O4—C8	1.251 (2)	C17—C16	1.388 (3)
O5—C20	1.218 (3)	C17—C18	1.374 (3)
N1—C15	1.330 (2)	C17—H17	0.9300
N1—C19	1.330 (2)	C18—H18	0.9300
N2—C20	1.340 (3)	C19—C18	1.379 (3)
N2—C21	1.458 (3)	C19—H19	0.9300
N2—C23	1.472 (3)	C21—C22	1.519 (4)
C1—C2	1.500 (2)	C21—H211	0.9700
C2—C3	1.388 (3)	C21—H212	0.9700
C2—C7	1.383 (3)	C22—H221	0.9600
C3—C4	1.384 (3)	C22—H222	0.9600
C3—H3	0.9300	C22—H223	0.9600
C4—C5	1.367 (4)	C23—C24	1.496 (4)
C5—C6	1.369 (4)	C23—H232	0.9700
C5—H5	0.9300	C23—H231	0.9700
C6—H6	0.9300	C24—H241	0.9600
C7—C6	1.392 (3)	C24—H242	0.9600
C7—H7	0.9300	C24—H243	0.9600
C8—C9	1.499 (2)		
O1—Cu1—Cu1 ⁱ	87.05 (4)	C10—C11—C12	118.7 (2)
O1—Cu1—N1	98.53 (6)	C10—C11—C12	121.7 (2)
O2 ⁱ —Cu1—Cu1 ⁱ	81.29 (4)	C12—C11—C12	119.57 (19)
O2 ⁱ —Cu1—O1	168.33 (6)	C11—C12—H12	120.5
O2 ⁱ —Cu1—O3	89.25 (6)	C13—C12—C11	119.1 (2)
O2 ⁱ —Cu1—O4 ⁱ	88.70 (6)	C13—C12—H12	120.5
O2 ⁱ —Cu1—N1	93.11 (6)	C12—C13—C14	120.5 (3)
O3—Cu1—Cu1 ⁱ	83.65 (4)	C12—C13—H13	119.8
O3—Cu1—O1	89.21 (6)	C14—C13—H13	119.8
O3—Cu1—N1	93.79 (6)	C9—C14—C13	119.9 (2)
O4 ⁱ —Cu1—Cu1 ⁱ	84.77 (4)	C9—C14—H14	120.0
O4 ⁱ —Cu1—O1	90.49 (6)	C13—C14—H14	120.0
O4 ⁱ —Cu1—O3	168.41 (5)	N1—C15—C16	123.41 (17)
O4 ⁱ —Cu1—N1	97.71 (6)	N1—C15—H15	118.3
N1—Cu1—Cu1 ⁱ	173.85 (4)	C16—C15—H15	118.3
C1—O1—Cu1	119.39 (12)	C15—C16—C17	117.64 (17)
C1—O2—Cu1 ⁱ	126.70 (12)	C15—C16—C20	122.09 (17)
C8—O3—Cu1	123.38 (11)	C17—C16—C20	119.91 (16)

C8—O4—Cu1 ⁱ	122.52 (12)	C16—C17—H17	120.4
C15—N1—Cu1	120.69 (12)	C18—C17—C16	119.14 (16)
C19—N1—Cu1	120.94 (12)	C18—C17—H17	120.4
C19—N1—C15	118.21 (15)	C17—C18—C19	119.09 (18)
C20—N2—C21	125.21 (18)	C17—C18—H18	120.5
C20—N2—C23	117.7 (2)	C19—C18—H18	120.5
C21—N2—C23	116.48 (19)	N1—C19—C18	122.49 (18)
O1—C1—O2	125.52 (16)	N1—C19—H19	118.8
O1—C1—C2	118.14 (16)	C18—C19—H19	118.8
O2—C1—C2	116.33 (16)	O5—C20—N2	123.2 (2)
C3—C2—C1	119.11 (17)	O5—C20—C16	118.15 (19)
C7—C2—C1	121.25 (17)	N2—C20—C16	118.64 (18)
C7—C2—C3	119.61 (18)	N2—C21—C22	113.6 (3)
C2—C3—H3	120.5	N2—C21—H211	108.9
C4—C3—C2	119.0 (2)	N2—C21—H212	108.9
C4—C3—H3	120.5	C22—C21—H211	108.9
C3—C4—C11	118.5 (2)	C22—C21—H212	108.9
C5—C4—C11	119.61 (17)	H211—C21—H212	107.7
C5—C4—C3	121.8 (2)	C21—C22—H221	109.5
C4—C5—C6	119.0 (2)	C21—C22—H222	109.5
C4—C5—H5	120.5	C21—C22—H223	109.5
C6—C5—H5	120.5	H221—C22—H222	109.5
C5—C6—C7	120.8 (2)	H221—C22—H223	109.5
C5—C6—H6	119.6	H222—C22—H223	109.5
C7—C6—H6	119.6	N2—C23—C24	113.1 (2)
C2—C7—C6	119.8 (2)	N2—C23—H232	109.0
C2—C7—H7	120.1	N2—C23—H231	109.0
C6—C7—H7	120.1	C24—C23—H232	109.0
O3—C8—C9	116.81 (15)	C24—C23—H231	109.0
O4—C8—O3	125.68 (16)	H232—C23—H231	107.8
O4—C8—C9	117.52 (16)	C23—C24—H243	109.5
C10—C9—C8	119.60 (18)	C23—C24—H242	109.5
C10—C9—C14	119.75 (19)	C23—C24—H241	109.5
C14—C9—C8	120.64 (18)	H243—C24—H242	109.5
C9—C10—H10	120.5	H243—C24—H241	109.5
C11—C10—C9	119.0 (2)	H242—C24—H241	109.5
C11—C10—H10	120.5		
Cu1 ⁱ —Cu1—O1—C1	-1.49 (13)	O1—C1—C2—C3	-179.12 (17)
O2 ⁱ —Cu1—O1—C1	-2.7 (4)	O1—C1—C2—C7	2.9 (3)
O3—Cu1—O1—C1	-85.17 (14)	O2—C1—C2—C3	1.7 (3)
O4 ⁱ —Cu1—O1—C1	83.25 (14)	O2—C1—C2—C7	-176.27 (18)
N1—Cu1—O1—C1	-178.88 (13)	C1—C2—C3—C4	-176.33 (17)
Cu1 ⁱ —Cu1—O3—C8	-0.31 (14)	C7—C2—C3—C4	1.7 (3)
O1—Cu1—O3—C8	86.81 (15)	C3—C2—C7—C6	-1.9 (3)
O2 ⁱ —Cu1—O3—C8	-81.63 (15)	C1—C2—C7—C6	176.0 (2)
O4 ⁱ —Cu1—O3—C8	-1.8 (4)	C2—C3—C4—C11	178.00 (15)
N1—Cu1—O3—C8	-174.70 (14)	C2—C3—C4—C5	-0.4 (3)

O1—Cu1—N1—C15	52.96 (15)	C11—C4—C5—C6	-179.0 (2)
O1—Cu1—N1—C19	-131.61 (16)	C3—C4—C5—C6	-0.7 (4)
O2 ⁱ —Cu1—N1—C15	-126.27 (15)	C4—C5—C6—C7	0.4 (4)
O2 ⁱ —Cu1—N1—C19	49.16 (16)	C2—C7—C6—C5	0.9 (4)
O3—Cu1—N1—C15	-36.82 (15)	O3—C8—C9—C10	-12.2 (2)
O3—Cu1—N1—C19	138.61 (16)	O3—C8—C9—C14	168.16 (18)
O4 ⁱ —Cu1—N1—C15	144.62 (14)	O4—C8—C9—C10	168.11 (17)
O4 ⁱ —Cu1—N1—C19	-39.95 (16)	O4—C8—C9—C14	-11.5 (3)
Cu1—O1—C1—O2	2.8 (3)	C8—C9—C10—C11	-179.51 (18)
Cu1—O1—C1—C2	-176.35 (11)	C14—C9—C10—C11	0.1 (3)
Cu1 ⁱ —O2—C1—O1	-2.6 (3)	C8—C9—C14—C13	-179.86 (19)
Cu1 ⁱ —O2—C1—C2	176.52 (12)	C10—C9—C14—C13	0.5 (3)
Cu1—O3—C8—O4	0.1 (3)	C9—C10—C11—C12	179.02 (16)
Cu1—O3—C8—C9	-179.54 (11)	C9—C10—C11—C12	-0.7 (3)
Cu1 ⁱ —O4—C8—O3	0.3 (3)	C10—C11—C12—C13	0.7 (4)
Cu1 ⁱ —O4—C8—C9	179.96 (11)	C12—C11—C12—C13	-179.1 (2)
Cu1—N1—C15—C16	175.23 (14)	C14—C13—C12—C11	0.0 (4)
C19—N1—C15—C16	-0.3 (3)	C9—C14—C13—C12	-0.6 (4)
Cu1—N1—C19—C18	-174.76 (16)	C17—C16—C15—N1	-0.9 (3)
C15—N1—C19—C18	0.8 (3)	C20—C16—C15—N1	-174.04 (18)
C21—N2—C20—O5	172.2 (2)	C15—C16—C20—O5	113.2 (2)
C21—N2—C20—C16	-7.7 (3)	C15—C16—C20—N2	-66.8 (3)
C23—N2—C20—O5	1.4 (3)	C17—C16—C20—O5	-59.7 (3)
C23—N2—C20—C16	-178.50 (19)	C17—C16—C20—N2	120.2 (2)
C20—N2—C21—C22	107.6 (3)	C18—C17—C16—C15	1.7 (3)
C23—N2—C21—C22	-81.5 (3)	C18—C17—C16—C20	174.99 (19)
C20—N2—C23—C24	83.6 (3)	C16—C17—C18—C19	-1.3 (3)
C21—N2—C23—C24	-88.0 (3)	N1—C19—C18—C17	0.0 (3)

Symmetry code: (i) $-x+1, -y+2, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C17—H17 \cdots O5 ⁱⁱ	0.93	2.45	3.221 (3)	140

Symmetry code: (ii) $-x+2, -y+2, -z+2$.