

Dichlorido{2-[*(E*)-phenyl(pyridin-2-yl- κ N)methylidene]-*N*-phenylhydrazinecarboxamide- κ^2 *N*²,O}copper(II)

N. Aiswarya,^a M. Sithambaresan,^{b*}
M. R. Prathapachandra Kurup^a and Seik Weng Ng^{c,d}

^aDepartment of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, India, ^bDepartment of Chemistry, Faculty of Science, Eastern University, Sri Lanka, Chenkalady, Sri Lanka, ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^dChemistry Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia
Correspondence e-mail: eesans@yahoo.com


Received 27 September 2013; accepted 30 September 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.004$ Å;
 R factor = 0.034; wR factor = 0.092; data-to-parameter ratio = 17.5.

The title compound, $[CuCl_2(C_{19}H_{16}N_4O)]$, contains a Cu^{II} atom *N,N'*,*O*-chelated by a neutral *N*-phenylhydrazinecarboxamide ligand and additionally coordinated by two Cl atoms, resulting in a distorted square-pyramidal geometry. The ligating atoms in the basal square plane of the complex comprise the azomethine N, the pyridine N, the amide O and one of the Cl atoms, whereas the other Cl atom occupies an apical position. The apical Cl atoms in adjacent layers function as hydrogen-bond acceptors to both NH groups. Intermolecular C–H···Cl and C–H···O interactions are also observed.

Related literature

For the biological applications of hydrazinecarboxamide and its derivatives, see: Beraldo & Gambino (2004); Kasuga *et al.* (2006); Rivadeneira *et al.* (2009); Shalini *et al.* (2009); Rodriguez-Arguelles *et al.* (2010). For the synthesis of related compounds, see: Kurup *et al.* (2011). For related structures, see: Kunnath *et al.* (2012). For the calculation of the trigonality index, see: Addison *et al.* (1984). For the graph-set notation, see: Etter *et al.* (1990).



Experimental

Crystal data

$[CuCl_2(C_{19}H_{16}N_4O)]$	$\gamma = 100.475 (1)^\circ$
$M_r = 450.81$	$V = 978.83 (7) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.4483 (5) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.8197 (3) \text{ \AA}$	$\mu = 1.41 \text{ mm}^{-1}$
$c = 11.5307 (4) \text{ \AA}$	$T = 293 \text{ K}$
$\alpha = 104.067 (1)^\circ$	$0.35 \times 0.32 \times 0.30 \text{ mm}$
$\beta = 103.026 (1)^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer	7149 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	4411 independent reflections
$T_{\min} = 0.614$, $T_{\max} = 0.649$	3550 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.092$	$\Delta\rho_{\max} = 0.34 \text{ e \AA}^{-3}$
$S = 1.01$	$\Delta\rho_{\min} = -0.40 \text{ e \AA}^{-3}$
4411 reflections	
252 parameters	
2 restraints	

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N3-H3'\cdots Cl^i$	0.85 (2)	2.40 (2)	3.1397 (18)	147 (2)
$N4-H4'\cdots Cl^i$	0.83 (2)	2.35 (2)	3.136 (2)	159 (2)
$C2-H2\cdots Cl^{ii}$	0.93	2.69	3.589 (4)	163
$C19-H19\cdots O1$	0.93	2.36	2.953 (4)	121

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 2, -z + 1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2/SAINT* (Bruker, 2004); data reduction: *SAINT/XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXL97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 2012) and *DIAMOND* (Brandenburg, 2010); software used to prepare material for publication: *SHELXL97* and *publCIF* (Westrip, 2010).

NA thanks the University Grants Commission (India) for a Junior Research Fellowship. We thank the Sophisticated Analytical Instruments Facility, Cochin University of S&T, for the diffraction measurements. MRPK thanks the University Grants Commission, New Delhi, for a UGC-BSR one-time grant to faculty. We also thank the Ministry of Higher Education of Malaysia (grant No. UM-C/HIR/MOHE/SC/12) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2644).

References

- Addison, A. W., Rao, J. N., Reedijk, J. & Verschoor, G. C. (1984). *Dalton Trans.* pp. 1349–1356.
- Beraldo, H. & Gambino, D. (2004). *Mini Rev. Med. Chem.* **4**, 31–39.
- Brandenburg, K. (2010). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *SADABS*, *APEX2*, *XPREP* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst. B* **46**, 256–262.

- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Kasuga, N. C., Onodera, K., Nakano, S., Hayashi, K. & Nomiya, K. (2006). *J. Inorg. Biochem.* **100**, 1176–1186.
- Kunnath, R. J., Prathapachandra Kurup, M. R. & Ng, S. W. (2012). *Acta Cryst. E* **68**, m1181.
- Kurup, M. R. P., Varghese, B., Sithambaresan, M., Krishnan, S., Sheeja, S. R. & Suresh, E. (2011). *Polyhedron*, **30**, 70–78.
- Rivadeneira, J., Barrio, D. A., Arrambide, G., Gambino, D., Bruzzone, L. & Etcheverry, S. B. (2009). *J. Inorg. Biochem.* **103**, 633–642.
- Rodriguez-Arguelles, M. C., Mosquera-Vazquez, S., Sanmartin-Matalobos, J., Garcia-Deibe, A. M., Pelizzetti, C. & Zani, F. (2010). *Polyhedron*, **29**, 864–870.
- Shalini, M., Yogeeshwari, P., Sriram, D. & Stables, J. P. (2009). *Biomed. Pharmacother.* **63**, 187–193.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2013). E69, m588–m589 [doi:10.1107/S1600536813026883]

Dichlorido{2-[*(E*)-phenyl(pyridin-2-yl- κ N)methylidene]-*N*-phenyl-hydrazinecarboxamide- κ^2 N²,O}copper(II)

N. Aiswarya, M. Sithambaresan, M. R. Prathapachandra Kurup and Seik Weng Ng

S1. Comment

Semicarbazones with versatile structural features are good ligands in both the neutral and anionic forms. Among the semicarbazones with diverse pharmacological activities (Beraldo & Gambino, 2004; Kasuga *et al.*, 2006; Rivadeneira *et al.*, 2009), the aryl semicarbazones were found to be devoid of sedative hypnotic activity and exhibited anticonvulsant activity with less neurotoxicity (Shalini *et al.*, 2009). The biological activity can be attributed to their ability to form chelates with transition metal ions by bonding *via* 'O' and azomethine 'N' atoms. Also, the chlorido complex of imidazole-2-carbaldehyde semicarbazone has been found to exhibit antimicrobial activity (Rodriguez-Arguelles *et al.*, 2010).

The title compound (Fig. 1) crystallizes in the triclinic space group $P\bar{1}$. The molecule adopts an *E* configuration with respect to C6=N2 bond and the tridentate ligand has its coordinating entities disposed in a *cis* fashion to each other.

The copper atom in the complex is *N,N',O* chelated by the neutral semicarbazone (Kunnath *et al.*, 2012). The C6=N2 [1.282 Å] and C13=O1 [1.229 Å] bond distances are very close to the formal C=N and C=O bond lengths [C=N; 1.28 Å and C=O; 1.21 Å] respectively confirming the azomethine bond formation and existence of semicarbazone in amido form. In addition to bond length and bond angle analysis, the trigonality index value confirms the coordination polyhedron to be a distorted square pyramidal (Addison *et al.*, 1984), with the apical chlorine atom out of the square plane by a distance of 2.450 Å. The apical chlorine atoms of the adjacent complex units function as hydrogen bond acceptors, generating a centrosymmetric dimer through a cyclic R_2^1 (6) association (Etter *et al.*, 1990). In addition to that a non-classical intermolecular C—H···Cl and an intramolecular C—H···O hydrogen bonds are also present in the molecular system (Fig. 2, Table 1).


Three types of Cu—Cl···Cg interactions are present in the complex (Fig. 3) with X···Cg distances of 3.9266 (16), 3.5545 (11) and 3.1361 (13) Å, respectively. C—H···π and N—H···π interactions are altogether absent in the molecule. Since the Cg—Cg distances are greater than 4 Å, the short ring interactions are not significant. Fig. 4 shows the packing diagram of the title compound along *c* axis.

S2. Experimental


The title compound was prepared by adapting a reported procedure (Kurup *et al.*, 2011). To the semicarbazone ligand synthesized by refluxing a mixture of hot methanolic solutions (25 ml) of 4-phenylsemicarbazide (0.151 g, 1 mmol) and 2-benzoylpyridine (0.183 g, 1 mmol), hot filtered methanolic solution (25 ml) of CuCl₂·2H₂O (0.170 g, 1 mmol) was added and refluxed for 2 h. The resulting green solution was cooled to room temperature. Green block shaped crystals were collected, washed with few drops of methanol and dried over P₄O₁₀ *in vacuo*. Single crystals suitable for X-ray analysis were obtained after slow evaporation of solution in air for few days. The compound was obtained in 75% yield (0.3375 g).

S3. Refinement

All H atoms on C were placed in calculated positions, guided by difference maps, with C—H bond distances of 0.93 Å. H atoms were assigned $U_{\text{iso}}(\text{H})$ values of 1.2 $U_{\text{eq}}(\text{carrier})$. Omitted owing to bad disagreement was reflection (0 0 1). H atoms of N3—H3' and N4—H4' bonds were located from difference maps and the bond distances are restrained to 0.88 ± 0.02 Å.

**Figure 1**

ORTEP view of the title compound, drawn with 50% probability displacement ellipsoids for the non-H atoms.

**Figure 2**

Graphical representation showing hydrogen-bonding interactions in the crystal structure of $C_{19}H_{16}Cl_2CuN_4O$.





Figure 3

Cu—Cl \cdots π interaction found in the title compound.

**Figure 4**

A view of the unit cell along c axis.

Dichlorido{2-[*(E*)-phenyl(pyridin-2-yl- κ N)methylidene]-*N*-phenylhydrazinecarboxamide- κ^2N^2,O }copper(II)

Crystal data

[CuCl₂(C₁₉H₁₆N₄O)]

$M_r = 450.81$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.4483 (5)$ Å

$b = 9.8197 (3)$ Å

$c = 11.5307 (4)$ Å

$\alpha = 104.067 (1)^\circ$

$\beta = 103.026 (1)^\circ$

$\gamma = 100.475 (1)^\circ$

$V = 978.83 (7)$ Å³

$Z = 2$

$F(000) = 458$

$D_x = 1.530$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3151 reflections

$\theta = 2.5\text{--}27.7^\circ$

$\mu = 1.41$ mm⁻¹

$T = 293$ K

Block, green

0.35 × 0.32 × 0.30 mm

Data collection

Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.33 pixels mm⁻¹

ω and φ scan

Absorption correction: multi-scan
(SADABS; Bruker, 2004)

$T_{\min} = 0.614$, $T_{\max} = 0.649$
 7149 measured reflections
 4411 independent reflections
 3550 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.3^\circ$
 $h = -9 \rightarrow 12$
 $k = -12 \rightarrow 12$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.092$
 $S = 1.01$
 4411 reflections
 252 parameters
 2 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0429P)^2 + 0.3399P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.40 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.56784 (3)	0.81865 (3)	0.45210 (3)	0.04217 (10)
Cl1	0.68970 (7)	0.63456 (5)	0.34892 (5)	0.04308 (14)
Cl2	0.54712 (10)	0.95006 (8)	0.32265 (7)	0.0657 (2)
O1	0.35179 (18)	0.68219 (17)	0.38879 (14)	0.0435 (4)
N1	0.7528 (2)	0.9491 (2)	0.5849 (2)	0.0474 (5)
N2	0.5556 (2)	0.74344 (17)	0.59382 (16)	0.0343 (4)
N3	0.4287 (2)	0.64128 (19)	0.57491 (17)	0.0385 (4)
H3'	0.418 (3)	0.593 (2)	0.625 (2)	0.048 (7)*
N4	0.2064 (2)	0.5040 (2)	0.43869 (19)	0.0459 (5)
C1	0.8558 (4)	1.0534 (3)	0.5712 (3)	0.0691 (9)
H1	0.8419	1.0733	0.4952	0.083*
C2	0.9811 (4)	1.1309 (4)	0.6673 (5)	0.0930 (12)
H2	1.0514	1.2026	0.6563	0.112*
C3	1.0019 (4)	1.1029 (4)	0.7775 (4)	0.0938 (12)
H3	1.0865	1.1555	0.8431	0.113*
C4	0.8967 (3)	0.9953 (3)	0.7934 (3)	0.0712 (8)
H4	0.9098	0.9743	0.8690	0.085*
C5	0.7732 (3)	0.9209 (2)	0.6946 (2)	0.0450 (5)
C6	0.6523 (2)	0.8054 (2)	0.7004 (2)	0.0358 (4)

C7	0.6428 (2)	0.7762 (2)	0.81791 (19)	0.0381 (5)
C8	0.6340 (3)	0.8871 (3)	0.9153 (2)	0.0500 (6)
H8	0.6375	0.9793	0.9066	0.060*
C9	0.6203 (3)	0.8607 (3)	1.0242 (2)	0.0596 (7)
H9	0.6128	0.9348	1.0884	0.071*
C10	0.6176 (3)	0.7256 (3)	1.0393 (2)	0.0587 (7)
H10	0.6090	0.7088	1.1137	0.070*
C11	0.6275 (3)	0.6155 (3)	0.9440 (2)	0.0564 (7)
H11	0.6263	0.5244	0.9544	0.068*
C12	0.6394 (3)	0.6395 (3)	0.8329 (2)	0.0458 (5)
H12	0.6450	0.5644	0.7684	0.055*
C13	0.3276 (2)	0.6125 (2)	0.46062 (19)	0.0364 (5)
C14	0.0774 (3)	0.4486 (3)	0.3355 (2)	0.0467 (5)
C15	-0.0277 (3)	0.3329 (3)	0.3403 (3)	0.0649 (8)
H15	-0.0097	0.2966	0.4081	0.078*
C16	-0.1589 (4)	0.2715 (4)	0.2447 (4)	0.0832 (10)
H16	-0.2297	0.1949	0.2488	0.100*
C17	-0.1846 (4)	0.3227 (5)	0.1446 (4)	0.0988 (13)
H17	-0.2719	0.2796	0.0793	0.119*
C18	-0.0819 (4)	0.4383 (5)	0.1398 (4)	0.0940 (12)
H18	-0.1011	0.4743	0.0719	0.113*
C19	0.0509 (3)	0.5020 (3)	0.2357 (3)	0.0664 (8)
H19	0.1206	0.5798	0.2319	0.080*
H4'	0.210 (3)	0.465 (3)	0.496 (2)	0.050 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.05420 (19)	0.03721 (16)	0.04289 (17)	0.00880 (12)	0.02232 (13)	0.01988 (12)
Cl1	0.0519 (3)	0.0338 (3)	0.0490 (3)	0.0081 (2)	0.0230 (3)	0.0161 (2)
Cl2	0.0992 (6)	0.0637 (4)	0.0735 (4)	0.0432 (4)	0.0528 (4)	0.0476 (4)
O1	0.0447 (9)	0.0477 (9)	0.0384 (8)	0.0059 (7)	0.0103 (7)	0.0190 (7)
N1	0.0500 (12)	0.0343 (9)	0.0629 (13)	0.0037 (9)	0.0314 (10)	0.0142 (9)
N2	0.0371 (10)	0.0284 (8)	0.0358 (9)	0.0010 (7)	0.0125 (8)	0.0099 (7)
N3	0.0438 (11)	0.0336 (9)	0.0337 (9)	-0.0034 (8)	0.0093 (8)	0.0135 (7)
N4	0.0431 (11)	0.0439 (11)	0.0445 (11)	-0.0019 (9)	0.0069 (9)	0.0167 (9)
C1	0.069 (2)	0.0500 (15)	0.102 (2)	0.0047 (14)	0.0523 (19)	0.0291 (15)
C2	0.057 (2)	0.063 (2)	0.156 (4)	-0.0107 (16)	0.044 (2)	0.033 (2)
C3	0.0433 (18)	0.080 (2)	0.129 (3)	-0.0197 (16)	0.004 (2)	0.024 (2)
C4	0.0431 (16)	0.0650 (18)	0.087 (2)	-0.0045 (13)	0.0021 (15)	0.0175 (16)
C5	0.0352 (12)	0.0374 (11)	0.0601 (15)	0.0044 (9)	0.0168 (11)	0.0107 (10)
C6	0.0362 (11)	0.0304 (10)	0.0405 (11)	0.0067 (8)	0.0133 (9)	0.0089 (8)
C7	0.0337 (11)	0.0398 (11)	0.0348 (11)	0.0055 (9)	0.0043 (9)	0.0081 (9)
C8	0.0583 (16)	0.0470 (13)	0.0391 (12)	0.0146 (12)	0.0068 (11)	0.0080 (10)
C9	0.0671 (19)	0.0687 (18)	0.0319 (12)	0.0138 (14)	0.0067 (12)	0.0041 (11)
C10	0.0535 (16)	0.081 (2)	0.0347 (12)	0.0047 (14)	0.0041 (11)	0.0220 (13)
C11	0.0546 (16)	0.0560 (15)	0.0559 (15)	0.0024 (12)	0.0057 (12)	0.0291 (13)
C12	0.0495 (14)	0.0413 (12)	0.0434 (12)	0.0069 (10)	0.0105 (11)	0.0124 (10)

C13	0.0398 (12)	0.0319 (10)	0.0367 (11)	0.0070 (9)	0.0122 (9)	0.0089 (8)
C14	0.0369 (13)	0.0430 (12)	0.0529 (14)	0.0084 (10)	0.0097 (11)	0.0049 (10)
C15	0.0482 (16)	0.0527 (15)	0.080 (2)	-0.0010 (13)	0.0132 (14)	0.0086 (14)
C16	0.0487 (18)	0.070 (2)	0.099 (3)	-0.0065 (15)	0.0032 (18)	-0.0016 (19)
C17	0.051 (2)	0.099 (3)	0.099 (3)	0.0014 (19)	-0.0168 (19)	-0.008 (2)
C18	0.072 (2)	0.114 (3)	0.077 (2)	0.022 (2)	-0.0126 (19)	0.026 (2)
C19	0.0487 (16)	0.0738 (19)	0.0654 (18)	0.0083 (14)	-0.0001 (13)	0.0205 (15)

Geometric parameters (\AA , $^{\circ}$)

Cu1—N2	1.9672 (16)	C6—C7	1.470 (3)
Cu1—N1	2.016 (2)	C7—C12	1.390 (3)
Cu1—O1	2.0865 (16)	C7—C8	1.391 (3)
Cu1—Cl2	2.1973 (6)	C8—C9	1.370 (4)
Cu1—Cl1	2.5175 (6)	C8—H8	0.9300
O1—C13	1.229 (3)	C9—C10	1.376 (4)
N1—C5	1.339 (3)	C9—H9	0.9300
N1—C1	1.344 (3)	C10—C11	1.375 (4)
N2—C6	1.282 (3)	C10—H10	0.9300
N2—N3	1.352 (2)	C11—C12	1.383 (3)
N3—C13	1.372 (3)	C11—H11	0.9300
N3—H3'	0.846 (16)	C12—H12	0.9300
N4—C13	1.343 (3)	C14—C19	1.369 (4)
N4—C14	1.410 (3)	C14—C15	1.388 (4)
N4—H4'	0.833 (16)	C15—C16	1.380 (4)
C1—C2	1.371 (5)	C15—H15	0.9300
C1—H1	0.9300	C16—C17	1.358 (5)
C2—C3	1.344 (6)	C16—H16	0.9300
C2—H2	0.9300	C17—C18	1.374 (6)
C3—C4	1.389 (4)	C17—H17	0.9300
C3—H3	0.9300	C18—C19	1.393 (4)
C4—C5	1.371 (4)	C18—H18	0.9300
C4—H4	0.9300	C19—H19	0.9300
C5—C6	1.483 (3)		
N2—Cu1—N1	78.70 (7)	C7—C6—C5	122.45 (19)
N2—Cu1—O1	77.84 (6)	C12—C7—C8	119.4 (2)
N1—Cu1—O1	154.03 (7)	C12—C7—C6	121.4 (2)
N2—Cu1—Cl2	162.21 (6)	C8—C7—C6	119.2 (2)
N1—Cu1—Cl2	99.08 (6)	C9—C8—C7	120.0 (2)
O1—Cu1—Cl2	99.86 (5)	C9—C8—H8	120.0
N2—Cu1—Cl1	96.69 (5)	C7—C8—H8	120.0
N1—Cu1—Cl1	97.98 (6)	C8—C9—C10	120.7 (2)
O1—Cu1—Cl1	95.64 (5)	C8—C9—H9	119.6
Cl2—Cu1—Cl1	101.09 (2)	C10—C9—H9	119.6
C13—O1—Cu1	112.00 (14)	C11—C10—C9	119.7 (2)
C5—N1—C1	119.2 (3)	C11—C10—H10	120.1
C5—N1—Cu1	114.21 (15)	C9—C10—H10	120.1

C1—N1—Cu1	126.5 (2)	C10—C11—C12	120.5 (2)
C6—N2—N3	123.84 (17)	C10—C11—H11	119.8
C6—N2—Cu1	120.05 (14)	C12—C11—H11	119.8
N3—N2—Cu1	115.29 (13)	C11—C12—C7	119.7 (2)
N2—N3—C13	113.53 (17)	C11—C12—H12	120.1
N2—N3—H3'	122.5 (18)	C7—C12—H12	120.1
C13—N3—H3'	123.5 (18)	O1—C13—N4	125.9 (2)
C13—N4—C14	129.7 (2)	O1—C13—N3	120.75 (19)
C13—N4—H4'	113.5 (19)	N4—C13—N3	113.39 (19)
C14—N4—H4'	116.8 (19)	C19—C14—C15	119.9 (3)
N1—C1—C2	121.2 (3)	C19—C14—N4	124.6 (2)
N1—C1—H1	119.4	C15—C14—N4	115.5 (2)
C2—C1—H1	119.4	C16—C15—C14	120.2 (3)
C3—C2—C1	119.8 (3)	C16—C15—H15	119.9
C3—C2—H2	120.1	C14—C15—H15	119.9
C1—C2—H2	120.1	C17—C16—C15	120.1 (3)
C2—C3—C4	119.9 (3)	C17—C16—H16	120.0
C2—C3—H3	120.1	C15—C16—H16	120.0
C4—C3—H3	120.1	C16—C17—C18	120.1 (3)
C5—C4—C3	118.2 (3)	C16—C17—H17	120.0
C5—C4—H4	120.9	C18—C17—H17	120.0
C3—C4—H4	120.9	C17—C18—C19	120.6 (4)
N1—C5—C4	121.8 (2)	C17—C18—H18	119.7
N1—C5—C6	114.9 (2)	C19—C18—H18	119.7
C4—C5—C6	123.3 (2)	C14—C19—C18	119.1 (3)
N2—C6—C7	125.61 (19)	C14—C19—H19	120.4
N2—C6—C5	111.80 (19)	C18—C19—H19	120.4
N2—Cu1—O1—C13	-6.89 (15)	N3—N2—C6—C5	-175.36 (19)
N1—Cu1—O1—C13	-32.7 (3)	Cu1—N2—C6—C5	-6.2 (2)
C12—Cu1—O1—C13	-168.91 (14)	N1—C5—C6—N2	6.1 (3)
C11—Cu1—O1—C13	88.77 (15)	C4—C5—C6—N2	-174.4 (2)
N2—Cu1—N1—C5	0.20 (16)	N1—C5—C6—C7	-169.7 (2)
O1—Cu1—N1—C5	25.9 (3)	C4—C5—C6—C7	9.7 (4)
C12—Cu1—N1—C5	162.26 (16)	N2—C6—C7—C12	61.0 (3)
C11—Cu1—N1—C5	-95.08 (16)	C5—C6—C7—C12	-123.7 (2)
N2—Cu1—N1—C1	177.3 (2)	N2—C6—C7—C8	-117.7 (3)
O1—Cu1—N1—C1	-157.0 (2)	C5—C6—C7—C8	57.6 (3)
C12—Cu1—N1—C1	-20.6 (2)	C12—C7—C8—C9	-0.8 (4)
C11—Cu1—N1—C1	82.0 (2)	C6—C7—C8—C9	177.9 (2)
N1—Cu1—N2—C6	3.66 (16)	C7—C8—C9—C10	1.1 (4)
O1—Cu1—N2—C6	-165.14 (18)	C8—C9—C10—C11	-0.5 (4)
C12—Cu1—N2—C6	-80.8 (3)	C9—C10—C11—C12	-0.4 (4)
C11—Cu1—N2—C6	100.51 (16)	C10—C11—C12—C7	0.7 (4)
N1—Cu1—N2—N3	173.68 (16)	C8—C7—C12—C11	-0.1 (4)
O1—Cu1—N2—N3	4.88 (14)	C6—C7—C12—C11	-178.8 (2)
C12—Cu1—N2—N3	89.2 (2)	Cu1—O1—C13—N4	-171.66 (19)
C11—Cu1—N2—N3	-89.48 (14)	Cu1—O1—C13—N3	8.0 (3)

C6—N2—N3—C13	167.1 (2)	C14—N4—C13—O1	−3.4 (4)
Cu1—N2—N3—C13	−2.5 (2)	C14—N4—C13—N3	176.9 (2)
C5—N1—C1—C2	0.1 (4)	N2—N3—C13—O1	−4.0 (3)
Cu1—N1—C1—C2	−176.9 (2)	N2—N3—C13—N4	175.67 (19)
N1—C1—C2—C3	−0.1 (6)	C13—N4—C14—C19	−1.9 (4)
C1—C2—C3—C4	0.2 (6)	C13—N4—C14—C15	178.9 (3)
C2—C3—C4—C5	−0.3 (6)	C19—C14—C15—C16	0.0 (5)
C1—N1—C5—C4	−0.2 (4)	N4—C14—C15—C16	179.3 (3)
Cu1—N1—C5—C4	177.2 (2)	C14—C15—C16—C17	1.0 (5)
C1—N1—C5—C6	179.3 (2)	C15—C16—C17—C18	−1.7 (6)
Cu1—N1—C5—C6	−3.4 (3)	C16—C17—C18—C19	1.4 (7)
C3—C4—C5—N1	0.3 (5)	C15—C14—C19—C18	−0.3 (5)
C3—C4—C5—C6	−179.1 (3)	N4—C14—C19—C18	−179.5 (3)
N3—N2—C6—C7	0.4 (3)	C17—C18—C19—C14	−0.4 (6)
Cu1—N2—C6—C7	169.48 (16)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3'···Cl1 ⁱ	0.85 (2)	2.40 (2)	3.1397 (18)	147 (2)
N4—H4'···Cl1 ⁱ	0.83 (2)	2.35 (2)	3.136 (2)	159 (2)
C2—H2···Cl1 ⁱⁱ	0.93	2.69	3.589 (4)	163
C19—H19···O1	0.93	2.36	2.953 (4)	121

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+2, -z+1$.