

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

Diaquabis(2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^2N^3,O$ )nickel(II) dihydrateAndrew P. Purdy<sup>a\*</sup> and Ray J. Butcher<sup>b</sup><sup>a</sup>Chemistry Division, Code 6100 Naval Research Laboratory, 4555 Overlook Avenue SW, Washington, DC 20375, USA, and <sup>b</sup>Department of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA

Correspondence e-mail: andrew.purdy@nrl.navy.mil


Received 13 October 2013; accepted 25 November 2013

Key indicators: single-crystal X-ray study;  $T = 123$  K; mean  $\sigma(C-C) = 0.004$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.112; data-to-parameter ratio = 14.1.

In the title complex,  $[Ni(C_6H_9N_2O_3S)_2(H_2O)_2] \cdot 2H_2O$ , the Ni<sup>II</sup> atom lies on an inversion center and is chelated by N and O atoms of two symmetry-equivalent imidazolesulfonate ligands in the basal plane, and two water O atoms in axial positions in an overall octahedral configuration. The crystal structure displays O—H...O and N—H...O hydrogen bonds, which connect the components into an extended three-dimensional network.

## Related literature

For examples of Ni–sulfonate complexes and MOFs, see: Lobana *et al.* (2004); Forbes & Sevov (2009); Kim *et al.* (2004); Yang *et al.* (2010). A small number of structurally characterized imidazole sulfonates are known, see: Kuhn *et al.* (2001, 2002); Chidambaram *et al.* (1988). The 2-ethyl-4-methyl-5-sulfonate ligand is described by Purdy *et al.* (2007) and Purdy & Butcher (2011).



## Experimental

## Crystal data

 $[Ni(C_6H_9N_2O_3S)_2(H_2O)_2] \cdot 2H_2O$   
 $M_r = 509.20$ Monoclinic,  $P2_1/a$  $a = 7.6037$  (2) Å $b = 16.8934$  (4) Å $c = 8.6574$  (3) Å $\beta = 111.303$  (3)° $V = 1036.08$  (5) Å<sup>3</sup> $Z = 2$ Cu  $K\alpha$  radiation $\mu = 3.77$  mm<sup>-1</sup> $T = 123$  K $0.52 \times 0.46 \times 0.35$  mm

## Data collection

Agilent Xcalibur Ruby Gemini diffractometer

Absorption correction: multi-scan

(CrysAlis PRO; Agilent, 2012)

 $T_{min} = 0.209$ ,  $T_{max} = 1.000$ 

2126 measured reflections

2126 independent reflections

2062 reflections with  $I > 2\sigma(I)$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$  $wR(F^2) = 0.112$  $S = 1.18$ 

2126 reflections

151 parameters

48 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{max} = 0.73$  e Å<sup>-3</sup> $\Delta\rho_{min} = -0.54$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W—H1W1...O2W	0.82 (2)	2.02 (2)	2.837 (3)	174 (4)
O1W—H1W2...O3 <sup>i</sup>	0.81 (2)	1.96 (2)	2.739 (3)	161 (4)
O2W—H2W1...O2 <sup>ii</sup>	0.80 (2)	1.97 (2)	2.751 (3)	167 (4)
O2W—H2W2...O2 <sup>iii</sup>	0.80 (2)	1.94 (2)	2.723 (3)	169 (4)
N2—H2A...O2W <sup>iv</sup>	0.88	1.94	2.818 (3)	174

Symmetry codes: (i)  $-x, -y + 1, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 1$ ; (iii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + 1$ ; (iv)  $-x + 1, -y + 1, -z + 2$ .

Data collection: CrysAlis PRO (Agilent, 2012); cell refinement: CrysAlis PRO; data reduction: CrysAlis PRO; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We thank The Office of Naval Research for financial support. RJB wishes to acknowledge the NSF–MRI program (grant CHE-0619278) for funds to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2177).

## References

- Agilent (2012). CrysAlis PRO and CrysAlis RED. Agilent Technologies, Yarnton, England.
- Chidambaram, S., Aravamudan, G. & Seshasayee, M. (1988). *Acta Cryst.* **C44**, 898–900.
- Forbes, T. Z. & Sevov, S. C. (2009). *Inorg. Chem.* **48**, 6873–6878.
- Kim, D. S., Forster, P. M., Le Toquin, R. & Cheetham, A. K. (2004). *Chem. Commun.* pp. 2148–2149.
- Kuhn, N., Eichele, K. & Walker, M. (2001). *Z. Anorg. Allg. Chem.* **627**, 2565–2567.

- Kuhn, N., Eichele, K., Walker, M., Berends, T. & Minkwitz, R. (2002). *Z. Anorg. Allg. Chem.* **628**, 2026–2032.
- Lobana, T. S., Kinoshita, I., Kimura, K., Nishioka, T., Shiomi, D. & Isobe, K. (2004). *Eur. J. Inorg. Chem.* pp. 356–367.
- Purdy, A. P. & Butcher, R. J. (2011). *Acta Cryst.* **E67**, m1303–m1304.
- Purdy, A. P., Gilardi, R., Luther, J. & Butcher, R. J. (2007). *Polyhedron*, **26**, 3930–3938.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yang, F., Wu, Z.-H. & Cai, J.-H. (2010). *Acta Cryst.* **E66**, m748.

## supporting information

*Acta Cryst.* (2014). E70, m18–m19 [https://doi.org/10.1107/S1600536813032169]

**Diaquabis(2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^2N^3,O$ )nickel(II) dihydrate****Andrew P. Purdy and Ray J. Butcher****S1. Comment**

The Ni atom lies on an inversion center and is chelated by N1 and O1 of the 2 symmetry equivalent imidazolesulfonate ligands in a plane, and 2 axial water molecules coordinated in an overall octahedral configuration. All of the Ni—O bond lengths (2.083 (2), 2.094 (2) Å) are about the same, and are in the normal range for octahedral coordinated Ni (2.05–2.12 Å). The largest deviation from the 90 ° angles of the octahedron is N1—Ni—O1, a result of the sulfonate group being attached to the imidazole ring. Two additional uncoordinated water molecules are present, but do not lie on any symmetry elements. There are no links other than hydrogen bonds between molecules, in contrast to the analogous unsolvated Cu(II) derivative (Purdy & Butcher, 2011). Hydrogen bonding links all the water molecules and N2, O2, and O3 into an extended structure in all three dimensions.

**S2. Experimental**

A solution of the potassium salt of the 2-ethyl-4-methyl-imidazole-5-sulfonic acid was prepared by combining 1 g (5.25 mmol) of the free acid with 1.5 equivalents of KOH solution, and diluting the solution to 1M based on K<sup>+</sup>. (All solutions were made with distilled water.) Two test reactions were done in vials with 0.5 M solutions of Ni(BF<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O and MnSO<sub>4</sub>·H<sub>2</sub>O, a 0.2 ml metered pipet was used for the additions, and each vial contained one addition of all three solutions. After 1 month, one vial were heated to a boil and allowed to cool and the other remained at room temperature. After one year, large blue crystals of the title compound grew in the solution that was not heated.


**Figure 1**

Diagram of  $C_{12}H_{22}N_4O_8S_2Ni$ , with uncoordinated water molecules not shown.




Figure 2

Packing diagram viewed down the *c* axis, displaying the hydrogen bonded interactions of both the coordinated and uncoordinated water molecules.

### Diaquabis(2-ethyl-5-methylimidazole-4-sulfonato- $\kappa^2N^3,O$ )nickel(II) dihydrate

#### Crystal data

$[\text{Ni}(\text{C}_6\text{H}_9\text{N}_2\text{O}_3\text{S})_2(\text{H}_2\text{O})_2] \cdot 2\text{H}_2\text{O}$

$M_r = 509.20$

Monoclinic,  $P2_1/a$

Hall symbol:  $-P\ 2yab$

$a = 7.6037\ (2)\ \text{\AA}$

$b = 16.8934\ (4)\ \text{\AA}$

$c = 8.6574\ (3)\ \text{\AA}$

$\beta = 111.303\ (3)^\circ$

$V = 1036.08\ (5)\ \text{\AA}^3$

$Z = 2$

$F(000) = 532$

$D_x = 1.632\ \text{Mg m}^{-3}$

Cu  $K\alpha$  radiation,  $\lambda = 1.54178\ \text{\AA}$

Cell parameters from 4614 reflections

$\theta = 5.2\text{--}75.5^\circ$

$\mu = 3.77\ \text{mm}^{-1}$

$T = 123\ \text{K}$

Block, blue

$0.52 \times 0.46 \times 0.35\ \text{mm}$

#### Data collection

Agilent Xcalibur Ruby Gemini  
diffractometer

Radiation source: Enhance (Cu) X-ray Source  
Graphite monochromator

Detector resolution:  $10.5081\ \text{pixels mm}^{-1}$

$\omega$  scans

Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.209$ ,  $T_{\max} = 1.000$   
 2126 measured reflections  
 2126 independent reflections  
 2062 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.000$

$\theta_{\max} = 75.8^\circ$ ,  $\theta_{\min} = 5.2^\circ$   
 $h = -9 \rightarrow 8$   
 $k = 0 \rightarrow 21$   
 $l = 0 \rightarrow 10$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.112$   
 $S = 1.18$   
 2126 reflections  
 151 parameters  
 48 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0411P)^2 + 2.2827P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.73 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.54 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni	0.5000	0.5000	0.5000	0.00455 (19)
S1	0.13784 (8)	0.60394 (4)	0.38692 (7)	0.00676 (18)
O1	0.2439 (3)	0.54424 (11)	0.3337 (2)	0.0082 (4)
O2	0.1485 (3)	0.68145 (12)	0.3149 (2)	0.0148 (4)
O3	-0.0536 (3)	0.58051 (13)	0.3598 (2)	0.0165 (4)
O1W	0.3581 (3)	0.40250 (11)	0.5452 (2)	0.0104 (4)
H1W1	0.422 (5)	0.378 (2)	0.629 (4)	0.027 (11)*
H1W2	0.267 (4)	0.417 (2)	0.563 (5)	0.027 (11)*
O2W	0.5754 (3)	0.30842 (12)	0.8196 (2)	0.0129 (4)
H2W1	0.667 (4)	0.309 (2)	0.795 (5)	0.014 (9)*
H2W2	0.519 (5)	0.2680 (16)	0.790 (5)	0.019 (10)*
N1	0.4281 (3)	0.56704 (13)	0.6667 (3)	0.0065 (4)
N2	0.3814 (3)	0.63709 (13)	0.8610 (3)	0.0094 (5)
H2A	0.3996	0.6567	0.9598	0.011*
C1	0.2651 (4)	0.61089 (15)	0.5997 (3)	0.0072 (5)
C2	0.2329 (4)	0.65519 (15)	0.7178 (3)	0.0085 (5)
C3	0.0834 (4)	0.71410 (17)	0.7081 (4)	0.0155 (6)
H3A	-0.0366	0.6972	0.6241	0.023*
H3B	0.1188	0.7659	0.6777	0.023*

H3C	0.0696	0.7179	0.8161	0.023*
C4	0.4953 (4)	0.58422 (15)	0.8257 (3)	0.0083 (5)
C5	0.6769 (4)	0.55417 (16)	0.9505 (3)	0.0110 (5)
H5A	0.6884	0.4968	0.9328	0.013*
H5B	0.6752	0.5615	1.0634	0.013*
C6	0.8474 (4)	0.59737 (17)	0.9371 (4)	0.0152 (6)
H6A	0.9637	0.5738	1.0147	0.023*
H6B	0.8420	0.6534	0.9641	0.023*
H6C	0.8457	0.5925	0.8237	0.023*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ni	0.0030 (3)	0.0058 (3)	0.0026 (3)	0.0012 (2)	-0.0017 (2)	-0.0004 (2)
S1	0.0039 (3)	0.0098 (3)	0.0042 (3)	0.0027 (2)	-0.0014 (2)	0.0007 (2)
O1	0.0067 (7)	0.0098 (7)	0.0051 (7)	0.0030 (6)	-0.0012 (5)	-0.0014 (6)
O2	0.0208 (11)	0.0099 (9)	0.0122 (9)	0.0059 (8)	0.0043 (8)	0.0056 (7)
O3	0.0112 (8)	0.0241 (8)	0.0131 (7)	-0.0003 (6)	0.0031 (6)	-0.0012 (6)
O1W	0.0087 (9)	0.0105 (9)	0.0108 (9)	0.0002 (7)	0.0023 (7)	0.0014 (7)
O2W	0.0164 (10)	0.0113 (9)	0.0105 (9)	-0.0051 (8)	0.0042 (8)	-0.0044 (7)
N1	0.0054 (8)	0.0070 (8)	0.0056 (7)	0.0000 (6)	0.0002 (6)	0.0002 (6)
N2	0.0121 (11)	0.0102 (11)	0.0050 (10)	-0.0014 (8)	0.0020 (8)	-0.0025 (8)
C1	0.0065 (8)	0.0075 (8)	0.0065 (8)	-0.0002 (7)	0.0008 (7)	-0.0005 (7)
C2	0.0078 (12)	0.0081 (11)	0.0085 (12)	-0.0012 (9)	0.0015 (10)	-0.0021 (9)
C3	0.0132 (14)	0.0129 (13)	0.0197 (14)	0.0028 (11)	0.0052 (11)	-0.0049 (11)
C4	0.0110 (13)	0.0059 (11)	0.0077 (12)	-0.0021 (9)	0.0029 (10)	-0.0003 (9)
C5	0.0115 (9)	0.0116 (9)	0.0081 (8)	0.0007 (7)	0.0013 (7)	0.0007 (7)
C6	0.0104 (13)	0.0177 (14)	0.0122 (13)	-0.0012 (10)	-0.0023 (10)	-0.0014 (10)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Ni—N1	2.058 (2)	N2—C4	1.354 (4)
Ni—N1 <sup>i</sup>	2.058 (2)	N2—C2	1.374 (3)
Ni—O1W	2.0825 (19)	N2—H2A	0.8800
Ni—O1W <sup>i</sup>	2.0825 (19)	C1—C2	1.359 (4)
Ni—O1	2.0941 (18)	C2—C3	1.490 (4)
Ni—O1 <sup>i</sup>	2.0941 (18)	C3—H3A	0.9800
S1—O3	1.443 (2)	C3—H3B	0.9800
S1—O2	1.465 (2)	C3—H3C	0.9800
S1—O1	1.4662 (18)	C4—C5	1.499 (4)
S1—C1	1.746 (3)	C5—C6	1.528 (4)
O1W—H1W1	0.823 (19)	C5—H5A	0.9900
O1W—H1W2	0.808 (19)	C5—H5B	0.9900
O2W—H2W1	0.798 (18)	C6—H6A	0.9800
O2W—H2W2	0.796 (18)	C6—H6B	0.9800
N1—C4	1.315 (3)	C6—H6C	0.9800
N1—C1	1.378 (3)		

N1—Ni—N1 <sup>i</sup>	180.0	C4—N2—H2A	125.5
N1—Ni—O1W	90.90 (8)	C2—N2—H2A	125.5
N1 <sup>i</sup> —Ni—O1W	89.10 (8)	C2—C1—N1	111.1 (2)
N1—Ni—O1W <sup>i</sup>	89.10 (8)	C2—C1—S1	130.4 (2)
N1 <sup>i</sup> —Ni—O1W <sup>i</sup>	90.90 (8)	N1—C1—S1	118.48 (19)
O1W—Ni—O1W <sup>i</sup>	180.00 (6)	C1—C2—N2	104.1 (2)
N1—Ni—O1	82.42 (8)	C1—C2—C3	131.9 (2)
N1 <sup>i</sup> —Ni—O1	97.58 (8)	N2—C2—C3	124.0 (2)
O1W—Ni—O1	89.75 (8)	C2—C3—H3A	109.5
O1W <sup>i</sup> —Ni—O1	90.25 (8)	C2—C3—H3B	109.5
N1—Ni—O1 <sup>i</sup>	97.58 (8)	H3A—C3—H3B	109.5
N1 <sup>i</sup> —Ni—O1 <sup>i</sup>	82.42 (8)	C2—C3—H3C	109.5
O1W—Ni—O1 <sup>i</sup>	90.25 (8)	H3A—C3—H3C	109.5
O1W <sup>i</sup> —Ni—O1 <sup>i</sup>	89.75 (8)	H3B—C3—H3C	109.5
O1—Ni—O1 <sup>i</sup>	180.00 (14)	N1—C4—N2	110.2 (2)
O3—S1—O2	112.61 (13)	N1—C4—C5	125.8 (2)
O3—S1—O1	113.45 (12)	N2—C4—C5	123.9 (2)
O2—S1—O1	111.06 (11)	C4—C5—C6	111.6 (2)
O3—S1—C1	109.24 (12)	C4—C5—H5A	109.3
O2—S1—C1	107.09 (12)	C6—C5—H5A	109.3
O1—S1—C1	102.73 (11)	C4—C5—H5B	109.3
S1—O1—Ni	120.66 (10)	C6—C5—H5B	109.3
Ni—O1W—H1W1	112 (3)	H5A—C5—H5B	108.0
Ni—O1W—H1W2	109 (3)	C5—C6—H6A	109.5
H1W1—O1W—H1W2	105 (3)	C5—C6—H6B	109.5
H2W1—O2W—H2W2	110 (3)	H6A—C6—H6B	109.5
C4—N1—C1	105.7 (2)	C5—C6—H6C	109.5
C4—N1—Ni	138.96 (19)	H6A—C6—H6C	109.5
C1—N1—Ni	115.34 (16)	H6B—C6—H6C	109.5
C4—N2—C2	109.0 (2)		
O3—S1—O1—Ni	-124.06 (13)	Ni—N1—C1—S1	1.0 (3)
O2—S1—O1—Ni	107.92 (14)	O3—S1—C1—C2	-57.4 (3)
C1—S1—O1—Ni	-6.27 (15)	O2—S1—C1—C2	64.8 (3)
N1—Ni—O1—S1	6.10 (13)	O1—S1—C1—C2	-178.1 (3)
N1 <sup>i</sup> —Ni—O1—S1	-173.90 (13)	O3—S1—C1—N1	124.0 (2)
O1W—Ni—O1—S1	97.04 (13)	O2—S1—C1—N1	-113.8 (2)
O1W <sup>i</sup> —Ni—O1—S1	-82.96 (13)	O1—S1—C1—N1	3.2 (2)
O1 <sup>i</sup> —Ni—O1—S1	89 (6)	N1—C1—C2—N2	-0.2 (3)
N1 <sup>i</sup> —Ni—N1—C4	81 (8)	S1—C1—C2—N2	-178.9 (2)
O1W—Ni—N1—C4	89.7 (3)	N1—C1—C2—C3	177.0 (3)
O1W <sup>i</sup> —Ni—N1—C4	-90.3 (3)	S1—C1—C2—C3	-1.7 (5)
O1—Ni—N1—C4	179.4 (3)	C4—N2—C2—C1	0.2 (3)
O1 <sup>i</sup> —Ni—N1—C4	-0.6 (3)	C4—N2—C2—C3	-177.3 (3)
N1 <sup>i</sup> —Ni—N1—C1	-102 (8)	C1—N1—C4—N2	-0.1 (3)
O1W—Ni—N1—C1	-93.13 (18)	Ni—N1—C4—N2	177.21 (19)
O1W <sup>i</sup> —Ni—N1—C1	86.87 (18)	C1—N1—C4—C5	-176.8 (2)
O1—Ni—N1—C1	-3.50 (17)	Ni—N1—C4—C5	0.5 (4)



O1 <sup>i</sup> —Ni—N1—C1	176.50 (17)	C2—N2—C4—N1	0.0 (3)
C4—N1—C1—C2	0.2 (3)	C2—N2—C4—C5	176.7 (2)
Ni—N1—C1—C2	-177.83 (17)	N1—C4—C5—C6	76.8 (3)
C4—N1—C1—S1	179.10 (18)	N2—C4—C5—C6	-99.4 (3)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .

*Hydrogen-bond geometry* ( $\text{\AA}$ ,  $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O1 <i>W</i> —H1 <i>W</i> 1 $\cdots$ O2 <i>W</i>	0.82 (2)	2.02 (2)	2.837 (3)	174 (4)
O1 <i>W</i> —H1 <i>W</i> 2 $\cdots$ O3 <sup>ii</sup>	0.81 (2)	1.96 (2)	2.739 (3)	161 (4)
O2 <i>W</i> —H2 <i>W</i> 1 $\cdots$ O2 <sup>i</sup>	0.80 (2)	1.97 (2)	2.751 (3)	167 (4)
O2 <i>W</i> —H2 <i>W</i> 1 $\cdots$ S1 <sup>i</sup>	0.80 (2)	2.92 (3)	3.602 (2)	145 (3)
O2 <i>W</i> —H2 <i>W</i> 2 $\cdots$ O2 <sup>iii</sup>	0.80 (2)	1.94 (2)	2.723 (3)	169 (4)
N2—H2 <i>A</i> $\cdots$ O2 <i>W</i> <sup>iv</sup>	0.88	1.94	2.818 (3)	174

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x, -y+1, -z+1$ ; (iii)  $-x+1/2, y-1/2, -z+1$ ; (iv)  $-x+1, -y+1, -z+2$ .