

9-[4-(Azidomethyl)phenyl]-9*H*-carbazole-3-carbonitrile

C. Ramathilagam,^a P. R. Umarani,^b N. Venkatesan,^c P. Rajakumar,^c B. Gunasekaran^{d*} and V. Manivannan^{e*}

^aDepartment of Physics, AMET University, Kanathur, Chennai 603 112, India,

^bPrincipal, Kundavai Nachiyar Govt College for Women, Thanjavur 613 007, India,

^cDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, ^dDepartment of Physics & Nano Technology, SRM University, SRM Nagar, Kattankulathur, Kancheepuram Dist., Chennai 603 203, Tamil Nadu, India, and ^eDepartment of Research and Development, PRIST University, Vallam, Thanjavur 613 403, Tamil Nadu, India

Correspondence e-mail: phdguna@gmail.com, crystallography2010@gmail.com


Received 16 January 2014; accepted 20 January 2014

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.065; wR factor = 0.230; data-to-parameter ratio = 17.3.

In the title compound $\text{C}_{20}\text{H}_{13}\text{N}_5$, the dihedral angle between the carbazole ring system (r.m.s. deviation = 0.027 Å) and the pendant benzene ring is $55.08(6)^\circ$. One of the azide N atoms is disordered over two positions in a 0.65 (2):0.35 (2) ratio. In the crystal, aromatic $\pi-\pi$ stacking is observed [minimum centroid–centroid separation = $3.6499(13)\text{ \AA}$] as well as inversion-dimers connected by pairs of weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the biological activity of carbazole derivatives, see: Ramsewak *et al.* (1999); Tachibana *et al.* (2001); Itoigawa *et al.* (2000); Friend *et al.* (1999). For related structures, see: Velmurugan *et al.* (2010); Ramathilagam *et al.* (2011)



Experimental

Crystal data



$M_r = 323.35$

Monoclinic, $P2_1/c$
 $a = 9.8457(2)\text{ \AA}$
 $b = 8.4032(2)\text{ \AA}$
 $c = 19.9127(5)\text{ \AA}$
 $\beta = 90.970(2)^\circ$
 $V = 1647.25(7)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.25 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.958$, $T_{\max} = 0.984$

15375 measured reflections
4076 independent reflections
2676 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.230$
 $S = 1.07$
4076 reflections
236 parameters

10 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.67\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$Cg3$ is the centroid of the C7–C12 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C14—H14··· $Cg3^i$	0.93	2.91	3.673 (2)	140
Symmetry code: (i) $-x + 1, -y + 1, -z$.				

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

CR acknowledge AMET University management, India, for their kind support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7185).

References

- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Friend, R. H., Gymer, R. W., Holmes, A. B., Burroughes, J. H., Mark, R. N., Taliani, C., Bradley, D. C., Dos Santos, D. A., Bredas, J. L., Logdlund, M. & Salaneck, W. R. (1999). *Nature (London)*, **397**, 121–127.
- Itoigawa, M., Kashiwada, Y., Ito, C., Furukawa, H., Tachibana, Y., Bastow, K. F. & Lee, K. H. (2000). *J. Nat. Prod.* **63**, 893–897.
- Ramathilagam, C., Venkatesan, N., Rajakumar, P., Umarani, P. R. & Manivannan, V. (2011). *Acta Cryst. E67*, o2796.
- Ramsewak, R. S., Nair, M. G., Strasburg, G. M., DeWitt, D. L. & Nitiss, J. L. (1999). *J. Agric. Food Chem.* **47**, 444–447.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Tachibana, Y., Kikuzaki, H., Lajis, N. H. & Nakatani, N. (2001). *J. Agric. Food Chem.* **49**, 5589–5594.
- Velmurugan, R., Sekar, M., Chandramohan, A., Ramesh, P. & Ponnuswamy, M. N. (2010). *Acta Cryst. E66*, o2965.

supporting information

Acta Cryst. (2014). E70, o197 [doi:10.1107/S1600536814001391]

9-[4-(Azidomethyl)phenyl]-9*H*-carbazole-3-carbonitrile

C. Ramathilagam, P. R. Umarani, N. Venkatesan, P. Rajakumar, B. Gunasekaran and V. Manivannan

S1. Comment

Carbazole derivatives possess antioxidative (Tachibana *et al.*, 2001), antitumor (Itoigawa *et al.*, 2000), anti-inflammatory and antimutagenic (Ramsewak *et al.*, 1999)activities. These derivatives also exhibit electroactivity and luminescence properties and are considered to be potential candidates for electronics such as colour display, organic semiconductor lasers and solar cells (Friend *et al.*, 1999).


The geometric parameters of the title molecule (Fig. 1) agree well with reported similar structure(Velmurugan *et al.*, 2010; Ramathilagam *et al.*, 2011). Phenyl ring makes the dihedral angle of 55.08 (6) ° with the carbazole ring system. In azidomethyl group one of the nitrogen atom is disordered over two position with site occupancies of 0.35 (2) and 0.65 (2). The sum of bond angles around the atom N1 [359.81 (17) °] indicates sp^2 hybridized. The crystal packing features weak C—H \cdots π [C14—H14 \cdots $Cg3(1 - x, 1 - y, -z)$ distance of 3.672 (7) Å, ($Cg3$ and is the centroid of the ring defined by the atoms C7—C12)] interactions.

S2. Experimental

9-(4-(bromomethyl)phenyl)-9*H*-carbazole-3-carbonitrile (1.0 mmol, 1.0 equiv) was dissolved in acetone/water (4:1, 8 ml). NaN₃ (1.5 mmol, 1.5 equiv.) was added and the mixture was heated at 60 °C for 5 h. The reaction mixture was cooled to room temperature, acetone was evaporated and the reaction mixture was diluted with H₂O (100 ml) and extracted with CHCl₃ (2 x 100 ml). The organic layer was washed with saturated NaCl (50 ml), dried over Na₂SO₄ and evaporated gave 9-(4-(azidomethyl)phenyl) -9*H*-carbazole-3-carbonitrile as pale yellow blocks with a yield of 84% and melting point 148 °C.

S3. Refinement

In azidomethyl group one of the nitrogen atom is disordered over two position. The site occupancy factors of disordered nitrogen atoms were refined as N3 = 0.35 (2) and N3A = 0.65 (2) during anisotropic refinement. The N3A—N4 bond distance was restrained to be 1.1500 (1) Å. The atom N3A exhibited elongation of the thermal ellipsoid and have been restrained (ISOR) to be more isotropic. The components of the anisotropic displacement parameters in direction of the bond N3—N4 and N4—N5, were restrained to be equal within an effective standard deviation of 0.001 using the DELU command in *SHELXL* (Sheldrick, 2008). H atoms were positioned geometrically and refined using riding model with C—H = 0.93 Å and $U_{iso}(\text{H})$ = 1.2Ueq(C) for aromatic C—H, C—H = 0.97 Å and $U_{iso}(\text{H})$ = 1.2Ueq(C) for CH₂.

**Figure 1**

The molecular structure of (I), with 30% probability displacement ellipsoids for non-H atoms.

9-[4-(Azidomethyl)phenyl]-9*H*-carbazole-3-carbonitrile

Crystal data

$C_{20}H_{13}N_5$
 $M_r = 323.35$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 9.8457 (2)$ Å
 $b = 8.4032 (2)$ Å
 $c = 19.9127 (5)$ Å
 $\beta = 90.970 (2)^\circ$
 $V = 1647.25 (7)$ Å³
 $Z = 4$

$F(000) = 672$
 $D_x = 1.304$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4076 reflections
 $\theta = 2.0\text{--}28.3^\circ$
 $\mu = 0.08$ mm⁻¹
 $T = 295$ K
Block, pale yellow
 $0.25 \times 0.20 \times 0.20$ mm

Data collection

Bruker kappa APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube

Graphite monochromator
Detector resolution: 0 pixels mm⁻¹
 ω and φ scans

Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.958$, $T_{\max} = 0.984$
 15375 measured reflections
 4076 independent reflections
 2676 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$
 $\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -12 \rightarrow 13$
 $k = -10 \rightarrow 11$
 $l = -26 \rightarrow 26$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.230$
 $S = 1.07$
 4076 reflections
 236 parameters
 10 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1219P)^2 + 0.4961P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.67 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.1427 (2)	0.6443 (3)	-0.02437 (10)	0.0462 (5)	
C2	0.0810 (2)	0.7908 (3)	-0.03579 (11)	0.0532 (5)	
H2	0.0891	0.8728	-0.0046	0.064*	
C3	0.0079 (3)	0.8108 (3)	-0.09424 (12)	0.0584 (6)	
H3	-0.0355	0.9073	-0.1025	0.070*	
C4	-0.0026 (2)	0.6891 (3)	-0.14171 (11)	0.0554 (6)	
C5	0.0600 (2)	0.5431 (3)	-0.13083 (10)	0.0502 (5)	
H5	0.0533	0.4627	-0.1628	0.060*	
C6	0.1326 (2)	0.5191 (2)	-0.07139 (10)	0.0453 (5)	
C7	0.2076 (2)	0.3867 (3)	-0.04382 (11)	0.0473 (5)	
C8	0.2302 (2)	0.2308 (3)	-0.06541 (12)	0.0541 (5)	
H8	0.1961	0.1962	-0.1067	0.065*	
C9	0.3039 (2)	0.1295 (3)	-0.02440 (13)	0.0602 (6)	
H9	0.3202	0.0258	-0.0384	0.072*	
C10	0.3538 (3)	0.1802 (3)	0.03750 (13)	0.0638 (6)	
H10	0.4019	0.1089	0.0646	0.077*	
C11	0.3343 (3)	0.3342 (3)	0.06016 (12)	0.0591 (6)	
H11	0.3687	0.3677	0.1016	0.071*	
C12	0.2609 (2)	0.4363 (3)	0.01842 (11)	0.0491 (5)	
C13	0.2663 (2)	0.6895 (2)	0.08503 (10)	0.0470 (5)	
C14	0.4037 (2)	0.7025 (3)	0.09936 (12)	0.0559 (6)	
H14	0.4666	0.6511	0.0726	0.067*	
C15	0.4469 (3)	0.7928 (3)	0.15393 (13)	0.0607 (6)	
H15	0.5393	0.8000	0.1640	0.073*	
C16	0.3556 (3)	0.8719 (3)	0.19349 (11)	0.0600 (6)	
C17	0.2184 (3)	0.8588 (3)	0.17812 (11)	0.0585 (6)	
H17	0.1556	0.9119	0.2044	0.070*	
C18	0.1733 (2)	0.7679 (3)	0.12418 (11)	0.0535 (5)	

H18	0.0809	0.7596	0.1144	0.064*
C19	-0.0782 (3)	0.7148 (3)	-0.20322 (13)	0.0720 (7)
C20	0.4028 (4)	0.9678 (4)	0.25333 (15)	0.0909 (10)
H20A	0.4985	0.9911	0.2479	0.109*
H20B	0.3548	1.0686	0.2523	0.109*
N1	0.22210 (19)	0.5936 (2)	0.02970 (9)	0.0501 (5)
N2	-0.1372 (4)	0.7353 (4)	-0.25201 (14)	0.1079 (10)
N3	0.4560 (12)	0.8595 (18)	0.3064 (5)	0.106 (4) 0.35 (2)
N3A	0.3867 (14)	0.9011 (9)	0.3169 (3)	0.134 (4) 0.65 (2)
N4	0.3659 (4)	0.7676 (7)	0.32487 (16)	0.1170 (13)
N5	0.3230 (6)	0.6426 (7)	0.3378 (3)	0.186 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0527 (11)	0.0444 (11)	0.0414 (10)	-0.0050 (9)	-0.0002 (8)	0.0002 (8)
C2	0.0671 (13)	0.0423 (12)	0.0502 (12)	-0.0001 (10)	-0.0010 (10)	-0.0016 (9)
C3	0.0714 (15)	0.0498 (13)	0.0538 (12)	0.0052 (11)	-0.0009 (11)	0.0063 (10)
C4	0.0656 (14)	0.0580 (14)	0.0425 (11)	-0.0009 (10)	-0.0018 (10)	0.0077 (9)
C5	0.0591 (12)	0.0503 (12)	0.0413 (10)	-0.0080 (9)	0.0015 (9)	-0.0012 (9)
C6	0.0499 (11)	0.0430 (11)	0.0431 (10)	-0.0062 (8)	0.0037 (8)	-0.0002 (8)
C7	0.0513 (11)	0.0444 (11)	0.0462 (11)	-0.0047 (8)	0.0020 (8)	-0.0014 (8)
C8	0.0584 (12)	0.0478 (12)	0.0561 (13)	-0.0058 (10)	0.0016 (10)	-0.0078 (10)
C9	0.0661 (14)	0.0428 (12)	0.0717 (15)	0.0011 (10)	0.0036 (12)	-0.0033 (10)
C10	0.0712 (15)	0.0511 (14)	0.0688 (15)	0.0082 (11)	-0.0046 (12)	0.0065 (11)
C11	0.0697 (14)	0.0531 (14)	0.0540 (13)	0.0034 (11)	-0.0085 (11)	0.0006 (10)
C12	0.0547 (11)	0.0431 (11)	0.0496 (11)	-0.0034 (9)	0.0000 (9)	-0.0013 (9)
C13	0.0587 (12)	0.0416 (11)	0.0405 (10)	-0.0067 (9)	-0.0041 (8)	-0.0005 (8)
C14	0.0579 (13)	0.0554 (13)	0.0543 (12)	-0.0036 (10)	0.0004 (10)	-0.0017 (10)
C15	0.0632 (14)	0.0579 (14)	0.0606 (13)	-0.0114 (11)	-0.0123 (11)	0.0004 (11)
C16	0.0848 (17)	0.0479 (13)	0.0468 (12)	-0.0102 (11)	-0.0128 (11)	-0.0002 (9)
C17	0.0772 (16)	0.0525 (13)	0.0458 (12)	0.0007 (11)	0.0028 (11)	-0.0061 (10)
C18	0.0587 (12)	0.0504 (13)	0.0512 (12)	-0.0028 (10)	-0.0015 (10)	-0.0019 (10)
C19	0.096 (2)	0.0680 (17)	0.0518 (14)	0.0063 (14)	-0.0078 (13)	0.0073 (12)
C20	0.123 (3)	0.083 (2)	0.0659 (18)	-0.0031 (18)	-0.0312 (17)	-0.0208 (15)
N1	0.0608 (11)	0.0419 (10)	0.0472 (10)	0.0001 (8)	-0.0076 (8)	-0.0047 (7)
N2	0.157 (3)	0.102 (2)	0.0629 (15)	0.021 (2)	-0.0350 (17)	0.0106 (15)
N3	0.074 (5)	0.200 (7)	0.044 (5)	0.028 (5)	-0.013 (3)	-0.012 (5)
N3A	0.157 (8)	0.172 (6)	0.073 (3)	-0.066 (5)	0.018 (4)	-0.039 (4)
N4	0.112 (3)	0.176 (4)	0.0636 (17)	0.046 (3)	0.0171 (17)	0.027 (2)
N5	0.194 (5)	0.181 (4)	0.187 (5)	0.079 (3)	0.087 (4)	0.105 (4)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.387 (3)	C12—N1	1.395 (3)
C1—C2	1.389 (3)	C13—C18	1.380 (3)
C1—C6	1.411 (3)	C13—C14	1.382 (3)
C2—C3	1.368 (3)	C13—N1	1.427 (3)

C2—H2	0.9300	C14—C15	1.386 (3)
C3—C4	1.396 (3)	C14—H14	0.9300
C3—H3	0.9300	C15—C16	1.377 (4)
C4—C5	1.389 (3)	C15—H15	0.9300
C4—C19	1.439 (3)	C16—C17	1.385 (4)
C5—C6	1.387 (3)	C16—C20	1.505 (3)
C5—H5	0.9300	C17—C18	1.385 (3)
C6—C7	1.439 (3)	C17—H17	0.9300
C7—C8	1.397 (3)	C18—H18	0.9300
C7—C12	1.401 (3)	C19—N2	1.137 (4)
C8—C9	1.378 (3)	C20—N3A	1.397 (8)
C8—H8	0.9300	C20—N3	1.483 (11)
C9—C10	1.387 (4)	C20—H20A	0.9700
C9—H9	0.9300	C20—H20B	0.9700
C10—C11	1.385 (4)	N3—N4	1.237 (14)
C10—H10	0.9300	N3A—N4	1.1513 (10)
C11—C12	1.389 (3)	N4—N5	1.163 (7)
C11—H11	0.9300		
N1—C1—C2	129.68 (19)	C18—C13—C14	120.1 (2)
N1—C1—C6	108.53 (18)	C18—C13—N1	120.60 (19)
C2—C1—C6	121.78 (19)	C14—C13—N1	119.3 (2)
C3—C2—C1	118.0 (2)	C13—C14—C15	119.5 (2)
C3—C2—H2	121.0	C13—C14—H14	120.3
C1—C2—H2	121.0	C15—C14—H14	120.3
C2—C3—C4	121.2 (2)	C16—C15—C14	121.2 (2)
C2—C3—H3	119.4	C16—C15—H15	119.4
C4—C3—H3	119.4	C14—C15—H15	119.4
C5—C4—C3	120.9 (2)	C15—C16—C17	118.6 (2)
C5—C4—C19	119.2 (2)	C15—C16—C20	121.1 (3)
C3—C4—C19	119.9 (2)	C17—C16—C20	120.3 (3)
C6—C5—C4	118.9 (2)	C16—C17—C18	120.9 (2)
C6—C5—H5	120.6	C16—C17—H17	119.5
C4—C5—H5	120.6	C18—C17—H17	119.5
C5—C6—C1	119.2 (2)	C13—C18—C17	119.6 (2)
C5—C6—C7	133.7 (2)	C13—C18—H18	120.2
C1—C6—C7	107.10 (18)	C17—C18—H18	120.2
C8—C7—C12	119.5 (2)	N2—C19—C4	179.6 (4)
C8—C7—C6	133.7 (2)	N3A—C20—C16	117.7 (3)
C12—C7—C6	106.80 (19)	N3—C20—C16	109.6 (6)
C9—C8—C7	118.8 (2)	N3A—C20—H20A	107.9
C9—C8—H8	120.6	N3—C20—H20A	82.6
C7—C8—H8	120.6	C16—C20—H20A	107.9
C8—C9—C10	120.8 (2)	N3A—C20—H20B	107.9
C8—C9—H9	119.6	N3—C20—H20B	135.8
C10—C9—H9	119.6	C16—C20—H20B	107.9
C11—C10—C9	121.8 (2)	H20A—C20—H20B	107.2
C11—C10—H10	119.1	C1—N1—C12	108.59 (17)

C9—C10—H10	119.1	C1—N1—C13	125.89 (18)
C10—C11—C12	117.1 (2)	C12—N1—C13	125.34 (18)
C10—C11—H11	121.4	N4—N3—C20	110.4 (7)
C12—C11—H11	121.4	N4—N3A—C20	122.6 (6)
C11—C12—N1	129.1 (2)	N3A—N4—N5	167.6 (8)
C11—C12—C7	121.9 (2)	N5—N4—N3	153.5 (8)
N1—C12—C7	108.97 (19)		
N1—C1—C2—C3	179.5 (2)	C14—C15—C16—C17	0.5 (4)
C6—C1—C2—C3	0.7 (3)	C14—C15—C16—C20	179.1 (2)
C1—C2—C3—C4	-1.1 (4)	C15—C16—C17—C18	0.1 (4)
C2—C3—C4—C5	0.5 (4)	C20—C16—C17—C18	-178.4 (2)
C2—C3—C4—C19	-178.9 (2)	C14—C13—C18—C17	-0.4 (3)
C3—C4—C5—C6	0.6 (3)	N1—C13—C18—C17	179.3 (2)
C19—C4—C5—C6	-180.0 (2)	C16—C17—C18—C13	-0.2 (3)
C4—C5—C6—C1	-1.0 (3)	C15—C16—C20—N3A	-103.0 (8)
C4—C5—C6—C7	179.6 (2)	C17—C16—C20—N3A	75.6 (8)
N1—C1—C6—C5	-178.63 (18)	C15—C16—C20—N3	-69.1 (6)
C2—C1—C6—C5	0.4 (3)	C17—C16—C20—N3	109.5 (6)
N1—C1—C6—C7	0.9 (2)	C2—C1—N1—C12	180.0 (2)
C2—C1—C6—C7	179.95 (19)	C6—C1—N1—C12	-1.1 (2)
C5—C6—C7—C8	-3.3 (4)	C2—C1—N1—C13	-4.8 (4)
C1—C6—C7—C8	177.3 (2)	C6—C1—N1—C13	174.15 (19)
C5—C6—C7—C12	179.1 (2)	C11—C12—N1—C1	-176.2 (2)
C1—C6—C7—C12	-0.4 (2)	C7—C12—N1—C1	0.9 (2)
C12—C7—C8—C9	0.5 (3)	C11—C12—N1—C13	8.5 (4)
C6—C7—C8—C9	-176.9 (2)	C7—C12—N1—C13	-174.42 (19)
C7—C8—C9—C10	0.6 (4)	C18—C13—N1—C1	57.2 (3)
C8—C9—C10—C11	-1.1 (4)	C14—C13—N1—C1	-123.1 (2)
C9—C10—C11—C12	0.6 (4)	C18—C13—N1—C12	-128.3 (2)
C10—C11—C12—N1	177.2 (2)	C14—C13—N1—C12	51.3 (3)
C10—C11—C12—C7	0.5 (4)	N3A—C20—N3—N4	50.2 (7)
C8—C7—C12—C11	-1.0 (3)	C16—C20—N3—N4	-61.0 (11)
C6—C7—C12—C11	177.0 (2)	N3—C20—N3A—N4	-66.7 (13)
C8—C7—C12—N1	-178.34 (19)	C16—C20—N3A—N4	16.2 (17)
C6—C7—C12—N1	-0.3 (2)	C20—N3A—N4—N5	-118 (3)
C18—C13—C14—C15	1.0 (3)	C20—N3A—N4—N3	68.5 (11)
N1—C13—C14—C15	-178.6 (2)	C20—N3—N4—N3A	-51.9 (9)
C13—C14—C15—C16	-1.1 (4)	C20—N3—N4—N5	131.1 (12)

Hydrogen-bond geometry (Å, °)

Cg3 is the centroid of the C7—C12 ring.

D—H···A	D—H	H···A	D···A	D—H···A
C14—H14···Cg3 ⁱ	0.93	2.91	3.673 (2)	140

Symmetry code: (i) $-x+1, -y+1, -z$.