

(R)-N-[(R)-2,2-Dichloro-1-phenyl-2-(phenylsulfonyl)ethyl]-2-methylpropane-2-sulfonamide

Haiji Huang, Ya Li* and Jingming Chu

College of Chemistry and Chemical Engineering, Shanghai University of Engineering Science, 333 Longteng Road, Shanghai, People's Republic of China
Correspondence e-mail: ya.li@sues.edu.cn


Received 9 December 2013; accepted 30 December 2013

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.038; wR factor = 0.086; data-to-parameter ratio = 17.2.

The title molecule, $\text{C}_{18}\text{H}_{21}\text{Cl}_2\text{NO}_3\text{S}_2$, contains one chiral carbon center and the absolute stereochemistry has been confirmed as *R*. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond occurs and the dihedral angle between the benzene rings is $64.5(1)^\circ$. In the crystal, the molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a zigzag chain structure extending along the *c*-axis direction.

Related literature

For the use of β -amino sulfones as enzyme inhibitors, see: Tamamura *et al.* (2003); Nakatani *et al.* (2008); Raja *et al.* (2009). For their use as synthetic intermediates, see: Pauly *et al.* (1994); de Blas *et al.* (1994); Carretero *et al.* (1997); Alonso *et al.* (1997). For their synthesis, see: Zhang *et al.* (2011). For the fluorinated analogue, see: Li & Hu (2005); Liu & Hu (2010).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{21}\text{Cl}_2\text{NO}_3\text{S}_2$

$M_r = 434.38$

Orthorhombic, $P2_12_12_1$

$a = 7.924(2)\text{ \AA}$

$b = 14.772(4)\text{ \AA}$

$c = 18.151(5)\text{ \AA}$

$V = 2124.6(10)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.52\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.39 \times 0.30 \times 0.26\text{ mm}$

Data collection

Bruker CCD area-detector

diffractometer

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2004)

$T_{\min} = 0.743$, $T_{\max} = 1.000$

11625 measured reflections

4174 independent reflections

3834 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.086$

$S = 0.99$

4174 reflections

242 parameters

H atoms treated by a mixture of
independent and constrained
refinement

$\Delta\rho_{\text{max}} = 0.28\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.20\text{ e \AA}^{-3}$

Absolute structure: Flack, 1983;

1786 Friedel pairs

Absolute structure parameter:

-0.04 (6)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C17—H17 \cdots O1 ⁱ	0.93	2.44	3.236 (4)	144
N1—H1 \cdots O2	0.81 (3)	2.19 (3)	2.889 (3)	145 (3)

Symmetry code: (i) $-x - \frac{1}{2}, -y, z - \frac{1}{2}$

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The Innovation Program of Shanghai Municipal Education Commission (12YZ155) and the Innovation Program of National University Students (201310856014) are gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2283).

References

- Alonso, D. A., Costa, A., Mancheno, B. & Najera, C. (1997). *Tetrahedron*, **53**, 4791–4814.
- Blas, J. de, Carretero, J. C. & Dominguez, E. (1994). *Tetrahedron Lett.* **35**, 4603–4606.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Carretero, J. C., Arrayas, R. G. & Storch de Garcia, J. (1997). *Tetrahedron Lett.* **38**, 8537–8540.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Li, Y. & Hu, J. (2005). *Angew. Chem. Int. Ed.* **44**, 5882–5886.
- Liu, J. & Hu, J. (2010). *Chem.-Eur. J.* **16**, 11443–11454.
- Nakatani, S., Hidaka, K., Ami, E., Nakahara, K., Sato, A., Nguyen, J.-T., Hamada, Y., Hori, Y., Ohnishi, N., Nagai, A., Kimura, T., Hayashi, Y. & Kiso, Y. (2008). *J. Med. Chem.* **51**, 2992–3004.
- Pauly, R., Sasaki, A. & Portier, P. (1994). *Tetrahedron Lett.* **35**, 237–240.
- Raja, S. N., Surber, B. W., Du, J. & Cross, J. L. (2009). *J. Labelled Compd. Radiopharm.* **52**, 98–102.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tamamura, H., Koh, Y., Ueda, S., Sasaki, Y., Yamasaki, T., Aoki, M., Maeda, K., Watai, Y., Arikuni, Y., Otaka, A., Mitsuya, H. & Fujii, N. (2003). *J. Med. Chem.* **46**, 1764–1768.
- Zhang, H., Li, Y., Xu, W., Zheng, W., Zhou, P. & Sun, Z. (2011). *Org. Biomol. Chem.* **9**, 6502–6505.

supporting information

Acta Cryst. (2014). E70, o119 [doi:10.1107/S1600536813034909]

(*R*)-*N*-[(*R*)-2,2-Dichloro-1-phenyl-2-(phenylsulfonyl)ethyl]-2-methylpropane-2-sulfinamide

Haiji Huang, Ya Li and Jingming Chu

S1. Comment

The β -amino sulfone motif is of significant importance because it was found to play important roles in a variety of bioactive compounds (Tamamura *et al.*, 2003; Nakatani *et al.*, 2008). Besides, β -amino sulfones are valuable intermediates in the preparation of a wide variety of interesting compounds (Pauly *et al.*, 1994; Zhang *et al.*, 2011). Herein, we report the crystal structure of a α,α -dichloro- β -amino sulfone, the title compound, $C_{18}H_{21}Cl_2NO_3S_2$, which may find potential use in the synthesis of bioactive compounds.


In this compound (Fig. 1), the benzene rings *A* (C3–C8) and *B* (C13–C18) are planar with r. m. s. deviations of 0.0053 Å and 0.0050 Å, respectively. The dihedral angle between *A* and *B* is 64.5 (1) $^\circ$. An intramolecular N1—H···O2 hydrogen-bonding interaction is present while a weak intermolecular C17—H···O1ⁱ hydrogen bond (Table 1, Figs. 2,3) results in a one-dimensional zigzag chain structure extending along *c*. Present also are 40 Å³ solvent accessible voids. The absolute configuration at the chiral centre [C2(*R*)] has been confirmed [absolute structure parameter = -0.04 (6) for 1786 Friedel pairs] (Flack, 1983).

S2. Experimental


Sodium bis(trimethylsilyl)amide (1.2 mmol, 1M in THF) was added slowly to a reaction mixture of (*Rs*)-*N*-benzylidene-2-methylpropane-2-sulfinamide (209 mg, 1.0 mmol) and (dichloromethylsulfonyl)benzene (225 mg, 1.0 mmol) in THF at 203 K. The mixture was stirred for 1 h at this temperature and then quenched with 1M HCl. The quenched mixture was extracted with EtOAc. After removal of the solvent, the residue was subjected to flash column chromatography to give the title compound (226 mg: yield, 52%). The obtained compound was recrystallized from ethyl acetate/hexane (1:1) to give colorless crystals.

S3. Refinement


All the H atoms of the phenyl groups were placed at calculated positions and treated as riding on the parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The methyl H atoms were placed at calculated positions and allowed to rotate, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. The H atoms of the amide group were located in a difference-Fourier map and refined isotropically. The absolute structure parameter [-0.04 (6); Flack, 1983] at the chiral center (C2) has been confirmed as *R* (1786 Friedel pairs).

**Figure 1**

Molecular structure of the title compound. The displacement ellipsoids are drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii.

**Figure 2**

Intramolecular N—H···O and intermolecular C—H···O interactions in the title compound, shown as dashed lines. For symmetry code *a*: $-x - 1/2, -y, z - 1/2$.

**Figure 3**

The zigzag chain formed via intermolecular C—H···O interactions, viewed along the *b* axis. The intramolecular hydrogen bonds are omitted for clarity.

(*R*)-*N*-[(*R*)-2,2-Dichloro-1-phenyl-2-(phenylsulfonyl)ethyl]-2-methylpropane-2-sulfonamide

Crystal data



$M_r = 434.38$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.924 (2)$ Å

$b = 14.772 (4)$ Å

$c = 18.151 (5)$ Å

$V = 2124.6 (10)$ Å³

$Z = 4$

$F(000) = 904$

$D_x = 1.358 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4807 reflections

$\theta = 5.3\text{--}51.2^\circ$

$\mu = 0.52 \text{ mm}^{-1}$

$T = 293$ K

Prismatic, colorless

$0.39 \times 0.30 \times 0.26$ mm

Data collection

Bruker CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2004)

$T_{\min} = 0.743$, $T_{\max} = 1.000$

11625 measured reflections

4174 independent reflections

3834 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.075$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.8^\circ$

$h = -9 \rightarrow 9$

$k = -18 \rightarrow 12$

$l = -22 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.086$

$S = 0.99$

4174 reflections

242 parameters

0 restraints

Primary atom site location: structure-invariant direct methods	$w = 1/[\sigma^2(F_o^2) + (0.0379P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
Secondary atom site location: difference Fourier map	$(\Delta/\sigma)_{\max} < 0.001$
Hydrogen site location: inferred from neighbouring sites	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$
H atoms treated by a mixture of independent and constrained refinement	Absolute structure: Flack, 1983: 1786 Friedel pairs Absolute structure parameter: -0.04 (6)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.31897 (9)	0.05580 (5)	0.41327 (3)	0.04581 (18)
S2	-0.02571 (8)	0.04757 (4)	0.21496 (3)	0.03700 (15)
C11	0.30446 (8)	-0.00843 (5)	0.15767 (3)	0.04829 (18)
C12	0.05808 (9)	-0.14312 (4)	0.19785 (4)	0.05009 (18)
N1	0.2849 (3)	0.04579 (15)	0.32217 (10)	0.0382 (5)
O1	0.1580 (3)	0.0660 (2)	0.45195 (11)	0.0862 (8)
O2	0.0522 (2)	0.13428 (12)	0.22302 (11)	0.0505 (5)
O3	-0.1488 (2)	0.01835 (13)	0.26710 (9)	0.0501 (5)
C1	0.1468 (3)	-0.03735 (16)	0.22190 (13)	0.0356 (5)
C2	0.2123 (3)	-0.04217 (15)	0.30187 (12)	0.0350 (5)
H2	0.1156	-0.0537	0.3342	0.042*
C3	0.3408 (3)	-0.11663 (16)	0.31436 (12)	0.0350 (5)
C4	0.5097 (3)	-0.10324 (17)	0.29887 (13)	0.0392 (6)
H4	0.5458	-0.0477	0.2805	0.047*
C5	0.6241 (4)	-0.17113 (19)	0.31038 (14)	0.0481 (7)
H5	0.7373	-0.1616	0.2993	0.058*
C6	0.5737 (4)	-0.2531 (2)	0.33810 (17)	0.0584 (8)
H6	0.6522	-0.2990	0.3456	0.070*
C7	0.4087 (5)	-0.2670 (2)	0.3544 (2)	0.0658 (9)
H7	0.3742	-0.3225	0.3732	0.079*
C8	0.2916 (4)	-0.19900 (19)	0.34330 (16)	0.0534 (7)
H8	0.1790	-0.2088	0.3554	0.064*
C9	0.4166 (4)	0.16808 (18)	0.41012 (14)	0.0459 (7)
C10	0.5815 (4)	0.1612 (2)	0.3691 (2)	0.0754 (10)
H10A	0.5600	0.1451	0.3188	0.113*
H10B	0.6509	0.1157	0.3916	0.113*
H10C	0.6386	0.2185	0.3708	0.113*
C11	0.4464 (5)	0.1909 (2)	0.49142 (18)	0.0764 (10)

H11A	0.5033	0.2481	0.4951	0.115*
H11B	0.5146	0.1446	0.5136	0.115*
H11C	0.3399	0.1943	0.5165	0.115*
C12	0.2961 (5)	0.2359 (2)	0.37614 (17)	0.0702 (10)
H12A	0.3346	0.2962	0.3865	0.105*
H12B	0.1855	0.2276	0.3966	0.105*
H12C	0.2920	0.2270	0.3238	0.105*
C13	-0.1098 (3)	0.04079 (18)	0.12566 (13)	0.0391 (6)
C14	-0.0407 (4)	0.0949 (2)	0.07116 (15)	0.0514 (7)
H14	0.0539	0.1305	0.0803	0.062*
C15	-0.1168 (4)	0.0943 (3)	0.00274 (17)	0.0673 (9)
H15	-0.0724	0.1292	-0.0352	0.081*
C16	-0.2563 (4)	0.0429 (2)	-0.00949 (16)	0.0656 (9)
H16	-0.3057	0.0432	-0.0560	0.079*
C17	-0.3265 (4)	-0.0098 (2)	0.04547 (17)	0.0581 (8)
H17	-0.4227	-0.0441	0.0364	0.070*
C18	-0.2513 (3)	-0.01063 (19)	0.11397 (15)	0.0463 (6)
H18	-0.2961	-0.0456	0.1518	0.056*
H1	0.237 (3)	0.0892 (18)	0.3050 (14)	0.041 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0602 (4)	0.0465 (4)	0.0308 (3)	-0.0052 (3)	-0.0030 (3)	0.0027 (3)
S2	0.0334 (3)	0.0386 (3)	0.0389 (3)	0.0012 (3)	-0.0006 (2)	-0.0038 (3)
Cl1	0.0409 (3)	0.0641 (4)	0.0399 (3)	0.0035 (3)	0.0099 (3)	0.0063 (3)
Cl2	0.0535 (4)	0.0381 (3)	0.0587 (4)	-0.0048 (3)	-0.0079 (3)	-0.0113 (3)
N1	0.0503 (13)	0.0327 (11)	0.0316 (10)	0.0010 (11)	-0.0046 (9)	0.0017 (9)
O1	0.0792 (15)	0.126 (2)	0.0538 (12)	-0.0324 (16)	0.0265 (12)	-0.0116 (13)
O2	0.0522 (11)	0.0372 (10)	0.0621 (12)	-0.0006 (9)	-0.0135 (10)	-0.0033 (8)
O3	0.0396 (9)	0.0694 (13)	0.0412 (9)	0.0064 (9)	0.0068 (8)	-0.0025 (9)
C1	0.0339 (12)	0.0355 (13)	0.0375 (12)	-0.0035 (10)	0.0007 (10)	-0.0024 (10)
C2	0.0373 (12)	0.0343 (12)	0.0334 (11)	-0.0039 (11)	0.0056 (10)	0.0023 (10)
C3	0.0373 (13)	0.0325 (13)	0.0352 (12)	-0.0022 (10)	-0.0024 (10)	-0.0008 (10)
C4	0.0414 (14)	0.0402 (14)	0.0360 (13)	0.0010 (11)	0.0054 (11)	0.0002 (11)
C5	0.0435 (15)	0.0576 (18)	0.0433 (14)	0.0084 (13)	0.0014 (12)	-0.0091 (13)
C6	0.063 (2)	0.0442 (17)	0.0677 (19)	0.0158 (15)	-0.0101 (16)	-0.0060 (15)
C7	0.073 (2)	0.0365 (16)	0.088 (2)	-0.0026 (15)	-0.0091 (19)	0.0145 (15)
C8	0.0458 (16)	0.0450 (16)	0.0693 (18)	-0.0078 (14)	-0.0022 (15)	0.0114 (14)
C9	0.0569 (17)	0.0404 (15)	0.0403 (14)	0.0007 (13)	-0.0041 (12)	-0.0078 (11)
C10	0.064 (2)	0.077 (2)	0.085 (2)	-0.024 (2)	0.0075 (18)	-0.0120 (19)
C11	0.103 (3)	0.071 (2)	0.0547 (18)	0.001 (2)	-0.025 (2)	-0.0192 (17)
C12	0.112 (3)	0.0453 (17)	0.0533 (17)	0.0220 (19)	-0.0169 (19)	-0.0043 (14)
C13	0.0370 (13)	0.0429 (14)	0.0373 (12)	0.0088 (12)	-0.0010 (10)	-0.0008 (11)
C14	0.0466 (15)	0.0557 (17)	0.0520 (17)	0.0031 (15)	0.0024 (13)	0.0057 (13)
C15	0.071 (2)	0.087 (2)	0.0443 (17)	0.017 (2)	0.0021 (16)	0.0115 (16)
C16	0.073 (2)	0.080 (2)	0.0440 (15)	0.022 (2)	-0.0168 (15)	-0.0116 (17)
C17	0.0516 (16)	0.0634 (19)	0.0594 (17)	0.0085 (16)	-0.0146 (15)	-0.0173 (15)

C18	0.0428 (14)	0.0472 (15)	0.0489 (15)	0.0021 (13)	-0.0045 (12)	-0.0064 (12)
-----	-------------	-------------	-------------	-------------	--------------	--------------

Geometric parameters (\AA , $\text{^{\circ}}$)

S1—O1	1.464 (2)	C8—H8	0.9300
S1—N1	1.682 (2)	C9—C10	1.507 (4)
S1—C9	1.831 (3)	C9—C12	1.516 (4)
S2—O3	1.4258 (18)	C9—C11	1.532 (4)
S2—O2	1.4296 (19)	C10—H10A	0.9600
S2—C13	1.755 (2)	C10—H10B	0.9600
S2—C1	1.860 (3)	C10—H10C	0.9600
C11—C1	1.761 (2)	C11—H11A	0.9600
C12—C1	1.768 (2)	C11—H11B	0.9600
N1—C2	1.468 (3)	C11—H11C	0.9600
N1—H1	0.81 (3)	C12—H12A	0.9600
C1—C2	1.543 (3)	C12—H12B	0.9600
C2—C3	1.516 (3)	C12—H12C	0.9600
C2—H2	0.9800	C13—C18	1.371 (4)
C3—C8	1.381 (4)	C13—C14	1.385 (4)
C3—C4	1.382 (3)	C14—C15	1.381 (4)
C4—C5	1.368 (4)	C14—H14	0.9300
C4—H4	0.9300	C15—C16	1.359 (5)
C5—C6	1.371 (4)	C15—H15	0.9300
C5—H5	0.9300	C16—C17	1.382 (5)
C6—C7	1.356 (5)	C16—H16	0.9300
C6—H6	0.9300	C17—C18	1.379 (4)
C7—C8	1.382 (4)	C17—H17	0.9300
C7—H7	0.9300	C18—H18	0.9300
O1—S1—N1	109.91 (13)	C10—C9—C12	112.9 (3)
O1—S1—C9	106.88 (15)	C10—C9—C11	110.9 (3)
N1—S1—C9	96.71 (11)	C12—C9—C11	110.1 (3)
O3—S2—O2	119.92 (12)	C10—C9—S1	108.7 (2)
O3—S2—C13	109.64 (12)	C12—C9—S1	110.2 (2)
O2—S2—C13	108.03 (12)	C11—C9—S1	103.6 (2)
O3—S2—C1	104.70 (11)	C9—C10—H10A	109.5
O2—S2—C1	106.24 (11)	C9—C10—H10B	109.5
C13—S2—C1	107.65 (11)	H10A—C10—H10B	109.5
C2—N1—S1	112.80 (16)	C9—C10—H10C	109.5
C2—N1—H1	115.1 (19)	H10A—C10—H10C	109.5
S1—N1—H1	112.7 (19)	H10B—C10—H10C	109.5
C2—C1—Cl1	113.30 (16)	C9—C11—H11A	109.5
C2—C1—Cl2	108.97 (16)	C9—C11—H11B	109.5
Cl1—C1—Cl2	109.45 (13)	H11A—C11—H11B	109.5
C2—C1—S2	110.00 (16)	C9—C11—H11C	109.5
Cl1—C1—S2	108.24 (12)	H11A—C11—H11C	109.5
Cl2—C1—S2	106.66 (12)	H11B—C11—H11C	109.5
N1—C2—C3	110.0 (2)	C9—C12—H12A	109.5

N1—C2—C1	109.09 (18)	C9—C12—H12B	109.5
C3—C2—C1	113.58 (19)	H12A—C12—H12B	109.5
N1—C2—H2	108.0	C9—C12—H12C	109.5
C3—C2—H2	108.0	H12A—C12—H12C	109.5
C1—C2—H2	108.0	H12B—C12—H12C	109.5
C8—C3—C4	118.5 (2)	C18—C13—C14	122.2 (2)
C8—C3—C2	120.5 (2)	C18—C13—S2	119.0 (2)
C4—C3—C2	121.1 (2)	C14—C13—S2	118.5 (2)
C5—C4—C3	120.4 (2)	C15—C14—C13	117.7 (3)
C5—C4—H4	119.8	C15—C14—H14	121.1
C3—C4—H4	119.8	C13—C14—H14	121.1
C4—C5—C6	120.7 (3)	C16—C15—C14	120.4 (3)
C4—C5—H5	119.6	C16—C15—H15	119.8
C6—C5—H5	119.6	C14—C15—H15	119.8
C7—C6—C5	119.6 (3)	C15—C16—C17	121.6 (3)
C7—C6—H6	120.2	C15—C16—H16	119.2
C5—C6—H6	120.2	C17—C16—H16	119.2
C6—C7—C8	120.4 (3)	C18—C17—C16	118.8 (3)
C6—C7—H7	119.8	C18—C17—H17	120.6
C8—C7—H7	119.8	C16—C17—H17	120.6
C3—C8—C7	120.4 (3)	C13—C18—C17	119.2 (3)
C3—C8—H8	119.8	C13—C18—H18	120.4
C7—C8—H8	119.8	C17—C18—H18	120.4
O1—S1—N1—C2	-72.4 (2)	C4—C5—C6—C7	-0.2 (4)
C9—S1—N1—C2	176.92 (18)	C5—C6—C7—C8	0.0 (5)
O3—S2—C1—C2	-56.98 (18)	C4—C3—C8—C7	-1.8 (4)
O2—S2—C1—C2	70.87 (18)	C2—C3—C8—C7	179.8 (3)
C13—S2—C1—C2	-173.60 (16)	C6—C7—C8—C3	1.0 (5)
O3—S2—C1—Cl1	178.74 (12)	O1—S1—C9—C10	-178.7 (2)
O2—S2—C1—Cl1	-53.41 (15)	N1—S1—C9—C10	-65.5 (2)
C13—S2—C1—Cl1	62.12 (15)	O1—S1—C9—C12	-54.5 (2)
O3—S2—C1—Cl2	61.06 (14)	N1—S1—C9—C12	58.8 (2)
O2—S2—C1—Cl2	-171.09 (12)	O1—S1—C9—C11	63.3 (2)
C13—S2—C1—Cl2	-55.56 (15)	N1—S1—C9—C11	176.5 (2)
S1—N1—C2—C3	-68.1 (2)	O3—S2—C13—C18	-17.1 (2)
S1—N1—C2—C1	166.75 (16)	O2—S2—C13—C18	-149.4 (2)
Cl1—C1—C2—N1	58.3 (2)	C1—S2—C13—C18	96.3 (2)
Cl2—C1—C2—N1	-179.58 (16)	O3—S2—C13—C14	156.5 (2)
S2—C1—C2—N1	-63.0 (2)	O2—S2—C13—C14	24.2 (2)
Cl1—C1—C2—C3	-64.7 (2)	C1—S2—C13—C14	-90.2 (2)
Cl2—C1—C2—C3	57.4 (2)	C18—C13—C14—C15	-1.6 (4)
S2—C1—C2—C3	173.99 (16)	S2—C13—C14—C15	-174.9 (2)
N1—C2—C3—C8	140.1 (2)	C13—C14—C15—C16	0.9 (4)
C1—C2—C3—C8	-97.3 (3)	C14—C15—C16—C17	0.2 (5)
N1—C2—C3—C4	-38.3 (3)	C15—C16—C17—C18	-0.7 (5)
C1—C2—C3—C4	84.2 (3)	C14—C13—C18—C17	1.0 (4)
C8—C3—C4—C5	1.6 (4)	S2—C13—C18—C17	174.3 (2)

C2—C3—C4—C5	−179.9 (2)	C16—C17—C18—C13	0.1 (4)
C3—C4—C5—C6	−0.7 (4)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C17—H17···O1 ⁱ	0.93	2.44	3.236 (4)	144
N1—H1···O2	0.81 (3)	2.19 (3)	2.889 (3)	145 (3)
N1—H1···S2	0.81 (3)	2.72 (3)	3.138 (2)	114 (2)

Symmetry code: (i) $-x-1/2, -y, z-1/2$.