

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Redetermination of 2-methyl-4-nitropyridine *N*-oxide

Max Peukert, Wilhelm Seichter and Edwin Weber*

Institut für Organische Chemie, TU Bergakademie Freiberg, Leipziger Strasse 29,
D-09596 Freiberg/Sachsen, Germany

Correspondence e-mail: edwin.weber@chemie.tu-freiberg.de


Received 22 January 2014; accepted 26 February 2014

Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å;
 R factor = 0.049; wR factor = 0.147; data-to-parameter ratio = 14.9.

An improved crystal structure of the title compound, $\text{C}_6\text{H}_6\text{N}_2\text{O}_3$, is reported. The structure, previously solved [Li *et al.* (1987). *Jiegou Huaxue (Chin. J. Struct. Chem.)*, **6**, 20–24] in the orthorhombic space group $Pca2_1$ and refined to $R = 0.067$, has been solved in the orthorhombic space group $Pbcm$ with data of enhanced quality, giving an improved structure ($R = 0.0485$). The molecule adopts a planar conformation with all atoms lying on a mirror plane. The crystal structure is composed of molecular sheets extending parallel to the ab plane and connected *via* $\text{C}-\text{H}\cdots\text{O}$ contacts involving ring H atoms and O atoms of the *N*-oxide and nitro groups, while van der Waals forces consolidate the stacking of the layers.

Related literature

For the synthesis and preparative aspects of pyridine-*N*-oxides, see: Fontenas *et al.* (1995); Katritzky & Lagowski (1971); Kilenyi (2001); Mosher *et al.* (1963). For the preparation of the title compound, see: Ashimori *et al.* (1990) and for potential applications, see: Elemans *et al.* (2009); Weber & Vögtle (1976); Winter *et al.* (2004). For the previous report of its crystal structure, see: Li *et al.* (1987). For non-classical hydrogen bonds, see: Desiraju & Steiner (1999).



Experimental

Crystal data

 $\text{C}_6\text{H}_6\text{N}_2\text{O}_3$ $M_r = 154.13$ Orthorhombic, $Pbcm$ $a = 8.6775$ (7) Å $b = 12.4069$ (10) Å $c = 6.1995$ (5) Å $V = 667.44$ (9) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.13$ mm⁻¹ $T = 153$ K $0.57 \times 0.30 \times 0.23$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

 $T_{\min} = 0.932$, $T_{\max} = 0.972$

19832 measured reflections

1100 independent reflections

973 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.028$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.049$ $wR(F^2) = 0.147$ $S = 1.10$

1100 reflections

74 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C2}-\text{H2A}\cdots\text{O1}^i$	0.95	2.29	3.225 (2)	169
$\text{C5}-\text{H5}\cdots\text{O2}^{ii}$	0.95	2.36	3.301 (2)	173

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT-NT (Bruker, 2008); data reduction: SAINT-NT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: SHELXL97.

The authors thank the German Research Foundation within the priority programme Porous Metal-Organic Frameworks (SPP 1362, MOFs).

Supporting information for this paper is available from the IUCr electronic archives (Reference: ZP2011).

References

- Ashimori, A., Ono, T., Uchida, T., Ohtaki, Y., Fukaja, C., Watanabe, M. & Yokoyama, K. (1990). *Chem. Pharm. Bull.* **38**, 2446–2458.
- Bruker (2008). *APEX2*, *SAINTE-NT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond In Structural Chemistry and Biology*, ch. 2. Oxford University Press.
- Elemans, J. A. A. W., Lei, S. & De Feyter, S. (2009). *Angew. Chem. Int. Ed.* **48**, 7298–7332.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Fontenas, C., Bejan, E., Ait Haddou, H. & Balavoine, G. G. A. (1995). *Synth. Commun.* **25**, 629–633.
- Katritzky, A. R. & Lagowski, J. M. (1971). In *Chemistry of the Heterocyclic N-Oxides*. New York: Academic Press.
- Kilenyi, S. N. (2001). In *Encyclopedia of Reagents for Organic Synthesis*, edited by L. A. Paquette. New York: Wiley.
- Li, S., Liu, S. & Wu, W. (1987). *Jiegou Huaxue (Chin. J. Struct. Chem.)*, **6**, 20–24.
- Mosher, H. S., Turner, L. & Carlsmith, A. (1963). *Org. Synth.*, Coll. Vol. 4, 828–830.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Weber, E. & Vögtle, F. (1976). *Chem. Ber.* **109**, 1803–1831.
- Winter, S., Seichter, W. & Weber, E. (2004). *J. Coord. Chem.* **57**, 991–1014.

supporting information

Acta Cryst. (2014). E70, o426–o427 [doi:10.1107/S1600536814004450]

Redetermination of 2-methyl-4-nitropyridine *N*-oxide

Max Peukert, Wilhelm Seichter and Edwin Weber

S1. Comment


Pyridine *N*-oxides are readily formed by oxidation of corresponding pyridines (Kilenyi, 2001; Mosher *et al.*, 1963). In contrast to the simple pyridines they facilitate an electrophilic substitution reaction in the ring position-4, hence being important intermediates in the synthesis of pyridine derivatives featuring a complex substitution pattern (Katritzky & Lagowski, 1971). Moreover, when 2-methylpyridine *N*-oxides are treated with trifluoroacetic anhydride, the Boekelheide reaction occurs to give 2-(hydroxymethyl)pyridines (Fontenas *et al.*, 1995) which are of relevance to make available chelating (Winter *et al.*, 2004), macrocyclic (Weber & Vögtle, 1976) and linker-type (Elemans *et al.*, 2009) ligands. In the course of a respective synthesis of the latter kind, the title compound was prepared and its structure redetermined. The previous crystal structure of the compound (reported in 1987 by Li *et al.*) has been solved in the orthorhombic space group *Pca*2₁ and refined to an *R*-value of 6.7%. The repeated analysis of the crystal structure with data of enhanced quality yields a crystal structure of space group *Pbcm* with nearly identical cell dimensions. The centrosymmetry of the crystal structure is sustained by the statistical analysis of *E*-values. The molecule is located on the crystallographic symmetry plane and thus adopts perfect planarity (Fig. 1). According to this, two-dimensional supramolecular aggregates extending parallel to the crystallographic *ab*-plane and with the molecules connected *via* C—H \cdots O hydrogen bonding (Desiraju & Steiner, 1999) that involves ring H atoms and both O atoms of the *N*-oxide (C—H \cdots O_{*N*-oxide} 2.29 Å, 169 °) and nitro groups (C—H \cdots O_{*nitro*} 2.36 Å, 173 °) represent the basic entities of the crystal structure (Fig. 2). As no other type of intermolecular interactions are observed, the crystal structure is stabilized by van der Waals forces in direction of the stacking axes of the molecular sheets.

S2. Experimental


The title compound was synthesized *via* nitration of 2-methylpyridine *N*-oxide following a described procedure (Ashimori *et al.*, 1990). Crystallization from toluene/chloroform (1/1) yielded yellow needles which were used for X-ray single-crystal structure analysis.

S3. Refinement

Aromatic H atoms were positioned geometrically and allowed to ride on their parent atoms, with C—H = 0.95 Å and $U_{\text{iso}} = 1.2 U_{\text{eq}}(\text{C})$.

**Figure 1**

Perspective view of the molecular structure of the title compound including the atom numbering. Anisotropic displacement parameters for non-hydrogen atoms are drawn at a 50% probability level.

**Figure 2**

Packing diagram of the title compound viewed down the *c*-axis. Hydrogen bonds are displayed as broken lines.

2-Methyl-4-nitropyridine *N*-oxide

Crystal data

$C_6H_6N_2O_3$

$M_r = 154.13$

Orthorhombic, *Pbcm*

Hall symbol: -P 2c 2b

$a = 8.6775$ (7) Å

$b = 12.4069$ (10) Å

$c = 6.1995$ (5) Å

$V = 667.44$ (9) Å³

$Z = 4$

$F(000) = 320$

$D_x = 1.534$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6590 reflections

$\theta = 2.4\text{--}35.0^\circ$

$\mu = 0.13$ mm⁻¹

$T = 153$ K

Column, yellow

$0.57 \times 0.30 \times 0.23$ mm

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.932$, $T_{\max} = 0.972$

19832 measured reflections

1100 independent reflections

973 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 30.4^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -12 \rightarrow 12$

$k = -17 \rightarrow 17$

$l = -8 \rightarrow 8$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.147$

$S = 1.10$

1100 reflections

74 parameters

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0871P)^2 + 0.2298P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. The C—H bonds of the methyl group were restrained to a target value of 0.89 (1) Å.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.52895 (16)	0.30510 (10)	0.2500	0.0309 (3)
O2	1.05609 (15)	-0.02519 (12)	0.2500	0.0355 (4)
O3	0.85937 (17)	-0.13544 (11)	0.2500	0.0441 (4)
N1	0.61843 (17)	0.22205 (10)	0.2500	0.0223 (3)
N2	0.91523 (17)	-0.04406 (12)	0.2500	0.0276 (3)
C1	0.55535 (18)	0.12002 (13)	0.2500	0.0216 (3)
C2	0.65356 (17)	0.03153 (12)	0.2500	0.0203 (3)
H2A	0.6128	-0.0395	0.2500	0.024*
C3	0.81182 (18)	0.04811 (12)	0.2500	0.0214 (3)
C4	0.87504 (19)	0.15129 (13)	0.2500	0.0249 (3)
H4	0.9834	0.1621	0.2500	0.030*
C5	0.77482 (19)	0.23649 (13)	0.2500	0.0247 (3)
H5	0.8150	0.3077	0.2500	0.030*
C6	0.38480 (19)	0.11425 (16)	0.2500	0.0283 (4)
H6A	0.3468 (19)	0.1469 (13)	0.133 (2)	0.040 (5)*
H6B	0.358 (3)	0.0444 (9)	0.2500	0.035 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0389 (7)	0.0217 (6)	0.0322 (7)	0.0136 (5)	0.000	0.000
O2	0.0209 (6)	0.0365 (7)	0.0492 (8)	0.0072 (5)	0.000	0.000
O3	0.0376 (8)	0.0197 (6)	0.0750 (12)	0.0050 (5)	0.000	0.000
N1	0.0277 (7)	0.0182 (6)	0.0210 (6)	0.0033 (5)	0.000	0.000
N2	0.0254 (6)	0.0243 (7)	0.0329 (7)	0.0049 (5)	0.000	0.000
C1	0.0232 (6)	0.0210 (7)	0.0206 (7)	0.0009 (5)	0.000	0.000
C2	0.0211 (7)	0.0170 (6)	0.0228 (7)	-0.0018 (5)	0.000	0.000
C3	0.0214 (7)	0.0182 (6)	0.0247 (7)	0.0023 (5)	0.000	0.000

C4	0.0272 (7)	0.0215 (7)	0.0259 (7)	-0.0047 (6)	0.000	0.000
C5	0.0286 (7)	0.0211 (7)	0.0244 (7)	-0.0050 (6)	0.000	0.000
C6	0.0202 (7)	0.0350 (9)	0.0296 (8)	0.0013 (6)	0.000	0.000

Geometric parameters (Å, °)

O1—N1	1.2902 (17)	C2—C3	1.389 (2)
O2—N2	1.244 (2)	C2—H2A	0.9500
O3—N2	1.233 (2)	C3—C4	1.393 (2)
N1—C5	1.369 (2)	C4—C5	1.369 (2)
N1—C1	1.379 (2)	C4—H4	0.9500
N2—C3	1.454 (2)	C5—H5	0.9500
C1—C2	1.390 (2)	C6—H6A	0.892 (9)
C1—C6	1.482 (2)	C6—H6B	0.898 (10)
O1—N1—C5	119.48 (14)	C2—C3—C4	121.72 (14)
O1—N1—C1	119.61 (14)	C2—C3—N2	119.61 (14)
C5—N1—C1	120.91 (13)	C4—C3—N2	118.68 (14)
O3—N2—O2	123.99 (15)	C5—C4—C3	117.36 (15)
O3—N2—C3	118.73 (14)	C5—C4—H4	121.3
O2—N2—C3	117.28 (14)	C3—C4—H4	121.3
N1—C1—C2	118.79 (14)	N1—C5—C4	121.92 (14)
N1—C1—C6	116.16 (14)	N1—C5—H5	119.0
C2—C1—C6	125.05 (15)	C4—C5—H5	119.0
C3—C2—C1	119.30 (14)	C1—C6—H6A	110.3 (11)
C3—C2—H2A	120.3	C1—C6—H6B	108.0 (16)
C1—C2—H2A	120.3	H6A—C6—H6B	109.9 (14)
O1—N1—C1—C2	180.0	O2—N2—C3—C2	180.0
C5—N1—C1—C2	0.0	O3—N2—C3—C4	180.0
O1—N1—C1—C6	0.0	O2—N2—C3—C4	0.0
C5—N1—C1—C6	180.0	C2—C3—C4—C5	0.0
N1—C1—C2—C3	0.0	N2—C3—C4—C5	180.0
C6—C1—C2—C3	180.0	O1—N1—C5—C4	180.0
C1—C2—C3—C4	0.0	C1—N1—C5—C4	0.0
C1—C2—C3—N2	180.0	C3—C4—C5—N1	0.0
O3—N2—C3—C2	0.0		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2A \cdots O1 ⁱ	0.95	2.29	3.225 (2)	169
C5—H5 \cdots O2 ⁱⁱ	0.95	2.36	3.301 (2)	173

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $-x+2, y+1/2, -z+1/2$.